

Yang-Mills Mass Gap Clay Institute Problem Solution

Cusack P*

Independent Researcher, Canada

Abstract

Here is the solution to the Yang-Mills Mass Gap Problem as described below. The Mass in the material universe is a balance between the resistance to mass formation (cuz=Pi-base e), squeezing Energy, and Gravity pulling it apart. There is an explanation as to why there are 116 sable elements in the material universe. The calculations are verified by predicting the size of the universe within experimental error.

Keywords: Quantum physics; Elementary particle theory; Yang-Mills theory

Introduction

The laws of quantum physics stand to the world of elementary particles in the way that Newton's laws of classical mechanics stand to the macroscopic world. Almost half a century ago, Yang and Mills introduced a remarkable new framework to describe elementary particles using structures that also occur in geometry. Quantum Yang-Mills theory is now the foundation of most of elementary particle theory, and its predictions have been tested at many experimental laboratories, but its mathematical foundation is still unclear. The successful use of Yang-Mills theory to describe the strong interactions of elementary particles depends on a subtle quantum mechanical property called the "mass gap": the quantum particles have positive masses, even though the classical waves travel at the speed of light. This property has been discovered by physicists from experiment and confirmed by computer simulations, but it still has not been understood from a theoretical point of view. Progress in establishing the existence of the Yang-Mills theory and a mass gap will require the introduction of fundamental new ideas both in physics and in mathematics.

Literature Search

As for a literature search on this topic, there are no Mathematical Physics models that I know about that assume a static universe in compression. All the theories purpose incorrectly that the universe is expanding. In order to understand the Mass Gap problem, we need to understand that there is a force on the exterior of the universe that is compressing the "dust" or "clay particles" together. Mass precipices. The only place you find this theory is in my 6 volume work entitled, Astrotheology, The Missing Link, and Cusack's Model of the Universe [1,2].

Explanation

To begin, the Total Energy in the universe follows the e^{-t} , because the universe exist where the functioun equals the derivative ($y=y'$ =telescopic) function while the Kinetic Energy follows the Ln function. The difference is the Potential Energy stored in Mass, called Rm and is equal to $cuz=Pi-e=0.4233$ (Figure 1). In my work Astrotheology, The Missing Link, I've calculated that Gravity= Pi/Ln 1.618. When $t=Pi$ or the human frequency is $1/Pi=31.8$ Hz, then humans perceive the Energy being squeezed into a Mass signal. Time is determined to follow the Golden Mean Equation. $t^2-t-1=0$, roots, $t=0.618, 1.618$. I call this the Energy parabola [3]. Space is formed by the cross product of E and t is:

$$s=|E||t| \sin \theta$$




Figure 1: Illustration of resistance to mass.

$$s=F$$

$$F=\sin \theta$$

We know in a harmonic system, the Cairnaut Second Order Linear Ordinary Differential Equation applies to the wave energy So,

$$\partial E^2 / \partial t - E = 0$$

$$I've shown G=E^1$$

Gravity is that irreducible value that occurs when,

$$y=y'$$

$$y=E \text{ (Figure 2).}$$

So the volume of this half cube is,

$$\frac{1}{2} * 1 * 1 * 4 / 3 = 6.66 = G$$

When we subtract the weak nuclear force of of $G=6.67$, we get,

*Corresponding author: Cusack P, Independent Researcher, Canada, Tel: (506) 214-3313; E-mail: St-Michael@Hotmail.Com

Received February 08, 2016; Accepted April 29, 2016; Published May 05, 2016

Citation: Cusack P (2016) Yang-Mills Mass Gap Clay Institute Problem Solution. J Phys Math 7: 171. doi:10.4172/2090-0902.1000171

Copyright: © 2016 Cusack P. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 4: Illustration of the universe inside and out.




Figure 5: Illustration 5 plate with a hole.

right to cause space, and mass to precipitate. The force acting on the inner universe is $\sin\theta$ [4].

$$E = 1/cuz, \mu = 0.27, \rho = 0.127, F = \pi - 2a$$

$$u/u_0 = f(r/a) \sin(\omega t - \phi)$$

$$u/u_0 = f(a/a) \sin(1 - 2a/\pi)$$

$$u/u_0 = f(1) \sin(\pi/\pi - 2a/\pi)$$

$$= \sin 1 = s = v = a \text{ (Figure 5)}$$

The periodic wave that travels in the outer core or cylinder of the tubular universe, reaches the right force ($F = \sin \theta = 8/3$) at the outer edge of the ellipsoid thereby squeezing the Energy by the right amount to precipitate mass and space and time.

$$V = 2\pi R^2 [(1 - \pi/4)r - R/6]$$

Substituting, $t = 1.0337, 60^\circ = 1/6^{\text{th}}$ circle

$$B = R [(5/6 - \pi/4) - R/24] / (1 - \pi/4) = 1.4467 - 4/3/6 = 0.2084$$

Finally we achieve

$$\sigma = E\varepsilon$$

$$F(\varepsilon) = 1/(\pi - e)$$

$$8/3(\varepsilon) = 1/0.4233 = 0.8859 = h = k$$

Where $v = 0.27 \omega a/c = 6$

$$(1) a/4.3079 = 6.67a = 28.7 = r \text{ (Figure 6)}$$

$$y = e^{-x}$$

$$y(0) = 2(4/3) = 8.3$$

$$y(x) = 4/3s = Ae^{-st}$$

$$s(0) = Ae^{(-0)t} = 8/3$$

$$A = 8/3 = 2.667s(x) = 4/3$$

$$= 2.667(e^{-4/3t})0.5$$

$$= e^{(-st)} \text{Ln}(1/2)$$

$$= \text{Ln}(e^{(-st)}t)$$

$$= 0.51990.52$$

$$t - 29.788^0 = 30^0 \text{ if } t = 0.5199t/22LY = 0.0236 = 1/0.4233 = 1/cuz = E = \text{youngs Modulus}$$

And

$$t = 0.5199t/66 = 0.0079$$

$$= 1/(126.9)$$

$$= 1/\rho$$

So there you have it $v = 0.27, E = 1/0.4233$, and $\rho = 0.127$ (Figures 7-9)

To verify the calculations of the outer and inner universe, the dimension of the outer ring was determined to be within experimental error [5]

$$4/3 + 28.9 + 4/3 = \text{Diameter}$$

$$(1/2) * (1/R2) = (1/2) * (1/c2)$$

$$R2 = 1/2(5.55) = 0.9009 = c2$$

$$C^2 = R$$

$$\text{Area} = \pi R^2 = \pi(c^2) \text{circum.} = 2\pi * R = 2\pi * c^2$$

$$\text{Area/Circumference} = 1/2$$





Figure 6: Illustration of plate with a hole and the exponential decline.



$$\sin \theta = 1/\sqrt{2} = c, c^2 = 1/2$$

$$N = |\nu| |\nu| \sin \theta$$

$$N' = |\nu| |\nu| \cos \theta$$

$$N = (0.27)(\sqrt{2}) \sin 74.34 = 0.3674$$

$$N' = \int 1/e = \int (1/x) = -1 = 1/e$$

$$N = \int 1/e = \int (1/x) = -1 = 1/e$$

$$N = \int 1/e = \int (1/x) = -1 = 1/e$$

$$N = \int 1/e = \int (1/x) = -1 = 1/e$$

$$N = \int 1/e = \int (1/x) = -1 = 1/e$$

Figure 7: Illustration of 7 dot and cross products.




Figure 8: Illustration of the cylindrical universe and wave dissipation.




Figure 9: Illustration 9 age of the universe.

$$\partial s / \partial t = |E| t | \cos \theta$$

$$\theta = 60^\circ$$

$$1/2(1) (1) 1/2 = 1/4$$

$$4 \text{Area} = \text{circu. } R = 1/2$$

So,

$$C^2 = 1/2c$$

$$c^2 = \frac{1}{2c} = \frac{1}{\sqrt{2}} = \sin \theta - \cos \theta$$

$$c = \frac{1}{\sqrt{2}} = 2.997929c = \text{cuz}$$

The Universe is 13.34+0.289=13.629, within 13.73+/-0.12 range=13.61. This confirms my entire theory, My calculations agree with observation.

$$F = Ma$$

$$8.3 = (M_{\text{proton}} + M_{\text{electron}}) = (943 + 1.602)a = 9.445a; a = 2.823$$

2.823/cuz = 6.668 ~ G = 6.67 To produce Mass, some much Work goes into the Density and so much goes into the Spring or overcoming the charge resistance. I calculate it as follows:

$$M_p + M_e = 943 + 1.602 = 944.6$$

Why are there 116 stable elements in the Periodic table whereas there should be 118?

$$944.6 \times 2 = 1189.2$$

$$\Delta F m = 119.2 / 1189.2 / 0.4233 = 2809$$

$$\Delta F = \sin \theta = 2809 - 26.667 = 0.86441$$

$$\arcsin \theta (86441) = 59.8157 \text{ degrees}$$

$$\sin 60^\circ - \sin 59.81 = 0.001613 \sim 1.618 = t$$

$$\lim_{\delta x \rightarrow 0} \Delta F / \Delta t = \sin \theta' = -\cos \theta = 534.246$$

$$\theta = 57^\circ$$

$$60^\circ - 57^\circ = 3^\circ$$

$$\sin 3^\circ = 0.05230.0523 / 2.667$$

$$= 196(\text{Infinity})$$

$$= 196/2$$

$$= 981$$

$$1/981$$

$$0.1019$$

$$N'$$

$$= (0.27)\sqrt{2} * \cos 74.34 \text{ deg}$$

$$1/N' = e$$

$$\Delta F = \{M p + M e-\} * 2 / \text{cuz}$$

$$\Delta F = \{M p + M e-\} * 2 / \{\pi - 1 / N'\}$$

$$N' = |v| |E| \cos \theta$$

$$2(4.486) / \Delta F = \pi - 1 / (0.27) \cos 60 \text{ (Mass=4.486)}$$

$$\rho = 0.1272$$

Now

$$\Delta F = 2809$$

$$\chi^* \rho = \Delta F$$

$$\chi^* = 2.2 - 0.83 \chi^{(1/3)} = 1.3022$$

Continuing,

$$2M / \Delta F = ET - 1 / Ev$$

$$((2 * 944.6) * 2) = \pi - 1 / v \cos \theta$$

$$[2 * (2M) - \rho] / [\Delta F] = \pi$$

Or

$$4Fc^2 \rho = Ft$$

$$F = 1 / \text{cuz} = E$$

Work Spring + Work Density = Work (Total)

$$4(t/\text{cuz})c^2 \rho = Ft$$

Or

$$P = [Ft - Mc^2] / 4$$


Cusack's Mass Formation Equations

$$\rho = \pi / 4 - \text{Re (Figure 10)}$$

If we take a stress cube out of the cylindrical wall, we find that, when theta = 60 degrees (Mohr coulomb Failure Plane), the sine wave force is enough to overcome the the resistance to mass $R_m 2.667 + 4 = 6.67 = G$,

$$\partial / L_0 = \epsilon = \sigma / E$$

$$= 4 / (1/\text{cuz}) = 1.6932$$



$$\partial/L0 = PL/AE$$

$$1.6392 = (2.667)L^3/AE$$

$$1.6392 = 2.667(4/3)\pi R^2 * (1/0.4233)$$

$$= 2/\pi = dM/dt * freq$$

$$\epsilon = freq * dM/dt$$

$$\epsilon = \sigma/E = freq * dM/dt$$

Note

$$\epsilon = \lim_{x \rightarrow \infty} \Delta / dt$$

$$\int \epsilon = \Delta$$

$$2E * 1 / \pi(M) = \sigma^2$$

$$2(1/0.4233) * (1/\pi)M = 1.5^2; M = 1 \text{ Mass Gap}$$

Finally,

Newton's Gravitational Formula:

$$F < G M1M2/R^2$$

$$F < 2/3 M(1)/(1)^2$$

$$1.5F < M$$

$$1.5(1) < M$$

$$1.5 < M \text{ QED}$$

Using Divergence, the Mass Gap can be found another way, as follows:

$$\text{div} \nabla = \lim_{\tau \rightarrow 0} \frac{1}{\tau} \times \int \text{ba} \nabla \delta \sigma$$

Where

$$\text{div} \nabla = \text{Scalar Field} = \text{Area of the circle with radius} = 1$$

$1/\tau = 1/c$ area of the vector field where $c = \text{speed of light}$. This is a complete and closed system in Hilbert Space.

$$\sigma = E \times \epsilon = (\pi - e) \times \epsilon = (0.4233)(1 - \sin 1)$$

$$\pi - e = c \times z = \text{Resistance of Energy to Mass formation.}$$

$$\epsilon = \delta L/L = \text{Moment} = 1 - \sin 1 = 0.1585 = 0.8415 = \sin 1$$

And,

$$\text{Area} = \text{circumference of a circle,}$$

$$y = y'$$

$$\pi R^2 = 2\pi R$$

$$R = 1$$

Why is the sin (1 rad) required?

$$S = |E||t|\sin\theta$$

$$E = 1, t = 1$$

$$s = \sin\theta$$

Dot Product = Cross Product

$$|E||t|\cos\theta = |E||t|\sin\theta$$

$$-\sin\theta = \cos\theta \text{ since } s = v = a = s' = s''$$

$$\Theta = 45 \text{ degrees}$$

$$\text{But } s = t = 1$$

$$\sin \theta = \sin t = \sin 1 = 0.8415$$

$$\text{Lorentz Transform} = \sqrt{1 + v^2/c^2}$$

$$= \sqrt{1 + 0.8415^2/2.99792} = 0.8481$$

$$\arcsin(0.8481) = 58 \text{ degrees } 1 \text{ rad}$$

So, continuing

$$\text{div} \nabla = 1/\nabla = 1/\delta \tau \times \nabla^2/2 * E * \epsilon$$

$$\text{div} \nabla * \delta \tau = \pi/2(1 - \sin 1)$$

$$\delta \tau = 1/c$$

So,

$$\text{div} \nabla = c/2 * \nabla^2 * \sigma$$

$$\text{div} \nabla / (\nabla^2 * \sigma) = 1.5 = \text{Mass gap}$$

$$A^2 / (\pi^2 * F) = 1.5$$

$$\pi^2 / \pi^2 / F = 1.5$$

$$1/F = \text{Mass Gap}$$


The Super force is $F = \sin \theta = \sin t$ where $t = 1$. (AstroTheology, Cusack's Universe)

$$1/\sin 1 \text{ rad} = 118.8 \text{ Mass of the last element 118 in the periodic table.}$$

The equation of the universe is,

$$\int \text{ba} 1/n! = E = ex$$

There are 118 chemical elements. So,



$$\ln(1/n!) = \ln(1/118!) = 4.482 = \text{Mass}$$

$$E = Mc^2 = 4.482(2.997929)^2 = 0.402 = \text{Reynold's Number.}$$

$$0.402/c = 0.402/2.9979 = 0.1341$$

Conjugate = $1 - 0.1341 = 0.866 = \sin 60 \text{ degrees} = \text{critical angle of repose for kaolinite Universe.}$

QED (Figure 11)

And, the Dirac delta function,

$$\delta = \{1/c \text{ for } |x| = |t| = c/2 = 3/2 = 1.5\} \text{ (Actually, } 1.499 \text{ as above)}$$

Conclusion

The mass gap exists because as we've seen in these last few

pages again, the gravity would pull apart any atoms above 6.67 or $n = 4 + \tau = 2.667$; and it takes $f = 2.667$ at 60 degrees to compress energy into the form mass. The resistance to massism = $cuz = \pi - e$.

References

1. Cusack P (2015) Astrotheology, the missing link, Cusack's model of the universe. Canada.
2. Beer FP, Johnston ER, Dewolf J, Mazurek DF (2012) Nanoscale mechanical tailoring of interfaces using self-assembled monolayers. Mechanics of Materials 98: 71-80.
3. Arnold V (2010) Introduction to Soil Dynamics. Springer
4. Jaffe A, Edward WN (2000) Witten Quantum Yang-Mills theory. Clay Mathematics Institute.
5. Byron FW, Fuller RW (1969) Mathematics of classical and quantum physics, Dover, Newyork.

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://omicsonline.com/open-access/physical-mathematics.php>