

ON SOME DETERMINANT EQUATION

By Hiraku TOYAMA

In his theory of communication, C.E. Shannon⁽¹⁾ determined the channel capacity of a discrete noiseless system by means of a determinant equation of the following type:

$$|E - A(z)| = 0,$$

where $A(z)$ is a square matrix dependent on a complex variable z .

In this note I will prove the existence of a real positive root of the smallest absolute value, which is assumed in Shannon's theory.

Theorem.

Let $A(z)$ be a matrix subject to the following conditions:

- (1) $A(z)$ is a square matrix of order n .
- (2) Every matrix element $A_{ik}(z)$ of $A(z)$ is an entire function of z

$$A_{ik}(z) = \sum_{m=0}^{\infty} A_{ik,m} z^m$$

and

$$A_{ik}(0) = 0$$

- (3) Every coefficient $A_{ik,m}$ of $A_{ik}(z)$ is non-negative

$$A_{ik,m} \geq 0.$$

- (4) At least one coefficient of the characteristic polynomial $|\lambda E - A(z)|$ is not constant.

Then the determinant equation

$$|E - A(z)| = 0$$

has a real positive root of the smallest⁽²⁾ absolute value.

Lemma 1. If all of the traces $\text{Tr}(A(z)), \text{Tr}(A^2(z)), \dots, \text{Tr}(A^n(z))$ of

$A^k(z)$ are constant, then the characteristic polynomial has constant coefficients.

Proof. Let $\lambda_1, \lambda_2, \dots, \lambda_n$ be eigenvalues of $A(z)$, then

$$\text{Tr}(A(z)) = \lambda_1 + \lambda_2 + \dots + \lambda_n = \text{const}$$

$$\text{Tr}(A^2(z)) = \lambda_1^2 + \lambda_2^2 + \dots + \lambda_n^2 = \text{const}$$

$$\text{Tr}(A^n(z)) = \lambda_1^n + \lambda_2^n + \dots + \lambda_n^n = \text{const}$$

From these equations follows

$$\lambda_1 + \lambda_2 + \dots + \lambda_n = \text{const}$$

$$\lambda_1 \lambda_2 + \lambda_1 \lambda_3 + \dots + \lambda_{n-1} \lambda_n = \text{const}.$$

$$\lambda_1 \lambda_2 \lambda_3 \dots \lambda_n = \text{const}$$

by the theorem of symmetric functions.

Lemma 2. The matrix $(E - A(z))^{-1}$ is not an entire function.

Proof. By Lemma 1, at least one $\text{Tr}(A^k(z))$ is not constant for $1 \leq k \leq n$.

Hence at least one diagonal element of $A^k(z)$ is not all constant.

$$A_{jj}^k(z) = \sum_{m=0}^{\infty} A_{jj,m}^k z^m$$

Let $A_{jj,p}^k$ be a non-zero coefficient of the smallest order in the above expansion. Because all the coefficients is non-negative, the following inequality holds:

$$A_{jj,sp}^{ks} \geq (A_{jj,p}^k)^s$$

In the expansion

$$\sum_{\ell=0}^{\infty} A(z)^\ell$$