

# A RELATION BETWEEN JACOBI AND LAGUERRE POLYNOMIALS

EARL D. RAINVILLE

1. **Introduction.** We shall obtain an apparently new expression for the Jacobi polynomial  $P_n^{(\alpha, \beta)}(x)$  as a series of products of Laguerre polynomials of orders  $\alpha$  and  $\beta$  and of different arguments.

2. **A relation between the Legendre and Laguerre polynomials.** Because of the popularity of the Legendre and the simple Laguerre polynomials, we shall prove that special case of the relation between Jacobi and Laguerre polynomials. The method of proof carries through with no alteration in the general case and need not be repeated.

The generating function

$$(1) \quad e^t J_0(2(ty)^{1/2}) = \sum_{n=0}^{\infty} L_n(y) \frac{t^n}{n!}$$

for the Laguerre polynomials may be found in the standard books. For the Legendre polynomials a special case of a well known result<sup>1</sup> of Bateman is

$$(2) \quad I_0((2t(x-1))^{1/2}) I_0((2t(x+1))^{1/2}) = \sum_{n=0}^{\infty} P_n(x) \frac{t^n}{(n!)^2}.$$

From (1) with  $y = (1-x)/2$  it follows, since  $J_0(iy) = I_0(y)$ , that

$$e^t I_0((2t(x-1))^{1/2}) = \sum_{n=0}^{\infty} L_n\left(\frac{1-x}{2}\right) \frac{t^n}{n!}.$$

Using (1) with  $y = (1+x)/2$  and with  $t$  replaced by  $(-t)$ , we may conclude that

$$e^{-t} I_0((2t(x+1))^{1/2}) = \sum_{n=0}^{\infty} (-1)^n L_n\left(\frac{1+x}{2}\right) \frac{t^n}{n!}.$$

Then, with the aid of the Cauchy product of two series, it may be seen that

---

Received by the editors September 19, 1944.

<sup>1</sup> H. Bateman, *A generalization of the Legendre polynomial*, Proc. London Math. Soc. (2) vol. 3 (1905) pp. 111-123. See his formula (8).