

Values of Brownian intersection exponents, I: Half-plane exponents

by

GREGORY F. LAWLER

*Duke University
Durham, NC, U.S.A.*

ODED SCHRAMM

*Microsoft Research
Redmond, WA, U.S.A.*

and

WENDELIN WERNER

*Université Paris-Sud
Orsay, France*

and

*The Weizmann Institute of Science
Rehovot, Israel*

1. Introduction

Theoretical physics predicts that conformal invariance plays a crucial role in the macroscopic behavior of a wide class of two-dimensional models in statistical physics (see, e.g., [4], [6]). For instance, by making the assumption that critical planar percolation behaves in a conformally invariant way in the scaling limit, and using ideas involving conformal field theory, Cardy [7] produced an exact formula for the limit, as $N \rightarrow \infty$, of the probability that, in two-dimensional critical percolation, there exists a cluster crossing the rectangle $[0, aN] \times [0, bN]$. Also, Duplantier and Saleur [13] predicted the “fractal dimension” of the hull of a very large percolation cluster. These are just two examples among many such predictions.

In 1988, Duplantier and Kwon [12] suggested that the ideas of conformal field theory can also be applied to predict the intersection exponents between random walks in \mathbf{Z}^2 (and Brownian motions in \mathbf{R}^2). They predicted, for instance, that if B and B' are independent planar Brownian motions (or simple random walks in \mathbf{Z}^2) started from distinct points in the upper half-plane $\mathbf{H} = \{(x, y) : y > 0\} = \{z \in \mathbf{C} : \text{Im}(z) > 0\}$, then when $n \rightarrow \infty$,

$$\mathbf{P}[B[0, n] \cap B'[0, n] = \emptyset] = n^{-\zeta + o(1)} \quad (1.1)$$

and

$$\mathbf{P}[B[0, n] \cap B'[0, n] = \emptyset \text{ and } B[0, n] \cup B'[0, n] \subset \mathbf{H}] = n^{-\tilde{\zeta} + o(1)}, \quad (1.2)$$

where

$$\zeta = \frac{5}{8}, \quad \tilde{\zeta} = \frac{5}{3}.$$

Very recently, Duplantier [11] gave another physical derivation of these conjectures based on “quantum gravity”.

In 1982, Mandelbrot [35] suggested that the Hausdorff dimension of the Brownian frontier (i.e., the boundary of a connected component of the complement of the path) is $\frac{4}{3}$, based on simulations and the analogy with the conjectured value for the fractal dimension of self-avoiding walks predicted by Nienhuis (also $\frac{4}{3}$; see, e.g., [33]).

To date, none of the physicists’ arguments have been made rigorous, and it seems very difficult to use their methods to produce proofs. Very recently, Kenyon [16], [17], [18] managed to derive the exact values of critical exponents for “loop-erased random walk” that theoretical physicists had predicted (Majumdar [34], Duplantier [10]). Kenyon’s methods involve the relation of the loop-erased walk to the uniform spanning tree and to domino tilings. Kenyon shows that the equations relating probabilities of some domino tiling events are discrete analogues of the Cauchy–Riemann equations, and therefore the probabilities can be approximated by analytic functions with prescribed boundary behavior. These methods do not seem applicable for the goals of the present paper.

For planar Brownian motions, it is easy to show, using subadditivity arguments and the scaling property, that there exist positive finite numbers ζ and $\tilde{\zeta}$ such that (1.1) and (1.2) are true. Up to the present paper, there was not even a mathematical heuristic arguing that the values of ζ and $\tilde{\zeta}$ are $\frac{5}{8}$ and $\frac{5}{3}$. Burdzy–Lawler [5] (see also [9], [24]) showed that the intersection exponents were indeed the same for simple random walks as for Brownian motions; Lawler [21] proved that the Hausdorff dimension of the set of cut points of a Brownian path is $2-2\zeta$. He also showed (see [22], [23]) that the dimension of the Brownian frontier (and more generally the whole multifractal spectrum of the Brownian frontier) can be expressed in terms of exponents defined analogously to ζ . As part of that work, he showed that the right-hand side of (1.1) can be replaced with $n^{-\zeta}g(n)$ where g is bounded away from 0 and infinity; we expect that the argument can be adapted to show that the same is true for (1.2).

Recently, Lawler and Werner [28] extended the definition of intersection exponents in a natural way to “non-integer packets of Brownian motions”, and derived certain functional relations between these exponents. These relations indicate that Mandelbrot’s conjecture that the dimension of the Brownian frontier is $\frac{4}{3}$ is indeed compatible with the predictions of Duplantier–Kwon. It turned out that intersection exponents in the half-plane play an important role in understanding exponents in the whole plane. Conformal invariance of planar Brownian motion is a crucial tool in the derivation of these relations. In particular, there is a measure on Brownian excursions in domains that has some strong conformal invariance properties, including a “restriction” (or “locality”) property.

In another paper, Lawler and Werner [29] showed that intersection exponents as-

sociated to any conformally invariant measure on sets with this restriction property are very closely related to the Brownian exponents. This provides a rigorous justification to the link between the conjectures regarding intersection exponents for planar Brownian motions and conjectures for intersection exponents of critical percolation clusters (see [13], [8], [3]), because percolation clusters are conjectured to be conformally invariant in the scaling limit—see, e.g., [19], [2]—and they should also have a restriction property (because of the independence properties of percolation). The question of how to compute these exponents remained open.

Independently, Schramm [42] defined a new class of conformally invariant stochastic processes indexed by a real parameter $\kappa \geq 0$, called SLE_κ (for stochastic Löwner evolution process with parameter κ). The definition of these processes is based on Löwner's ordinary differential equation that encodes in a conformally invariant way a continuous family of shrinking domains (see, e.g., [32], [37]). More precisely, [42] defines a family of conformal maps g_t from subsets D_t of \mathbf{H} onto \mathbf{H} by the equation

$$\partial_t g_t(z) = \frac{-2}{\beta_{\kappa t} - g_t(z)}, \quad (1.3)$$

where β is a standard Brownian motion on the real line. (Actually, in [42], instead of (1.3), the corresponding equation for the inverse maps g_t^{-1} is considered.) The domain D_t can be defined as the set of $z_0 \in \mathbf{H}$ such that a solution $g_s(z_0)$ of this equation exists for $s \in [0, t]$. When t increases, the set $K_t = \mathbf{H} \setminus D_t$ increases: Loosely speaking, $(K_t, t \geq 0)$ can be viewed as a growing “hull” that is penetrating the half-plane. By applying a conformal homeomorphism $f: \mathbf{H} \rightarrow D$, SLE_κ can similarly be defined in any simply-connected domain $D \subsetneq \mathbf{C}$.

In [42], the main focus is on the case $\kappa = 2$, which is conjectured there to correspond to the scaling limit of loop-erased random walks, but the conjecture that SLE_6 corresponds to the scaling limit of critical percolation cluster boundaries is also mentioned. In particular, it is possible to compute explicitly the probability that an SLE_6 crosses a rectangle of size $a \times b$. It turns out that this result is exactly Cardy's formula. This gives a mathematical proof for Cardy's formula, assuming the still open conjecture that SLE_6 is indeed the scaling limit of percolation cluster boundaries.

The main goal of the present paper is to prove some of the conjectured values of intersection exponents of Brownian motion in a half-plane.

THEOREM 1.1. *Let B^1, \dots, B^p denote p independent planar Brownian motions ($p \geq 2$) started from distinct points in the upper half-plane \mathbf{H} . Then, when $t \rightarrow \infty$,*

$$\mathbf{P}[\forall i \neq j \in \{1, \dots, p\}, B^i[0, t] \cap B^j[0, t] = \emptyset \text{ and } B^i[0, t] \subset \mathbf{H}] = t^{-\tilde{\zeta}_p + o(1)},$$

where

$$\tilde{\zeta}_p = \frac{1}{6}p(2p+1).$$

These values have been predicted by Duplantier and Kwon [12]. In particular, $\tilde{\zeta} = \tilde{\zeta}_2 = \frac{5}{3}$.

We also establish the exact value (and confirm some of the conjectures stated in [28], [11]) of more general intersection exponents between packets of Brownian motions in the half-plane; see Theorem 4.1.

The proof of Theorem 1.1 uses a combination of ideas from the papers [28], [29], [42]. However, to make the paper more accessible and self-contained, we attempt to review and explain all the necessary background. The reader who wishes to see complete proofs for all stated theorems has to be familiar with the basics of stochastic calculus and conformal mapping theory, and read about the excursion measure and the cascade relations from [28].

Although, at present, a proof of the conjecture that SLE_6 is the scaling limit of critical percolation cluster boundaries seems out of reach, this conjecture does lead one to believe that SLE_6 must satisfy a “locality” property; namely, it is not affected by the boundary of a domain when it is in the interior. This locality property for SLE_6 is stated more precisely and proved in §2. It is worthwhile to note that the locality property does not hold for the SLE_κ -processes when $\kappa \neq 6$.

In §3, we prove that SLE_6 satisfies a generalization of Cardy’s formula for percolation crossings probabilities. From this, exponents associated with the SLE_6 -process are computed.

In §4, universality ideas from [29] are used to compute the half-plane Brownian exponents from the SLE_6 -exponents, which completes the proof of Theorem 1.1.

In a final short §5, the conjectured relationship between SLE_6 and critical percolation is discussed. It is demonstrated that this conjecture implies a formula from the physics literature [13], [8], [3] for the exponents corresponding to the event that there are k disjoint percolation crossings of a long rectangle.

In the subsequent papers [25], [26], [27], we determine the exponents in the full plane and the remaining half-plane exponents. In particular, we prove that $\zeta = \frac{5}{8}$, and also establish Mandelbrot’s conjecture that the Hausdorff dimension of the frontier of planar Brownian motion is $\frac{4}{3}$.

It might be worthwhile to explain why the Brownian intersection exponents are accessible through SLE_6 , but are difficult to compute directly. In a way, the SLE_6 -process is simpler, since K_t continuously grows from its outer boundary. This means that when studying its evolution, one can essentially forget its interior, and only keep track of the exterior of K_t . By conformal invariance, this reduces problems to finitely many dimensions. The situation with planar Brownian motion is completely different, since it may enter holes it has surrounded and emerge to the exterior someplace else.

Many computations with SLE_κ are readily convertible to PDE problems, and in the presence of enough symmetry, some variables can often be eliminated, converting the PDE to an ODE.

2. SLE_6 and its locality property

2.1. The definition of chordal SLE_κ and some basic properties

Let $(\beta_t, t \geq 0)$ be a standard real-valued Brownian motion starting at $\beta_0=0$, let $\kappa > 0$, and let $W_t^\kappa = \beta_{\kappa t}$. Consider the ordinary differential equation

$$\partial_t g_t(z) = \frac{-2}{W_t^\kappa - g_t(z)} \tag{2.1}$$

with $g_0(z)=z$. For every $z_0 \in \mathbf{H}$ and every $T > 0$, either there is a solution of (2.1) for $t \in [0, T]$ and for all z in a neighborhood of z_0 , or there is some $t_0 \in (0, T]$ such that the solution exists for $t \in [0, t_0)$ and $\lim_{t \nearrow t_0} g_t(z) = W_{t_0}^\kappa$. Let D_T be the (open) set of $z \in \mathbf{H}$ such that the former is true, and let K_T be the set of $z \in \mathbf{H}$ such that the latter holds. By considering the inverse flow $\partial_t G_t(z) = 2(W_{T-t}^\kappa - G_t(z))^{-1}$, it is easy to see that $g_t(D_t) = \mathbf{H}$, and that $g_t: D_t \rightarrow \mathbf{H}$ is conformal. The process $g_t, t \geq 0$, will be called the *chordal stochastic Löwner evolution process* with parameter κ , or just SLE_κ ; see [42]. In [42], a variation of this process, which we now call *radial SLE_κ* was also studied. In the current paper, we will not use radial SLE_κ , and therefore the word ‘‘chordal’’ will usually be omitted. (However, radial SLE_κ plays a major role in a subsequent paper [25].) The set $K_t = \mathbf{H} \setminus D_t$ will be called the *hull* of the SLE. The process W_t^κ will be called the *driving process* of the SLE.

It is easy to verify that each of the maps g_t satisfies the *hydrodynamic normalization* at infinity:

$$\lim_{z \rightarrow \infty} g(z) - z = 0. \tag{2.2}$$

Remarks. It will be shown [40] that for all $\kappa \neq 8$ the hull K_t of SLE_κ is a.s. generated by a path. More precisely, a.s. the map $\gamma(t) := g_t^{-1}(W_t^\kappa)$ is a well-defined continuous path in $\overline{\mathbf{H}}$, and for every $t \geq 0$ the domain D_t is the unbounded connected component of $\mathbf{H} \setminus \gamma([0, t])$. There, it will also be shown that when $\kappa \leq 4$ a.s. K_t is a simple path for all $t > 0$. This is not the case when $\kappa > 4$ [42]. However, these results will not be needed for the current paper or for [25], [26], [27].

Löwner [32] considered the equation

$$\partial_t g_t(z) = g_t(z) \frac{\zeta(t) + g_t(z)}{\zeta(t) - g_t(z)},$$

with $g_0(z)=z$, where z is in the unit disk, and $\zeta(t)$ is a parameter. He used this equation in the study of extremal problems for classes of normalized conformal mappings. In Löwner's differential equation, the maps g_t satisfies $g_t(0)=0$. The equation (2.1) is an analogue of Löwner's equation in the half-plane, where the boundary point ∞ is fixed instead of 0, and $\zeta(t)$ is chosen to be scaled Brownian motion.

Marshall and Rohde [36] study conditions on $\zeta(t)$ which imply that K_t is a simple path.

We now note some basic properties of SLE_κ .

PROPOSITION 2.1. (i) [Scaling] SLE_κ is scale-invariant in the following sense. Let K_t be the hull of SLE_κ , and let $\alpha > 0$. Then the process $t \mapsto \alpha^{-1/2} K_{\alpha t}$ has the same law as $t \mapsto K_t$.

(ii) [Stationarity] Let g_t be an SLE_κ -process in \mathbf{H} , driven by W_t^κ , and let τ be any stopping time. Set $\tilde{g}_t(z) = g_{\tau+t} \circ g_\tau^{-1}(z + W_\tau^\kappa) - W_\tau^\kappa$. Then \tilde{g}_t is an SLE_κ -process in \mathbf{H} starting at 0, which is independent from $\{g_t : t \in [0, \tau]\}$.

Proof. (i) If K_t is driven by W_t^κ , then $\alpha^{-1/2} K_{\alpha t}$ is driven by $\alpha^{-1/2} W_{\alpha t}^\kappa$, which has the same law as W_t^κ .

(ii) The process \tilde{g}_t is driven by $W_{t+\tau}^\kappa - W_\tau^\kappa$. □

We now consider the definition of SLE_κ in domains other than \mathbf{H} .

Let $f: D \rightarrow \mathbf{H}$ be a conformal homeomorphism from some simply-connected domain D . Let f_t be the solution of (2.1) with $f_0(z) = f(z)$. Then $(f_t, t \geq 0)$ will be called the SLE_κ in D starting at f . If g_t is the solution of (2.1) with $g_0(z) = z$, then we have $f_t = g_t \circ f$. If K_t is the hull associated to g_t , then the hull associated with f_t is just $f^{-1}(K_t)$.

Suppose that ∂D is a Jordan curve in \mathbf{C} , and let $a, b \in \partial D$ be distinct. Then we may find such an $f: D \rightarrow \mathbf{H}$ with $f(a) = 0$ and $f(b) = \infty$. Let K_t^f be the SLE_κ -hull associated with the SLE_κ -process starting at f . If $f^*: D \rightarrow \mathbf{H}$ is another such map with $f^*(a) = 0$ and $f^*(b) = \infty$, then $f^*(z) = \alpha f(z)$ for some $\alpha > 0$. By Proposition 2.1, the corresponding SLE_κ -hull $K_t^{f^*}$ has the same law as a linear time change of K_t^f . This makes it natural to consider K_t^f as a process from a to b in D , and to ignore the role of f . However, when D is not a Jordan curve, some care may be needed since the conformal map f does not necessarily extend continuously to the boundary. Partly for that reason, we have chosen to stress the importance of the conformal parameterization f .

2.2. The locality property

The main result of this section can be loosely described as follows: an SLE_6 process does not feel where the boundary of the domain lies as long as it does not hit it. This is consistent with the conjecture [42] that the SLE_6 -process is the scaling limit of percolation cluster boundaries, which is explained in §5. This restriction property can therefore be viewed as additional evidence in favor of this conjecture. This feature is special to SLE_6 ; it is not shared by SLE_κ when $\kappa \neq 6$.

Such properties were studied in [29] and called “complete conformal invariance” (when combined with a conformal invariance property). As pointed out there, all processes with complete conformal invariance have closely related intersection exponents.

Let us first state a general local version of this result. We will say that the path γ is nice if it is a continuous simple path $\gamma: [0, 1] \rightarrow \bar{\mathbf{H}}$ such that $\gamma(0), \gamma(1) \in \mathbf{R} \setminus \{0\}$ and $\gamma(0, 1) \subset \mathbf{H}$. We then call the connected component $N = N(\gamma)$ of $\mathbf{H} \setminus \gamma[0, 1]$ such that $0 \in \partial N$ a nice neighborhood of 0 in \mathbf{H} . Note that N can be bounded or unbounded, depending on the sign of $\gamma(0)\gamma(1)$. When N is a nice neighborhood of 0, one can find a conformal homeomorphism $\psi = \psi_N$ from N onto \mathbf{H} such that $\psi(0) = 0$, $\psi'(0) = 1$ and $\psi^{-1}(\infty)$ is equal to ∞ if N is unbounded and to $\gamma(1)$ if N is bounded.

THEOREM 2.2 (locality). *Let $f: D \rightarrow \mathbf{H}$ be a conformal homeomorphism from a domain $D \subset \mathbf{C}$ onto \mathbf{H} . Suppose that N is a nice neighborhood of 0 in \mathbf{H} . Define $D^* = f^{-1}(N)$ and let f^* be the conformal homeomorphism $\psi_N \circ f$ from D^* onto \mathbf{H} . Let $K_t \subset D$ be the hull of SLE_6 starting at f , and let $\tau := \sup\{t: \bar{K}_t \cap \partial D^* \cap D = \emptyset\}$. Let K_t^* denote SLE_6 in D^* started at f^* , and let $\tau^* := \sup\{t: \bar{K}_t^* \cap \partial D^* \cap D = \emptyset\}$.*

Then the law of $(K_t, t < \tau)$ is that of a time change of $(K_t^, t < \tau^*)$.*

Note that in this theorem, we have not made any regularity assumption on the boundary of the domain D .

A consequence of this result is that, modulo time change, one can define the hull of SLE_6 in a non-simply-connected domain with finitely many boundary components, since such a domain looks locally like a simply-connected domain.

This property implies the following “global” restriction properties. For convenience only, we will state them under some assumptions on the boundaries of the domains.

COROLLARY 2.3 (splitting property). *Let D denote a simply-connected domain such that ∂D is a Jordan curve. Let a, b and b' denote three distinct points on ∂D , and let I denote the connected component of $\partial D \setminus \{b, b'\}$ that does not contain a . Let $(K_t, t \geq 0)$ (resp. K'_t) denote an SLE_6 in D from a to b (resp. from a to b'). Let T (resp. T') denote the first time at which \bar{K}_t (resp. \bar{K}'_t) intersects I . Then $(K_t, t < T)$ and $(K'_t, t < T')$ have the same law up to time change.*

COROLLARY 2.4 (restriction property). *Let $D^* \subset D$ denote two simply-connected domains, and assume that ∂D is a Jordan curve. Suppose that $I := \partial D^* \setminus \partial D$ is connected. Take two distinct points a and b in $\partial D \cap \partial D^* \setminus \bar{I}$.*

Let $(K_t, t \geq 0)$ denote SLE₆ from a to b in D , and $T := \sup\{t : \bar{K}_t \cap I = \emptyset\}$. Similarly, let $(K_t^, t \geq 0)$ be SLE₆ from a to b in D^* , and $T^* := \sup\{t : \bar{K}_t^* \cap I = \emptyset\}$. Then, $(K_t, t < T)$ and $(K_t^*, t < T^*)$ have the same law up to time change.*

In the present paper, we will use these results when D is a rectangle.

Proof of Corollary 2.3 (assuming Theorem 2.2). This is just a consequence of the fact that in Theorem 2.2 with $D = \mathbf{H}$ and bounded N , one can replace γ by $\beta(s) := \gamma(1-s)$. Then we get that the law of SLE₆ in N from 0 to $\gamma(0)$ is that of a time change of SLE₆ in N from 0 to $\gamma(1)$ up to their hitting times of γ . The result in a general domain follows by mapping it conformally onto a nice neighborhood N with a mapped to 0 and $\{b, b'\}$ to $\{\gamma(0), \gamma(1)\}$. \square

Proof of Corollary 2.4 (assuming Theorem 2.2). By approximation, it suffices to consider the case where I is a simple path. Let f denote a conformal map from D onto \mathbf{H} , with $f(a) = 0$ and $f(b) = \infty$. Define γ in such a way that $\gamma[0, 1] = f(I)$; note that $D^* = f^{-1}(N(\gamma))$. As $b \in \partial D^* \setminus I$, $N(\gamma)$ is unbounded. Hence, by Theorem 2.2, the law of SLE₆ in D from a to b stopped when it hits I is (up to time change) the same as that of SLE₆ in D^* from a to b stopped when its closure hits I . \square

In order to prove Theorem 2.2, we will establish

LEMMA 2.5. *Under the assumptions of Theorem 2.2, define for any fixed $s < 1$, $L_s = \gamma(0, s]$ and*

$$T = \sup\{t \geq 0 : \bar{K}_t \cap \bar{L}_s = \emptyset\}.$$

For all $t \leq T$, let $g_{s,t}$ denote the conformal homeomorphism taking $\mathbf{H} \setminus (K_t \cup L_s)$ onto \mathbf{H} with the hydrodynamic normalization. Then, the process $(g_{s,t}, t < T)$ has the same law as a time change of SLE₆ in $\mathbf{H} \setminus L_s$ starting at $g_{s,0} - g_{s,0}(0)$, up to the time when the closure of its hull intersects \bar{L}_s .

Proof of Theorem 2.2 (assuming Lemma 2.5). In the setting of the lemma, let

$$h_{s,t}(z) = \frac{g_{s,t}(z) - g_{s,0}(0)}{g'_{s,0}(0)}.$$

By Lemma 2.5 and Proposition 2.1, $t \mapsto h_{s,t}$ has the same law as a time change of SLE₆ starting at $h_{s,0}$. Note that $h_{s,0}(0) = 0$, $h'_{s,0}(0) = 1$. Hence, it follows easily that for all $z \in N(\gamma)$,

$$\lim_{s \rightarrow 1} h_{s,0}(z) = \psi_N(z).$$

By continuity, if we let $h_{1,t} = \lim_{s \rightarrow 1} h_{s,t}$, then $t \mapsto h_{1,t}$ has the same law as a time-changed SLE_6 (in N) started from ψ_N . The proof is completed by noting that the hull of $h_{1,t}$ is K_t . \square

The idea in the proof of Lemma 2.5 is to study how the process $g_{s,t}$ changes as s increases. For this, we will need to use some of the properties of solutions to (2.1) where W^κ is replaced by other continuous functions, and to study how (deterministic) families of conformal maps can be represented in this way with some driving function.

2.3. Deterministic expanding hulls

2.3.1. *Definition and first properties.* If $(U_t, t \in [0, a])$ is a continuous real-valued function, then the process defined by

$$\partial_t g_t(z) = \frac{-2}{U_t - g_t(z)} \tag{2.3}$$

and $g_0(z) = z$ will be called the *Löwner evolution with driving function U_t* . Note that g_t satisfies the hydrodynamic normalization (2.2). Moreover,

$$g_t(z) = z + 2tz^{-1} + a_2(t)z^{-2} + \dots, \quad z \rightarrow \infty, \tag{2.4}$$

for some functions $a_j(t)$, $j = 2, 3, \dots$. As above, we let $D_t \subset \mathbf{H}$ denote the domain of g_t , and let $K_t := \mathbf{H} \setminus D_t$. K_t will be called the *expanding hull* of the process g_t .

We now address the question of which processes K_t can appear as the expanding hull driven by a continuous function U_t . We say that a bounded set $K \subset \mathbf{H}$ is a *hull* if $\mathbf{H} \setminus K$ is open and simply-connected. The Riemann mapping theorem tells us that for each hull K , there is a unique conformal homeomorphism $g_K: \mathbf{H} \setminus K \rightarrow \mathbf{H}$ which satisfies the hydrodynamic normalization (2.2). Let

$$A(K) = A(g_K) := \frac{1}{2} \lim_{z \rightarrow \infty} z(g_K(z) - z);$$

that is, $g(z) = z + 2A(g)z^{-1} + \dots$ near ∞ . Observe that $A(K)$ is real, because $g_K(x)$ is real when $x \in \mathbf{R}$ and $|x|$ is sufficiently large. Moreover, $A(K) \geq 0$, because $\text{Im}(z - g_K(z))$ is a harmonic function which vanishes at infinity and has non-negative boundary values. Note that

$$A(g \circ h) = A(g) + A(h)$$

if g and h satisfy the hydrodynamic normalization. It follows that $A(K) \leq A(L)$ when $K \subset L$, since $g_L = g_{g_K(L \setminus K)} \circ g_K$.

The quantity $A(g)$ is similar to capacity, and plays an analogous role for the equation (2.1) as capacity plays for Löwner’s equation.

THEOREM 2.6. *Let $(K_t, t \in [0, a])$ be an increasing family of hulls. Then the following are equivalent:*

(1) *For all $t \in [0, a]$, $A(K_t) = t$, and for each $\varepsilon > 0$ there is a $\delta > 0$ such that for each $t \in [0, a - \delta]$ there is a bounded connected set $S \subset \mathbf{H} \setminus K_t$ with $\text{diam}(S) < \varepsilon$ and such that S disconnects $K_{t+\delta} \setminus K_t$ from infinity in $\mathbf{H} \setminus K_t$.*

(2) *There is some continuous $U: [0, a] \rightarrow \mathbf{R}$ such that K_t is driven by U_t .*

In [37] a similar theorem is proved for Löwner’s differential equation in the disk.

Note that K_t may change discontinuously, in the Hausdorff metric, as t increases. For example, consider $\widehat{K}_t := \{\exp(is) : 0 < s \leq t\}$ when $t < \pi$, and $\widehat{K}_\pi := \{z \in \mathbf{H} : |z| \leq 1\}$ and $\widehat{K}_{t+\pi} := K_\pi \cup (-1, -1+it]$, $t > 0$, say, and let $K_t := \widehat{K}_{\phi(t)}$ where ϕ is chosen to satisfy $A(K_{\phi(t)}) = t$.

LEMMA 2.7. *Let $r > 0$ and $x_0 \in \mathbf{R}$, and suppose that K is a hull contained in the disk $\{z : |z - x_0| < r\}$. Then*

$$\left| g_K^{-1}(z) - z + \frac{2A(K)}{z - x_0} \right| \leq \frac{CrA(K)}{|z - x_0|^2}$$

for all $z \in \mathbf{H}$ with $|z - x_0| > Cr$, where $C > 0$ is an absolute constant.

Proof of Lemma 2.7. For notational simplicity, we assume that $x_0 = 0$. Clearly, this does not entail any loss of generality. By approximation, we may assume that K has smooth boundary. Let $I \subset \mathbf{R}$ be the smallest interval in \mathbf{R} containing $\{g_K(x) : x \in \partial K \cap \mathbf{H}\}$, and let $f := g_K^{-1}$. Let f_I be the restriction of f to I . Let f^* denote the extension of f to $\mathbf{C} \setminus I$, by Schwarz reflection. The Cauchy formula gives

$$2\pi i f^*(w) = \int_{|z|=R} \frac{f^*(z)}{z-w} dz + \int_I \frac{f_I(x) - \overline{f_I(x)}}{x-w} dx,$$

provided that $R > |w|$, $R > \max\{|x| : x \in I\}$ and $w \in \mathbf{C} \setminus I$. Since $f^*(z) = z - 2A(K)z^{-1} + \dots$ near ∞ ,

$$\lim_{R \rightarrow \infty} \int_{|z|=R} \frac{f^*(z)}{z-w} dz = \lim_{R \rightarrow \infty} \int_{|z|=R} \frac{z}{z-w} dz = 2\pi i w.$$

Consequently, we have

$$f^*(w) - w = \frac{1}{\pi} \int_I \frac{\text{Im}(f_I(x))}{x-w} dx.$$

Multiplying by w and taking $w \rightarrow \infty$ gives

$$A(g_K) = -A(f^*) = \frac{1}{2\pi} \int_I \text{Im}(f_I(x)) dx. \tag{2.5}$$

Moreover,

$$f^*(w) - w - 2A(f^*)w^{-1} = \frac{1}{\pi} \int_I \operatorname{Im}(f_I(x)) \left(\frac{1}{x-w} + \frac{1}{w} \right) dx,$$

and therefore

$$\begin{aligned} |f^*(w) - w - 2A(f^*)w^{-1}| &\leq \frac{1}{\pi} \int_I \operatorname{Im}(f_I(x)) \sup\{|(x-w)^{-1} + w^{-1}| : x \in I\} dx \\ &= -2A(f^*) \sup\left\{ \left| \frac{x}{(x-w)w} \right| : x \in I \right\}. \end{aligned}$$

Hence, the proof will be complete once we demonstrate that there is some constant c_0 such that $I \subset [-c_0r, c_0r]$. This is easily done, as follows. Define $G(z) := g_K(rz)/r$ for $|z| > 1$, and write $G(z) = z + a_1z^{-1} + a_2z^{-2} + \dots$. The Area Theorem (see, e.g., [41]) gives $1 \geq \sum_{j=1}^\infty j|a_j|^2$. In particular, $|a_j| \leq 1$ for $j \geq 1$. Consequently, we have $|G(z) - z| \leq 1$ for $|z| \geq 2$. By Rouché’s theorem (e.g. [41]), it follows that $G(\{|z| \geq 2\}) \supset \{|z| > 3\}$. Consequently, $g_K(\mathbf{H} \setminus K) \supset \{|z| > 3r\}$, which gives $I \subset [-3r, 3r]$. \square

For convenience, we adopt the notation

$$K_{t,u} := g_{K_t}(K_{t+u} \setminus K_t).$$

Proof of Theorem 2.6. We start with (1) implies (2). Let $R := \sup\{|z| : z \in K_a\}$ and $Q := \{z \in \mathbf{H} : |z| > R + 2\}$. Let t, δ, ε and S be as in the statement of the theorem, and let $s \in \partial S$. Suppose that $\varepsilon < 1$ and $r \in [\varepsilon, \sqrt{\varepsilon}]$. Then there is an arc β_r of the circle of radius r about s such that $\beta_r \subset \mathbf{H} \setminus K_t$ and $K_t \cup \mathbf{R} \cup \beta_r$ separates $K_{t+\delta} \setminus K_t$ from Q . It therefore follows that the extremal length of the set of arcs in $\mathbf{H} \setminus K_t$ which separate $K_{t+\delta} \setminus K_t$ from Q in $\mathbf{H} \setminus K_t$ is at most $\operatorname{const}/\log(1/\varepsilon)$. (For the definition and basic properties of extremal length, see [1], [31]. The terms extremal length and extremal distance have the same meaning.) Since extremal length is invariant under conformal maps, it follows that the extremal length of the set of paths in \mathbf{H} that separate $K_{t,\delta}$ from $g_{K_t}(Q)$ is at most $\operatorname{const}/\log(1/\varepsilon)$. Because the diameter of $g_{K_t}(\mathbf{H} \setminus Q)$ is bounded by some function of R (this follows since g_{K_t} has the hydrodynamic normalization), we conclude that at least one of these arcs has length less than $\operatorname{const}/\log(1/\varepsilon)$. Consequently, this is a bound on the diameter of $K_{t,\delta}$. Observe that this bound is uniform for all $t \in [0, a - \delta]$. For each $t < a$, we then define U_t to be the point in the intersection $\bigcap_{u>0} \bar{K}_{t,u}$. We have an upper bound on $\operatorname{diam}(K_{t,\delta})$ which tends to zero uniformly as $\delta \rightarrow 0$, and therefore $\lim_{\delta \rightarrow 0} g_{K_{t,\delta}}(z) - z = 0$ uniformly for $z \in \mathbf{H} \setminus K_{t,\delta}$ and $t \leq a - \delta$. This implies that U_t is uniformly continuous on $[0, a)$ and can be extended continuously to $[0, a]$.

Now let $z_0 \in \mathbf{H} \setminus K_a$. Then there is some $c > 0$ such that $\operatorname{Im}(g_{K_t}(z_0)) > c$ for all $t \in [0, a]$. Lemma 2.7 applied with $K = K_{t,u}$, $z = g_{K_{t+u}}(z_0)$ and $x_0 = U_{t+u}$ gives

$$\frac{g_{K_t}(z_0) - g_{K_{t+u}}(z_0)}{u} + \frac{2}{g_{K_{t+u}}(z_0) - U_{t+u}} \rightarrow 0$$

as $\delta \rightarrow 0$. As $g_{K_t}(z_0)$ and U_t are continuous in t , we may therefore conclude that

$$\partial_t g_{K_t}(z_0) = \frac{2}{g_{K_t}(z_0) - U_t},$$

which gives (2).

The proof that (2) implies (1) is easy. Let $\varepsilon > 0$. Given $0 \leq t \leq t+u < a$, let $\varrho(t, u) := u + \max\{U(t') - U(t'') : t', t'' \in [t, t+u]\}$. Observe that $\text{diam}(K_{t,u}) \rightarrow 0$ if $\varrho(t, u) \rightarrow 0$, and $\varrho(t, u) \rightarrow 0$ if $u \rightarrow 0$. Consequently, the extremal length of the set of paths separating $K_{t,u}$ from $\{z \in \mathbf{H} : |z| > 1\}$ in \mathbf{H} goes to zero as $\varrho(t, u) \rightarrow 0$. This implies that there is a path β in this set such that $\text{diam}(g_{K_t}^{-1}(\beta)) < \varepsilon$, provided that u is small. We then just take $S = g_{K_t}^{-1}(\beta)$. \square

2.3.2. Time-modified expanding hulls and restriction. Let $(K_t, t \in [0, a])$ denote a family of hulls, and suppose that there is a monotone increasing homeomorphism $\phi: [0, a] \rightarrow [0, \hat{a}]$ such that $(K_{\phi(t)}, t \in [0, \hat{a}])$ is an expanding hull driven by some function $t \mapsto \hat{U}_t$. If additionally ϕ is continuously differentiable in $[0, a]$ and $\phi'(t) > 0$ for each $t \in [0, a]$, then we call $(K_t, t \in [0, a])$ a *time-modified* expanding hull, with driving function $U_t := \hat{U}_{\phi^{-1}(t)}$. Note that, in this case, $\phi^{-1}(t) = A(K_t)$, and that

$$\partial_t g_{K_t}(z) = \frac{2\partial_t A(K_t)}{g_{K_t}(z) - U_t}. \quad (2.6)$$

Note that in our terminology, an expanding hull is always a time-modified expanding hull.

LEMMA 2.8. *Let $(K_t, t \in [0, a])$ be a time-modified expanding hull, with driving function $(U_t, t \in [0, a])$. Let D be a relatively open subset of $\overline{\mathbf{H}}$ which contains \overline{K}_a , and set $D_{\mathbf{R}} := D \cap \mathbf{R}$. Let $G: D \rightarrow \overline{\mathbf{H}}$ be conformal in $D \setminus D_{\mathbf{R}}$ and continuous in D , and suppose that $G(D_{\mathbf{R}}) \subset \mathbf{R}$. Then $(G(K_t), t \in [0, a])$ is a time-modified expanding hull. Moreover,*

$$\partial_t A(G(K_t)) = G'(U_0)^2 \partial_t A(K_t) \quad \text{at } t = 0. \quad (2.7)$$

Proof. We first prove (2.7). The proof will be based on (2.5). Note first that if $K' = aK$ then $A(K') = a^2 A(K)$. Therefore, we may assume that $G'(U_0) = 1$. Similarly, with no loss of generality, we assume that $U_0 = G(U_0) = 0$. By the reflection principle, G is analytic in D .

Set $\hat{K}_t := G(K_t)$. Let $I_t \subset \mathbf{R}$ be the interval corresponding to $\overline{\partial K_t} \cap \overline{\mathbf{H}}$ under g_{K_t} , and let \hat{I}_t be the interval corresponding to $\overline{\partial \hat{K}_t} \cap \mathbf{H}$ under $g_{\hat{K}_t}$. Let $\varepsilon > 0$, and let $D_\varepsilon := \{z \in D : |1 - G'(z)| < \varepsilon\}$. Let β be some arc in $D_\varepsilon \setminus \{0\}$ that separates 0 from ∞ in $\overline{\mathbf{H}}$. Consider the map

$$h_t = g_{\hat{K}_t} \circ G \circ g_{K_t}^{-1}.$$

It is well defined in a neighborhood of I_t provided that $K_t \cap \beta = \emptyset$ (for instance), and this holds when t is small. This map may be continued analytically by reflecting in the real axis, and therefore the maximum principle implies that

$$\sup\{h'_t(x) : x \in I_t\} \leq \sup\{|h'_t(z)| : z \in g_{K_t}(\beta)\}$$

when $K_t \cap \beta = \emptyset$. Note that $g_{K_t}(z) - z \rightarrow 0$ and $g_{\widehat{K}_t}(z) - z \rightarrow 0$ as $t \searrow 0$, and therefore $g'_{K_t}(z) \rightarrow 1$ and $g'_{\widehat{K}_t}(z) \rightarrow 1$ on β . Consequently, for small t we have

$$\sup\{h'_t(x) : x \in I_t\} < 1 + 2\varepsilon. \tag{2.8}$$

Note that for z close to $U_0 = 0$ we have $\text{Im}(G(z)) \leq (1 + \varepsilon) \text{Im}(z)$. Using (2.5), this inequality and (2.8), we get

$$\begin{aligned} A(G(K_t)) &= \frac{1}{2\pi} \int_{I_t} \text{Im}(g_{\widehat{K}_t}^{-1}(x)) \, dx \\ &= \frac{1}{2\pi} \int_{I_t} \text{Im}(G \circ g_{\widehat{K}_t}^{-1}(x)) h'_t(x) \, dx \\ &\leq \frac{1}{2\pi} \int_{I_t} (1 + \varepsilon) \text{Im}(g_{\widehat{K}_t}^{-1}(x)) (1 + 2\varepsilon) \, dx \\ &= (1 + \varepsilon)(1 + 2\varepsilon) A(K_t) \end{aligned}$$

for small $t > 0$. (Note that $g_{\widehat{K}_t}^{-1}(x)$ is not defined for every $x \in I_t$, but it is defined for almost every $x \in I_t$.) By symmetry, we also have a similar inequality in the other direction. This proves (2.7).

By Theorem 2.6, to show that $G(K_t)$ is a time-modified expanding hull, it suffices to show that $A(G(K_t))$ is continuously differentiable in t , with derivative bounded away from 0. Let $G_t := g_{\widehat{K}_t} \circ G \circ g_{K_t}^{-1}$. Then G_t is analytic in $g_{K_t}(D \setminus K_t)$ and depends continuously on t . Hence $G'_t(U_t)$ is continuous in t . Since $A(G(K_{t+u})) = A(G(K_t)) + A(G_t(K_{t,u}))$, it follows that $\partial_t A(G(K_t)) = G'_t(U_t)^2 \partial_t A(K_t)$, which completes the proof. \square

For future reference, we note that when $g_t = g_{K_t}$, satisfies the differential equation (2.6), we have the formula

$$\partial_t \log g'_t(z) = - \frac{2\partial_t A(g_t)}{(g_t(z) - U_t)^2}, \tag{2.9}$$

which is obtained by differentiating (2.6) with respect to z .

2.3.3. *Pairs of time-modified expanding hulls.* We now discuss the situation where there are two disjoint expanding hulls.

Let $(L_s, s \in [0, s_0])$ and $(K_t, t \in [0, t_0])$ be a pair of time-modified expanding hulls such that $\bar{L}_{s_0} \cap \bar{K}_{t_0} = \emptyset$. Let $g_{s,t} := g_{L_s \cup K_t}$, $g_t := g_{K_t}$, $\hat{g}_s := g_{L_s}$ and $a(s, t) := A(g_{s,t})$. Then for each $s \in [0, s_0]$ and $t \in [0, t_0]$ we have

$$g_{s,t} = g_{g_t(L_s)} \circ g_t.$$

Therefore,

$$\partial_s g_{s,t}(z) = \frac{2\partial_s a(s,t)}{g_{s,t}(z) - U^1(s,t)},$$

where $s \mapsto U^1(s,t)$ is the driving function for the time-modified expanding hulls $s \mapsto g_t(L_s)$. Similarly,

$$\partial_t g_{s,t}(z) = \frac{2\partial_t a(s,t)}{g_{s,t}(z) - U^2(s,t)},$$

where $t \mapsto U^2(s,t)$ is the driving function for the time-modified expanding hulls $t \mapsto \hat{g}_s(K_t)$. Although we do not know that $g_t^{-1}(U^2(0,t))$ is well defined, $g_{s,t} \circ g_t^{-1}$ is analytic in a neighborhood of $U^2(0,t)$, by the reflection principle. Hence, it is clear that $U^2(s,t) = g_{s,t} \circ g_t^{-1}(U^2(0,t))$ (see, for example, the construction of U_t in the proof of Theorem 2.6), and therefore

$$\partial_s U^2(s,t) = \frac{2\partial_s a(s,t)}{U^2(s,t) - U^1(s,t)}. \quad (2.10)$$

We will now prove the formula

$$\partial_s \partial_t a(s,t) = \frac{-4\partial_s a(s,t)\partial_t a(s,t)}{(U^2(s,t) - U^1(s,t))^2}. \quad (2.11)$$

From (2.7) we have

$$\partial_t a(0,s) = \hat{g}'_s(U^2(0,0))^2 \partial_t a(0,0) = \hat{g}'_s(U^2(0,0))^2,$$

and using (2.9), we obtain

$$\partial_s \log \partial_t a(0,s) = \frac{-4\partial_s A(\hat{g}_s)}{(\hat{g}_s(U^2(0,0)) - U^1(s,0))^2} = \frac{-4\partial_s a(0,s)}{(U^2(s,0) - U^1(s,0))^2}.$$

This verifies (2.11) for the case $t=0$. The general case is similarly obtained.

2.4. Proof of Lemma 2.5

We will now prove Lemma 2.5; this is the core of the proof of the locality property. We assume that $\gamma: [0, s_1] \rightarrow \overline{\mathbf{H}}$ is a continuous simple path with $\gamma(0) \in \mathbf{R} \setminus \{0\}$ and $L_s := \gamma(0, s_1] \subset \mathbf{H}$. With no loss of generality, assume that γ is parameterized so that $A(L_s) = s$. By Theorem 2.6, $(L_s, s \in [0, s_1])$ is a time-modified expanding hull, and by Lemma 2.8, for each s , $t \mapsto g_{L_s}(K_t)$ is a time-modified expanding hull, and for each t , $s \mapsto g_{K_t}(L_s)$ is a time-modified expanding hull. Let $t \mapsto W(s, t)$ be the process driving $t \mapsto g_{L_s}(K_t)$, let $U(s, t)$ be the process driving $s \mapsto g_{K_t}(L_s)$, and let $Y_t = W(0, t)$ be the process driving K_t . As above, let $a(s, t) = A(g_{s,t})$. For simplicity, $W(s, t)$ will be abbreviated to W , $U(s, t)$ to U , $a(s, t)$ to a , etc.

Our aim is to show that $(W(s_1, t), t \geq 0)$ is a continuous martingale (up to the stopping time T), and that its quadratic variation (for background on stochastic calculus, see, e.g., [15], [39]) is

$$\langle W(s_1, \cdot) \rangle_t = 6(a(s_1, t) - a(s_1, 0)).$$

Indeed, if this is true, let $\phi(t)$ be the inverse of the map $t \mapsto a(s_1, t) - a(s_1, 0)$, and define $\widetilde{W}(t) = W(s_1, \phi(t))$. Then $\widetilde{W}(\frac{1}{6}t)$ is a Brownian motion, so that $t \mapsto g_{s_1, 0}(K_{\phi(t)}) - g_{s_1, 0}(0)$ is an SLE_6 -process, as required. Note that this will in fact give a precise expression for the time change in Lemma 2.5 and Theorem 2.2.

Before giving the mathematically rigorous proof, we first present a formal, non-rigorous derivation of the fact that $W(s, \cdot)$ is a martingale. In this derivation, κ will be kept as a variable, in order to stress where the assumption $\kappa = 6$ plays a role (it will not be so apparent in our proof).

Non-rigorous argument. The first goal is to show that the quadratic variation $\langle W \rangle_t$ of $t \mapsto W(s, t)$ satisfies

$$\partial_t \langle W \rangle_t = \kappa \partial_t a, \tag{2.12}$$

for each s, t . It is clear that this holds when $s = 0$, since K_t is SLE_6 . We have

$$\begin{aligned} \partial_s \partial_t \langle W \rangle_t &= \partial_t \partial_s \langle W \rangle_t \\ &= 2 \partial_t \langle \partial_s W, W \rangle_t \\ &= 2 \partial_t \langle 2(\partial_s a)(W - U)^{-1}, W \rangle_t \quad (\text{by (2.10)}) \\ &= -4(\partial_s a)(W - U)^{-2} \partial_t \langle W \rangle_t \quad (\text{by It\^o's formula}) \\ &= (\partial_t a)^{-1} (\partial_s \partial_t a) \partial_t \langle W \rangle_t \quad (\text{by (2.11)}). \end{aligned}$$

Consequently,

$$(\partial_t a)^2 \partial_s (\partial_t \langle W \rangle_t / \partial_t a) = \partial_t a \partial_s \partial_t \langle W \rangle_t - \partial_t \langle W \rangle_t \partial_s \partial_t a = 0,$$

which means that $\partial_t \langle W \rangle_t / \partial_t a$ does not depend on s . Since (2.12) holds when $s=0$, this proves (2.12).

We now show that $t \mapsto W(s, t)$ is a martingale. The dt -term in Itô's formula for the ∂_t -derivative of

$$\partial_s W(s, t) = \frac{2\partial_s a}{W-U}$$

is

$$\frac{2\partial_t \partial_s a}{W-U} + 2 \frac{\partial_s a}{(W-U)^3} \partial_t \langle W \rangle_t - 4 \frac{\partial_s a}{(W-U)^3} \partial_t a,$$

where the first summand comes from differentiating $\partial_s a$, the second summand is the diffusion term in Itô's formula, and the last summand comes from differentiating with respect to U and using (2.10) for $\partial_t U$. Using (2.11) and (2.12), this becomes

$$(3 - \frac{1}{2}\kappa) \frac{\partial_t \partial_s a}{W-U},$$

which vanishes when $\kappa=6$. Hence $t \mapsto \partial_s W(s, t)$ is a martingale. As

$$W(s, t) = Y_t + \int_0^s \partial_s W(s', t) ds',$$

it follows that $t \mapsto W(s, t)$ is a martingale. This completes the informal proof.

The problem with the above argument is that we do not know that $t \mapsto W(s, t)$ is a semi-martingale, and hence cannot apply stochastic calculus to it. Moreover, we need to check that there is sufficient regularity to justify the equality $\partial_s \partial_t \langle W \rangle_t = \partial_t \partial_s \langle W \rangle_t$.

To rectify the situation, set

$$V(s, t') := W(s, 0) + \int_0^{t'} \sqrt{\partial_t a(s, t)} dY_t.$$

Then $t \mapsto V(s, t)$ is clearly a martingale. The rest of this subsection will be devoted to the proof of the fact that $V=W$. Recall that

$$T = \sup\{t \geq 0 : \bar{K}_t \cap \bar{L}_{s_1} = \emptyset\}.$$

We will need the following fact:

LEMMA 2.9. *There exists a continuous version of V on $[0, s_1] \times [0, T]$.*

Proof. In order to keep some quantities bounded, we have to stop the processes slightly before T . Let us fix $\varepsilon \in (0, 1)$, and define

$$T_1^\varepsilon = \inf\{t > 0 : \inf_{s \leq s_1} |W(s, t) - U(s, t)| \leq \varepsilon\}.$$

Define for any $s \leq s_1$ and $t_0 \geq 0$

$$\tilde{V}(s, t_0) = \tilde{V}^\varepsilon(s, t_0) := \int_0^{\min(t_0, T_1^\varepsilon)} \sqrt{\partial_t a} dY_t.$$

As

$$\sup_{n \geq 1} T_1^{1/n} = T,$$

it is sufficient to show existence of a continuous version (on $[0, s_1] \times \mathbf{R}_+$) of \tilde{V} .

Let

$$\tilde{a} = \tilde{a}(s, t) := 1_{\{t \leq T_1^\varepsilon\}} \sqrt{\partial_t a}.$$

Note that $\partial_t a(0, t) = \partial_s a(s, 0) = 1$. Hence, from (2.11) it follows that $\partial_s a \leq 1$ and $\partial_t a \leq 1$ for all $s \leq s_1, t < T$. Using (2.11) again, we get

$$|\partial_s \tilde{a}| = 1_{\{t \leq T_1^\varepsilon\}} \frac{2\sqrt{\partial_t a} \partial_s a}{(W-U)^2} \leq 2\varepsilon^{-2}.$$

Hence, for all $t \geq 0$, for all s, s' in $[0, s_1]$,

$$|\tilde{a}(s, t) - \tilde{a}(s', t)| \leq 2\varepsilon^{-2} |s - s'|.$$

But

$$\mathbf{E}[(\tilde{V}(s, t_0) - \tilde{V}(s', t'_0))^4] \leq 16\mathbf{E}\left[\left(\int_{t_0}^{t'_0} \tilde{a}(s', t) dY_t\right)^4\right] + 16\mathbf{E}\left[\left(\int_0^{t_0} (\tilde{a}(s, t) - \tilde{a}(s', t)) dY_t\right)^4\right],$$

and using, for instance, the Burkholder–Davis–Gundy inequality for $p=4$ (see, e.g., [39, IV.4]), we see that there exists a constant $c_1 = c_1(\varepsilon)$ such that for all $t_0, t'_0 \geq 0$ and $s, s' \in [0, s_1]$,

$$\begin{aligned} \mathbf{E}[(\tilde{V}(s, t_0) - \tilde{V}(s', t'_0))^4] &\leq c_1 \mathbf{E}[(t_0 - t'_0)^2] + c_1 \mathbf{E}\left[\left(\int_0^{t_0} (s - s')^2 dt\right)^2\right] \\ &\leq c_1(t_0 - t'_0)^2 + c_1 t_0^2 (s - s')^4, \end{aligned}$$

and the existence of a continuous version of \tilde{V} then easily follows from Kolmogorov’s lemma (see, e.g., [39, I.(1.8)]). □

From now on, we will use a version of V that is continuous on $[0, s_1] \times [0, T)$. Define

$$\begin{aligned} T_2^\varepsilon &:= \inf \{t \geq 0 : \sup_{s \leq s_1} |V(s, t) - W(s, t)| \geq 1\}, \\ T_3^\varepsilon &:= \inf \{t \geq 0 : \inf_{s \leq s_1} |V(s, t) - U(s, t)| \leq \varepsilon\}, \\ T^\varepsilon &:= \min(T_1^\varepsilon, T_2^\varepsilon, T_3^\varepsilon). \end{aligned}$$

Note that for all $s \leq s_1$ and $t < T$,

$$\partial_s \sqrt{\partial_t a} = \frac{-2\sqrt{\partial_t a} \partial_s a}{(W-U)^2}.$$

The process $\partial_s \sqrt{\partial_t a}$ remains bounded before T^ε (uniformly in $s \leq s_1$), and it is a measurable function of (s, t) . By Fubini's theorem for stochastic integrals ([15, Lemma III.4.1]), we have that for all $s_0 \leq s_1$, for all $t_0 \geq 0$, almost surely

$$\begin{aligned} \int_0^{s_0} \int_0^{t'_0} \frac{-2\sqrt{\partial_t a} \partial_s a}{(W-U)^2} dY_t ds &= \int_0^{t'_0} \left(\int_0^{s_0} \partial_s \sqrt{\partial_t a} ds \right) dY_t \\ &= \int_0^{t'_0} (\sqrt{\partial_t a}(s_0, t) - \sqrt{\partial_t a}(0, t)) dY_t \\ &= V(s_0, t'_0) - V(0, t'_0) - (W(s_0, 0) - W(0, 0)), \end{aligned} \quad (2.13)$$

where $t'_0 := \min(t_0, T^\varepsilon)$. On the other hand, using Itô's formula, we now compute

$$\begin{aligned} \frac{2\partial_s a(s, t'_0)}{U(s, t'_0) - V(s, t'_0)} &= \frac{2}{U(s, 0) - V(s, 0)} + \int_0^{t'_0} \frac{2\partial_s a \sqrt{\partial_t a}}{(U-V)^2} dY_t \\ &\quad + 2 \int_0^{t'_0} \left(\frac{\partial_t \partial_s a}{U-V} - \frac{\partial_s a \partial_t U}{(U-V)^2} + \frac{\partial_s a \partial_t \langle V \rangle_t}{(U-V)^3} \right) dt \\ &= -\partial_s W(s, 0) + \int_0^{t'_0} \tilde{b}_1 dY_t + \int_0^{t'_0} (V-W)b_2 dt, \end{aligned} \quad (2.14)$$

where (using $\partial_t \langle V \rangle_t = \partial_t a \partial_t \langle Y \rangle_t = \kappa \partial_t a = 6\partial_t a$)

$$\begin{aligned} \tilde{b}_1(s, t) &:= \frac{2\partial_s a \sqrt{\partial_t a}}{(U-V)^2}, \\ b_2(s, t) &:= \frac{4\partial_s a \partial_t a}{(U-W)^2 (U-V)^2} \left(5 + 3 \frac{W-V}{V-U} \right). \end{aligned}$$

Note that for all $s \leq s_1$ and $t \leq T^\varepsilon$,

$$|b_2(s, t)| \leq 16\varepsilon^{-5}.$$

By integrating (2.14) with respect to s and subtracting (2.13) from it, we get

$$\begin{aligned} V(s_0, t'_0) - V(0, t'_0) &+ \int_0^{s_0} \frac{2\partial_s a(s, t'_0)}{U(s, t'_0) - V(s, t'_0)} ds \\ &= \int_0^{s_0} \int_0^{t'_0} (V-W)b_2(s, t) dt ds \\ &\quad + \int_0^{s_0} \left(\int_0^{t'_0} \frac{-2\sqrt{\partial_t a} \partial_s a}{(W-U)^2} dY_t + \int_0^{t'_0} \tilde{b}_1(s, t) dY_t \right) ds \\ &= \int_0^{s_0} \int_0^{t'_0} (V-W)b_2 dt ds + \int_0^{s_0} \int_0^{t'_0} (V-W)b_1 dY_t ds, \end{aligned} \quad (2.15)$$

where (after some simplifications)

$$b_1(s, t) := \frac{2\partial_s a \sqrt{\partial_t a}}{(V-U)^2(W-U)^2} ((U-W) + (U-V)).$$

Note that for all $s \in [0, s_1]$ and $t \leq T^\epsilon$,

$$|b_1(s, t)| \leq 4\epsilon^{-3}.$$

But we know on the other hand that

$$W(s_0, t'_0) - W(0, t'_0) = \int_0^{s_0} \frac{2\partial_s a(s, t'_0)}{W(s, t'_0) - U(s, t'_0)} ds. \tag{2.16}$$

Subtracting this equation from (2.15), one gets

$$\begin{aligned} V(s_0, t'_0) - W(s_0, t'_0) &= \int_0^{s_0} b_3(s, t'_0)(V(s, t'_0) - W(s, t'_0)) ds \\ &\quad + \int_0^{s_0} \int_0^{t'_0} (V-W)b_2 dt ds + \int_0^{s_0} \int_0^{t'_0} (V-W)b_1 dY_t ds, \end{aligned}$$

where

$$b_3(s, t) := \frac{-2\partial_s a}{(U-W)(U-V)}.$$

Again b_3 remains uniformly bounded before T^ϵ .

We now define

$$H(s, t) = V(s, t) - W(s, t).$$

Hence, for all $t_0 \leq T$,

$$H(s_0, t_0) = \int_0^{s_0} b_3(s, t_0)H(s, t_0) ds + \int_0^{s_0} \int_0^{t_0} b_2 H ds dt + \int_0^{s_0} \int_0^{t_0} b_1 H ds dY_t$$

and $|b_1|, |b_2|, |b_3|$ are all bounded by some constant $c_2 = c_2(\epsilon)$ on $[0, s_1] \times [0, T^\epsilon]$. This equation and an argument similar to Gronwall's lemma will show that $H = 0$.

Let us fix $t_1 > 0$. For any $t \geq 0$, define

$$\tau(t) = \min(t, t_1, T^\epsilon).$$

We will use the notation $\tau_0 = \tau(t_0)$. It is easy to see that there exists a $c_3 = c_3(\epsilon, t_1, s_1)$ such that for all $t_0 \geq 0$ and $s_0 \in [0, s_1]$,

$$H(s_0, \tau_0)^2 \leq c_3 \int_0^{s_0} H(s, \tau_0)^2 ds + c_3 \int_0^{\tau_0} \int_0^{s_0} H^2 ds dt + c_3 \left(\int_0^{\tau_0} \int_0^{s_0} b_1 H ds dY_t \right)^2.$$

The Burkholder–Davis–Gundy inequality for $p=2$ (see, e.g., [39, IV.4]) shows that there exist constants $c_4=c_4(\varepsilon, t_1, s_1)$ and $c_5=c_5(\varepsilon, t_1, s_1)$ such that for all $s \in [0, s_1]$ and $t_0 \geq 0$,

$$\begin{aligned} \mathbf{E} \left[\sup_{u \leq t_0} \left(\int_0^{\tau(u)} \int_0^{s_0} b_1 H \, ds \, dY_t \right)^2 \right] &\leq c_4 \mathbf{E} \left[\int_0^{\tau_0} \left(\int_0^{s_0} b_1 H \, ds \right)^2 dt \right] \\ &\leq c_5 \int_0^{t_0} \int_0^{s_0} \mathbf{E}[H(s, \tau(t))^2] \, ds \, dt. \end{aligned}$$

Let us now define

$$h(s_0, t_0) = \mathbf{E} \left[\sup_{t \leq t_0} H(s_0, \tau(t))^2 \right].$$

Then

$$h(s_0, t_0) \leq c_6 \left(\int_0^{s_0} h(s, t_0) \, ds + \int_0^{t_0} \int_0^{s_0} h(s, t) \, ds \, dt \right).$$

We also know that $h(s, t)$ is bounded by 1 (because $|H| \leq 1$). Hence, it is straightforward to prove by induction that for all $s_0 \in [0, s_1]$, $t_0 \geq 0$ and $p=1, 2, \dots$,

$$h(s_0, t_0) \leq \frac{c_6^p s_0^p (1+t_0)^p}{p!},$$

so that $h(s_0, t_0)=0$. In particular (using the continuity of V and W), this shows that $W=V$ almost surely on all sets $[0, s_1] \times [0, \min(t_1, T^\varepsilon)]$. As this is true for all ε and t_1 , we conclude that $V=W$ on $[0, s_1] \times [0, T)$. Lemma 2.5 follows, and thereby also Theorem 2.2. \square

3. Exponents for SLE_6

3.1. Statement

In the present section, we are going to compute intersection exponents associated with SLE_6 .

Suppose that $D \subset \mathbf{C}$ is a Jordan domain; that is, ∂D is a simple closed curve in \mathbf{C} . Let $a, b \in \partial D$ be two distinct points on the boundary of D . As explained in §2.1, the $\text{SLE}_6(K_t, t \geq 0)$ from a to b in D is well defined, up to a linear time change.

Now suppose that $I \subset \partial D$ is an arc with $b \in I$ but $a \notin I$. Let

$$\tau_I := \sup\{t \geq 0 : \bar{K}_t \cap I = \emptyset\}.$$

By Corollary 2.3, up to a time change, the law of the process $(K_t, t < \tau_I)$ does not change if we replace b by another point $b' \in I$. Set

$$S = S(a, I, D) := \bigcup_{t < \tau_I} K_t,$$

and call this set the *hull from a to I in D*. It does not depend on b .

Suppose that $L > 0$, and let $\mathcal{R} = \mathcal{R}(L)$ denote the rectangle with corners

$$A_1 := 0, \quad A_2 := L, \quad A_3 := L + i\pi, \quad A_4 := i\pi. \tag{3.1}$$

Let \mathcal{S} denote the closure of the hull from A_4 to $[A_1, A_2] \cup [A_2, A_3]$ in \mathcal{R} .

In the following, we will use the terminology π -*extremal distance* instead of “ π times the extremal distance”. For instance, the π -extremal distance between the vertical sides of \mathcal{R} in \mathcal{R} is L .

When $\mathcal{S} \cap [A_1, A_2] = \emptyset$, let \mathcal{L} be the π -extremal distance between $[A_1, A_4]$ and $[A_2, A_3]$ in $\mathcal{R} \setminus \mathcal{S}$. Otherwise, put $\mathcal{L} = \infty$.

In the sequel, we will use the function

$$u(\lambda) = \frac{1}{6} (6\lambda + 1 + \sqrt{24\lambda + 1}). \tag{3.2}$$

The main goal of this section is to prove the following result.

THEOREM 3.1.

$$\mathbf{E}[1_{\{\mathcal{L} < \infty\}} \exp(-\lambda \mathcal{L})] = \exp(-u(\lambda)L + O(1)(\lambda + 1)) \quad \text{as } L \rightarrow \infty, \tag{3.3}$$

for any $\lambda \geq 0$ (where $O(1)$ denotes an arbitrary quantity whose absolute value is bounded by a constant which does not depend on L or λ).

In particular, when $\lambda = 0$,

$$\mathbf{P}[\mathcal{S} \cap [A_1, A_2] = \emptyset] = \mathbf{P}[\mathcal{L} < \infty] = \exp(-\frac{1}{3}L + O(1)) \quad \text{as } L \rightarrow \infty. \tag{3.4}$$

3.2. Generalized Cardy’s formula

By conformal invariance, we may work in the half-plane \mathbf{H} . Map the rectangle \mathcal{R} conformally onto \mathbf{H} so that A_1 is mapped to 1, A_2 is mapped to ∞ , A_3 is mapped to 0, and then the image $x = x(L) \in (0, 1)$ of A_4 is determined for us. Let K_t be the hull of an SLE₆-process $g_t = g_{K_t}$ in \mathbf{H} , with driving process $W(t)$, which is started at $W(0) = x$ (that is, K_t is a translation by x of the standard SLE₆ starting at 0). In order to emphasize the dependence on x , we will use the notation \mathbf{P}_x and \mathbf{E}_x for probability and expectation.

Set

$$\begin{aligned} T_0 &:= \sup\{t \geq 0 : \bar{K}_t \cap (-\infty, 0] = \emptyset\}, \\ T_1 &:= \sup\{t \geq 0 : \bar{K}_t \cap [1, \infty) = \emptyset\}, \\ T &:= \min\{T_0, T_1\}. \end{aligned}$$

As will be demonstrated, $T_0, T_1 < \infty$ a.s. Let

$$f_t(z) := \frac{g_t(z) - g_t(0)}{g_t(1) - g_t(0)},$$

for $t < T$, which is just g_t renormalized to fix 0, 1 and ∞ . It turns out that $f_T := \lim_{t \nearrow T} f_t$ exists a.s. On the event $T_1 > T_0$, f_T uniformizes the quadrilateral

$$(\mathbf{H} \setminus K_T; 1, \infty, \min(\bar{K}_T \cap \mathbf{R}), \max(\bar{K}_T \cap \mathbf{R}))$$

to the form

$$(\mathbf{H}; 1, \infty, 0, f_T(\max(\bar{K}_T \cap \mathbf{R}))).$$

Therefore, we want to know the distribution of

$$1 - f_T(\max(\bar{K}_T \cap \mathbf{R})),$$

and how it depends on x (especially when $x \in (0, 1)$ is close to 1). In this subsection, we will calculate something very closely related: the distribution of $f'_T(1)$ and how it depends on x .

Set

$$\Lambda(1-x, b) := \mathbf{E}_x[1_{\{T_0 < T_1\}} f'_T(1)^b]$$

for $b \geq 0$ and $x \in (0, 1)$. Recall the definition of the hypergeometric function ${}_2F_1$ (see, e.g., [30]):

$${}_2F_1(a_0, a_1, a_2; x) = \sum_{n=0}^{\infty} \frac{(a_0)_n (a_1)_n}{(a_2)_n n!} x^n,$$

where $(a)_n = \prod_{j=1}^n (a+j-1)$ and $(a)_0 = 1$. Note that ${}_2F_1(a_0, a_1, a_2; 0) = 1$.

THEOREM 3.2. *For all $b \geq 0$, $x \in (0, 1)$,*

$$\Lambda(x, b) = \frac{\sqrt{\pi} 2^{-2\hat{b}} \Gamma(\frac{5}{6} + \hat{b})}{\Gamma(\frac{1}{3}) \Gamma(1 + \hat{b})} x^{1/6 + \hat{b}} {}_2F_1(\frac{1}{6} + \hat{b}, \frac{1}{2} + \hat{b}, 1 + 2\hat{b}; x),$$

where

$$\hat{b} = \frac{1}{6} \sqrt{1 + 24b},$$

and ${}_2F_1$ is the hypergeometric function.

Setting $b=0$, we obtain Cardy's formula [7]. Thus, this result can be thought of as a generalization of Cardy's formula.

Note that Theorem 3.2 determines completely the law of $1_{\{T_0 < T_1\}} f'_T(1)$. In particular, the Laplace transform of the conditional law of $\log 1/f'_T(1)$ given $\{T_0 < T_1\}$ is $\Lambda(1-x, b)/\Lambda(1-x, 0)$.

Proof. We first observe that $T_1 < \infty$ almost surely. Indeed, $g_t(1) - W(t)$ is a Bessel process with index $\frac{5}{3}$, with time linearly scaled, and hence hits 0 almost surely in finite time (e.g. [39]). Similarly, $g_t(0) - W(t)$ hits 0 almost surely. It is clear that T_1 is the time t when $g_t(1) - W(t)$ hits 0, and T_0 is the time when $g_t(0) - W(t)$ hits 0. It follows from Theorem 2.6 and $x \in (0, 1)$ that almost surely $T_0 \neq T_1$. We may also conclude that

$$\lim_{x \rightarrow 0} \mathbf{P}_x[T_0 < T_1] = 1, \quad \lim_{x \rightarrow 1} \mathbf{P}_x[T_0 < T_1] = 0. \tag{3.5}$$

The next goal is to prove that

$$\lim_{t \nearrow T} f'_t(1) > 0 \quad \text{if and only if} \quad T_0 < T_1. \tag{3.6}$$

If $T_0 < T_1$, then $\bar{K}_T \cap [1, \infty) = \emptyset$. Therefore f_T is defined and conformal near 1, and $f'_T(1) > 0$, by the reflection principle. On the other hand, if $T_1 < T_0$, then $\bar{K}_T \cap [1, \infty) \neq \emptyset$. We claim that

$$1 \notin \bar{K}_{T_1} \quad \text{almost surely}; \tag{3.7}$$

since $\bar{K}_{T_1} \cap [1, \infty) \neq \emptyset$, this means that K_{T_1} separates 1 from ∞ in \mathbf{H} . Indeed, let $\phi: \mathbf{H} \rightarrow \mathbf{H}$ be the anti-conformal automorphism that fixes x and exchanges 1 and ∞ . $T_1 < \infty$ a.s. and $\sup_{t < T_1} |W(t)| < \infty$ a.s. imply that K_{T_1} is bounded a.s., which is the same as saying that $\phi(K_t)$ stays bounded away from 1 as $t \nearrow T_1$. But Corollary 2.3 and invariance under reflection imply that up to time T_1 the law of K_t is the same as a time change of the law of $\phi(K_t)$. Hence, a.s. K_t stays bounded away from 1 as $t \nearrow T_1$, proving (3.7). It follows from (3.7) that $\lim_{t \nearrow T_1} f'_t(1) = 0$ a.s. on the event $T_1 < T_0$ (observe that, given (3.7), the extremal length from a neighborhood of 0 to a neighborhood of 1 in $\mathbf{H} \setminus K_t$ tends to ∞ as $t \nearrow T_1$), and (3.6) is established.

Define the renormalized version of $W(t)$,

$$Z(t) := \frac{W(t) - g_t(0)}{g_t(1) - g_t(0)},$$

and the new time parameter

$$s = s(t) := \int_0^t \frac{dt}{(g_t(1) - g_t(0))^2}, \quad t < T.$$

Set $s_0 := \lim_{t \nearrow T} s(t)$. Since $T_1 \neq T_0$ a.s., $\inf\{g_t(1) - g_t(0) : t < T\} > 0$ a.s., and hence $s_0 < \infty$ a.s. Let $t(s)$ denote the inverse to the map $t \mapsto s(t)$. A direct calculation gives

$$\partial_s(f_{t(s)}(z)) = \frac{-2}{Z(t) - f_t(z)} + \frac{2(1 - f_t(z))}{Z(t)} - \frac{2f_t(z)}{1 - Z(t)}$$

and

$$dZ(t) = \frac{dW_t}{g_t(1) - g_t(0)} + \frac{2 dt}{(g_t(1) - g_t(0))^2} \left(\frac{1}{Z(t)} + \frac{1}{Z(t) - 1} \right).$$

We now use the notation

$$\tilde{Z}(s) := Z(t(s)), \quad \tilde{f}_s(z) := f_{t(s)}(z).$$

Then,

$$d\tilde{Z}_s = dX_s + \frac{2(1 - 2\tilde{Z}(s)) ds}{\tilde{Z}(s)(1 - \tilde{Z}(s))} = dX_s + \left(\frac{2}{\tilde{Z}(s)} - \frac{2}{1 - \tilde{Z}(s)} \right) ds, \quad (3.8)$$

where $(X_s, s \geq 0)$ has the same law as $(W(t), t \geq 0)$; i.e., it is a Brownian motion with time rescaled by a factor of 6. Also,

$$\partial_s(\tilde{f}_s(z)) = \frac{-2}{\tilde{Z}(s) - \tilde{f}_s(z)} + \frac{2(1 - \tilde{f}_s(z))}{\tilde{Z}(s)} - \frac{2\tilde{f}_s(z)}{1 - \tilde{Z}(s)}. \quad (3.9)$$

These two equations describe the evolution of $\tilde{f}_s(z)$. Note that $s(T) = s_0$ is the first time at which $\tilde{Z}(s)$ hits 0 or 1.

We now assume that $b > 0$. Differentiating (3.9) with respect to z gives

$$\partial_s(\log \tilde{f}'_s(z)) = \frac{-2}{(\tilde{Z}(s) - \tilde{f}_s(z))^2} - \frac{2}{\tilde{Z}(s)} - \frac{2}{1 - \tilde{Z}(s)} \quad (3.10)$$

(the Cauchy integral formula, for example, shows that we may indeed differentiate, but this is also legitimate since ∂_s and ∂_z commute in this case). We are particularly interested in

$$\alpha(s) := \log \tilde{f}'_s(1) = \log f'_{t(s)}(1),$$

which satisfies

$$\partial_s \alpha(s) = \frac{-2}{(\tilde{Z}(s) - 1)^2} - \frac{2}{\tilde{Z}(s)} - \frac{2}{1 - \tilde{Z}(s)}. \quad (3.11)$$

Note that the equations (3.8) and (3.11) describe the evolution of the Markov process $(\tilde{Z}(s), \alpha(s))$. The process stops at s_0 . Define

$$y(x, v) := \mathbf{E}[\exp(b\alpha(s_0)) \mid \tilde{Z}(0) = x, \alpha(0) = v],$$

where the expectation corresponds to the Markov process started from $\tilde{Z}(0) = x$ and $\alpha(0) = v$. From the definition of y it follows that

$$y(x, 0) = \Lambda(1 - x, b),$$

since $\lim_{t \nearrow T} f_t = f_T$ in a neighborhood of 1 on the event $T_0 < T_1$ and (3.6) holds. It is standard that such a function $y(x, v)$ is C^∞ , and the strong Markov property ensures that the process

$$Y = y(\tilde{Z}(s), \alpha(s))$$

is a local martingale. The drift term in Itô's formula for dY must vanish, which gives

$$0 = \frac{2(1-2x)}{x(1-x)} \partial_x y + 3 \partial_{xx}^2 y + \left(\frac{-2}{(1-x)^2} - \frac{2}{x} - \frac{2}{1-x} \right) \partial_v y. \quad (3.12)$$

As

$$\alpha(s) = \alpha(0) + \int_0^s (\partial_s \alpha(s')) ds',$$

we get

$$y(x, v) = \exp(bv) y(x, 0).$$

Set

$$h(x) := y(1-x, 0) = \Lambda(x, b),$$

so that $y(x, v) = \exp(bv) h(1-x)$. Hence (3.12) becomes

$$-2bh(x) + 2x(1-2x)h'(x) + 3x^2(1-x)h''(x) = 0. \quad (3.13)$$

The second statement in (3.5) implies that

$$\lim_{x \searrow 0} h(x) = 0,$$

while

$$\lim_{x \nearrow 1} h(x) = 1$$

holds, since when x is close to 0, K_{T_0} is likely to be small, by scale invariance, for example. The differential equation (3.13) can be solved explicitly by looking for solutions of the type $h(x) = x^c z(x)$: two linearly independent solutions are ($i=1, 2$)

$$h_i(x) = x^{1/6+b_i} {}_2F_1\left(\frac{1}{6}+b_i, \frac{1}{2}+b_i, 1+2b_i; x\right),$$

where

$$b_1 = -b_2 = \frac{1}{6} \sqrt{1+24b}.$$

Recall that ${}_2F_1(a_0, a_1, a_2; 0) = 1$. The function $h(x)$ must be a linear combination of h_1 and h_2 . However, $\lim_{x \searrow 0} h(x) = 0 = \lim_{x \searrow 0} h_1(x)$, but $\lim_{x \searrow 0} h_2(x) = \infty$. Hence, $h(x) = ch_1(x)$ for some constant c . The equality $h(1) = 1$ and knowledge of the value at $x=1$ of

hypergeometric functions (see, e.g., [30]) allows the determination of c , and establishes the theorem in the case $b > 0$. The case $b = 0$ follows by taking a limit as $b \searrow 0$. \square

Remark. With the same proof, Theorem 3.2 generalizes to SLE_κ with $\kappa > 4$, and gives

$$\Lambda_\kappa(x, b) = C(b, \kappa) x^{1/2 - 2/\kappa + \hat{b}_\kappa} {}_2F_1\left(\frac{1}{2} - 2/\kappa + \hat{b}_\kappa, 6/\kappa - \frac{1}{2} + \hat{b}_\kappa, 1 + 2\hat{b}_\kappa; x\right),$$

where

$$\hat{b}_\kappa := \frac{\sqrt{(\kappa - 4)^2 + 16\kappa b}}{2\kappa}, \quad C(b, \kappa) := \frac{\Gamma(\frac{3}{2} - 6/\kappa + \hat{b}_\kappa) \Gamma(\frac{1}{2} + 2/\kappa + \hat{b}_\kappa)}{\Gamma(1 - 4/\kappa) \Gamma(1 + 2\hat{b}_\kappa)}.$$

3.3. Determination of the SLE_6 -exponents

For every $t \geq 0$, set

$$M_t := \max(\bar{K}_t \cap \mathbf{R}).$$

The following lemma shows that our understanding of the derivative $f'_T(1)$ gives information on $f_T(M_T)$ itself.

LEMMA 3.3. *In the above setting, let $N_T := f_T(M_T)$. For $b \geq 0$, set*

$$\Theta(x, b) := E_{1-x}[1_{\{T_0 < T_1\}}(1 - N_T)^b].$$

Then

$$\left(\frac{1}{2}x\right)^b \Lambda\left(\frac{1}{2}x, b\right) \leq \Theta(x, b) \leq x^b \Lambda(x, b). \tag{3.14}$$

Note that $\Theta(x, b)$ is close to the quantity we are after, since $-\log(1 - N_T)$ is approximately the extremal length of the quadrilateral

$$(\mathbf{H} \setminus K_T; \min(\bar{K}_T \cap \mathbf{R}), \max(\bar{K}_T \cap \mathbf{R}), 1, \infty).$$

Proof. It follows easily from (3.10) that $f'_t(z)$ is non-decreasing in z (viewed as a real variable), as long as $z \geq M_t$. Therefore,

$$1 - N_T = \int_{M_T}^1 f'_T(z) dz \leq (1 - M_T) f'_T(1) \leq x f'_T(1).$$

This gives the right-hand inequality in (3.14).

To get the other inequality, consider some fixed $x^* > 1$ (x^* should be thought of as close to 1; we will eventually take $x^* = 1/(1 - x)$). Let $\tilde{x}^* = f_T(x^*)$. Then

$$\tilde{x}^* - N_T \geq \tilde{x}^* - 1 = \int_1^{x^*} f'_T(z) dz \geq (x^* - 1) f'_T(1).$$

This gives

$$E_{1-x}[1_{\{T_0 < T_1\}}(\tilde{x}^* - N_T)^b] \geq (x^* - 1)^b \Lambda(x, b). \tag{3.15}$$

A simple scaling argument will give an upper bound of the left-hand side of this inequality in terms of Θ . Let

$$T^* = \inf\{t \geq 0 : \bar{K}_t \cap \mathbf{R} \setminus (0, x^*) \neq \emptyset\}.$$

Note that $T \leq T^* < \infty$ a.s. and that $T = T^*$ if $T_0 < T_1$. For each $t \leq T^*$, let f_t^* be the conformal map from the unbounded component of $\mathbf{H} \setminus K_t$ to \mathbf{H} , which fixes the points $\infty, 0, x^*$. For all $t \leq T$,

$$f_t^*(z) = \frac{x^*}{f_t(x^*)} f_t(z).$$

Note that

$$f_t(x^*) \leq x^* = f_t^*(x^*).$$

Then,

$$1_{\{T_0 < T_1\}}(\tilde{x}^* - N_T) \leq 1_{\{T_0 < T_1\}}(x^* - f_T^*(M_T)).$$

Hence,

$$1_{\{T_0 < T_1\}}(\tilde{x}^* - N_T) \leq 1_{\{M_{T^*} < x^*\}}(x^* - f_{T^*}^*(M_{T^*})), \tag{3.16}$$

since on the event $T_0 < T_1$, we have $T = T^*$ and $M_T = M_{T^*}$. However, by scale invariance, when $W(0) = 1 - x$ the random variable

$$1_{\{M_{T^*} < x^*\}}(x^* - f_{T^*}^*(M_{T^*}))$$

has the same law as the random variable

$$x^* 1_{\{M_T < 1\}}(1 - f_T(M_T))$$

does when $W(0) = (1 - x)/x^*$. Thus, combining (3.16) and (3.15) gives

$$(x^*)^b \Theta(1 - (1 - x)/x^*, b) \geq (x^* - 1)^b \Lambda(x, b).$$

We take $x^* = 1/(1 - x)$, say, and get

$$\Theta(2x, b) \geq \Theta(2x - x^2, b) \geq x^b \Lambda(x, b),$$

which gives the left-hand side of (3.14). □

Proof of Theorem 3.1. We are now ready to derive Theorem 3.1 by combining Theorem 3.2 and Lemma 3.3.

In the setting of the theorem, let $\phi: \mathcal{R}(L) \rightarrow \mathbf{H}$ be the conformal homeomorphism satisfying $\phi(A_1)=1$, $\phi(A_2)=\infty$ and $\phi(A_3)=0$. Set $x=x(L):=\phi(A_4)$. By conformal invariance, the law of \mathcal{L} is the same as that of the π -extremal distance \mathcal{L}^* from $(-\infty, 0]$ to $(N_T, 1)$ in \mathbf{H} (with the notations of Lemma 3.3). Considering the map $z \mapsto \log(z-1)$ makes it clear that

$$L = -\log(1-x) + O(1), \quad (3.17)$$

for $L > 1$, and similarly

$$\mathcal{L}^* = -\log(1-N_T) + O(1). \quad (3.18)$$

For $L > 1$ (note also that $\mathcal{L} \geq L$),

$$\begin{aligned} \mathbf{E}[1_{\{\mathcal{L} < \infty\}} \exp(-\lambda \mathcal{L})] &= \mathbf{E}[1_{\{\mathcal{L}^* < \infty\}} \exp(-\lambda \mathcal{L}^*)] \\ &= \exp(O(1)) \mathbf{E}[1_{\{N_T < 1\}} (1-N_T)^\lambda] \quad (\text{by (3.18)}) \\ &= \exp(O(1)) \Theta(1-x, \lambda) \\ &= \exp(O(\lambda+1)) (1-x)^{u(\lambda)} \quad (\text{by Lemma 3.3 and Theorem 3.2}) \\ &= \exp(O(\lambda+1)) \exp(-u(\lambda)L) \quad (\text{by (3.17)}), \end{aligned}$$

which completes the proof of Theorem 3.1. \square

4. The Brownian half-plane exponents

We are now ready to combine the results collected so far and a “universality” idea similar to that developed in [29] to compute the exact value of some Brownian intersection exponents in the half-plane.

4.1. Definitions and background

In this short subsection, we quickly review some results on intersection exponents between independent planar Brownian motions. For details and complete proofs of these results, see [28], [29].

Suppose that $n+p$ independent planar Brownian motions β^1, \dots, β^n and $\gamma^1, \dots, \gamma^p$ are started from points $\beta^1(0) = \dots = \beta^n(0) = 0$ and $\gamma^1(0) = \dots = \gamma^p(0) = 1$ in the complex plane, and consider the probability $f_{n,p}(t)$ that for all $j \leq n$ and $l \leq p$, the paths of β^j up to time t and of γ^l up to time t do not intersect; more precisely,

$$f_{n,p}(t) := \mathbf{P} \left[\left(\bigcup_{j=1}^n \beta^j[0, t] \right) \cap \left(\bigcup_{l=1}^p \gamma^l[0, t] \right) = \emptyset \right].$$

It is easy to see that as $t \rightarrow \infty$ this probability decays roughly like a power of t . The (n, p) -intersection exponent $\xi(n, p)$ is defined as twice this power, i.e.,

$$f_{n,p}(t) = (\sqrt{t})^{-\xi(n,p)+o(1)}, \quad t \rightarrow \infty.$$

We call $\xi(n, p)$ the intersection exponent between one packet of n Brownian motions and one packet of p Brownian motions (for a list of references on Brownian intersection exponents, see [28]). Note that the exponent ζ described in the introduction is $\frac{1}{2}\xi(1, 1)$. It turns out to be more convenient to use this definition as a power of \sqrt{t} , i.e., of the space parameter. A Brownian motion travels very roughly to distance \sqrt{t} in time t : recall that if β is a planar Brownian motion started from 0, say, and T_R denotes its hitting time of the circle of radius R about 0, then for all $\delta > 0$, the probability that $T_R \notin (R^{2-\delta}, R^{2+\delta})$ decays as $R \rightarrow \infty$ faster than any negative power of R . This facilitates an easy conversion between the time-based definition of intersection exponents and a definition where the particles die when they exit a large ball.

Similarly, one can define corresponding probabilities for intersection exponents in a half-plane,

$$\tilde{f}_{n,p}(t) := \mathbf{P}[\forall j \leq n, \forall l \leq p, \beta^j[0, t] \cap \gamma^l[0, t] = \emptyset \text{ and } \beta^j[0, t] \cup \gamma^l[0, t] \subset \mathcal{H}],$$

where \mathcal{H} is some half-plane containing the two starting points ($\tilde{f}_{n,p}(t)$ will depend on \mathcal{H}). In plain words, we are looking at the probability that all Brownian motions stay in the half-plane and that all β 's avoid all γ 's. It is also easy to see that there exists a $\tilde{\xi}(n, p)$ (which does not depend on \mathcal{H}) such that

$$\tilde{f}_{n,p}(t) = (\sqrt{t})^{-\tilde{\xi}(n,p)+o(1)}, \quad t \rightarrow \infty.$$

Note that $\tilde{\zeta}$ described in the introduction is $\frac{1}{2}\tilde{\xi}(1, 1)$.

One can also define intersection exponents $\xi(n_1, \dots, n_p)$ and $\tilde{\xi}(n_1, \dots, n_p)$ involving more packets of Brownian motions. (For a more detailed discussion of this, see [28]). For instance, if B^1, B^2, B^3, B^4 denote four Brownian motions started from different points, the exponent $\xi(2, 1, 1)$ is defined by

$$\begin{aligned} \mathbf{P}[\text{the three sets } B^1[0, t] \cup B^2[0, t], B^3[0, t], B^4[0, t] \text{ are disjoint}] \\ = t^{-\xi(2,1,1)/2+o(1)}, \quad t \rightarrow \infty. \end{aligned}$$

One of the results of [28] is that there is a natural and rigorous way to generalize the definition of intersection exponents between packets of Brownian motions to the case where each packet of Brownian motions is the union of a “non-integer number” of paths; for

the half-plane exponents, one can define the exponents $\tilde{\xi}(u_1, \dots, u_p)$, where $u_1, \dots, u_p \geq 0$. These generalized exponents satisfy the so-called cascade relations (see [28]): for any $1 \leq q \leq p-1$,

$$\tilde{\xi}(u_0, \dots, u_p) = \tilde{\xi}(u_0, \dots, u_{q-1}, \tilde{\xi}(u_q, \dots, u_p)). \tag{4.1}$$

Moreover, $\tilde{\xi}$ is invariant under a permutation of its arguments.

There exists (see [28], [29]) a characterization of these exponents in terms of the so-called Brownian excursions that turns out to be useful. For any bounded simply-connected open domain D , there exists a Brownian excursion measure μ_D in D . This is an infinite measure on paths $(B(t), t \leq \tau)$ in D such that $B(0, \tau) \subset D$ and $B(0), B(\tau) \in \partial D$ (these can be viewed as prime ends if necessary). $x_s := B(0)$ and $x_e := B(\tau)$ are the starting point and terminal point of the excursion. One possible definition of μ_D is the following: Suppose first that D is the unit disc. For any $s > 0$ define the measure P^s on Brownian paths (modulo continuous increasing time change) started uniformly on the circle of radius $\exp(-s)$, and killed when they exit D . Note for any $s_0 > s$, the killed Brownian path defined under the probability measure P^s has a probability s/s_0 to intersect the circle of radius $\exp(-s_0)$. Then, define

$$\mu_D := \lim_{s \searrow 0} (2\pi/s) P^s.$$

One can then easily check that for any Möbius transformation ϕ from D onto D , $\phi(\mu_D) = \mu_D$. This makes it possible to extend the definition of μ_D to any simply-connected domain D , by conformal invariance. These Brownian excursions also have a “restriction” property [29], as the Brownian paths only feel the boundary of D when they hit it (and get killed).

Suppose for a moment that $\mathcal{R} = \mathcal{R}(L) \subset \mathbf{C}$ is the rectangle with corners given by (3.1), and that \mathcal{B} is the trace of the Brownian excursion $(B(t), t \leq \tau)$ in \mathcal{R} . Define the event

$$E_1 = \{B(0) \in [A_1, A_4] \text{ and } B(\tau) \in [A_2, A_3]\},$$

i.e., B crosses the rectangle from the left to the right. (Although $\mu_{\mathcal{R}}$ is an infinite measure, $\mu_{\mathcal{R}}(E_1)$ is finite.) When E_1 holds, let \mathcal{R}_B^+ be the component of $\mathcal{R} \setminus \mathcal{B}$ above \mathcal{B} , and let \mathcal{R}_B^- be the component of $\mathcal{R} \setminus \mathcal{B}$ below \mathcal{B} . Let \mathcal{L}_B^- (resp. \mathcal{L}_B^+) denote the π -extremal distance between $[A_1, x_s]$ and $[A_2, x_e]$ in \mathcal{R}_B^- (resp. $[x_s, A_4]$ and $[x_e, A_3]$) in \mathcal{R}_B^+ .

Then, for any $\alpha \geq 0$ and $\alpha' \geq 0$, the exponent $\tilde{\xi}(\alpha, 1, \alpha') = \tilde{\xi}(1, \tilde{\xi}(\alpha, \alpha'))$ is characterized by

$$\mathbf{E}_{\mu_{\mathcal{R}}} [1_{E_1} \exp(-\alpha \mathcal{L}_B^+ - \alpha' \mathcal{L}_B^-)] = \exp(-\tilde{\xi}(\alpha', 1, \alpha)L + o(L)), \tag{4.2}$$

when $L \rightarrow \infty$, where $\mathbf{E}_{\mu_{\mathcal{R}}}$ denotes expectation (that is, integration) with respect to the measure $\mu_{\mathcal{R}}$. Similarly,

$$\mathbf{E}_{\mu_{\mathcal{R}}} [1_{E_1} \exp(-\alpha \mathcal{L}_B^+)] = \exp(-\tilde{\xi}(1, \alpha)L + o(L)), \quad L \rightarrow \infty. \tag{4.3}$$

See [29]. It will also be important later that $\tilde{\xi}$ is continuous in its arguments, and that $\lambda \mapsto \tilde{\xi}(1, \lambda)$ is strictly monotone.

4.2. Statement and proof

For any $p \geq 0$, we put

$$v_p = \frac{1}{6}p(p+1).$$

Let \mathcal{V} denote the set of numbers $\{v_p : p \in \mathbf{N}\}$. Note that the smallest values in \mathcal{V} are $0, \frac{1}{3}, 1, 2, \frac{10}{3}, 5, 7$.

We are now ready to prove the following result:

THEOREM 4.1. *For any $k \geq 2$, $\alpha_1, \dots, \alpha_{k-1}$ in \mathcal{V} , and for all $\alpha_k \in \mathbf{R}_+$,*

$$\tilde{\xi}(\alpha_1, \dots, \alpha_k) = \frac{1}{24}((\sqrt{24\alpha_1+1} + \dots + \sqrt{24\alpha_k+1} - (k-1))^2 - 1). \tag{4.4}$$

It is immediate to verify that this theorem implies Theorem 1.1.

Remark. In [26], Theorem 4.1 is extended to all non-negative reals $\alpha_1, \dots, \alpha_k$.

Theorem 4.1 is a consequence of the cascade relations and the following lemma, which is the special case of the theorem with $k=2$, $\alpha_1 = \frac{1}{3}$:

LEMMA 4.2. *For any $\lambda > 0$,*

$$\tilde{\xi}\left(\frac{1}{3}, \lambda\right) = u(\lambda),$$

where $u(\lambda)$ is given by (3.2).

Proof of Theorem 4.1 (assuming Lemma 4.2). Define $U(\lambda) = \sqrt{24\lambda+1} - 1$, for all $\lambda \geq 0$. Lemma 4.2 implies immediately that for all $\lambda \geq 0$,

$$U\left(\tilde{\xi}\left(\frac{1}{3}, \lambda\right)\right) = U(\lambda) + 2 = U(\lambda) + U\left(\frac{1}{3}\right)$$

and (for all integer p) $v_{p+1} = \tilde{\xi}\left(\frac{1}{3}, v_p\right)$. The cascade relations then imply that for all integers p_1, \dots, p_{k-1} ,

$$\tilde{\xi}(v_{p_1}, \dots, v_{p_{k-1}}, \lambda) = U^{-1}(2(p_1 + \dots + p_{k-1}) + U(\lambda)).$$

This is (4.4). □

Proof of Lemma 4.2. For convenience, we again work in a rectangle rather than in the upper half-plane. Let $\mathcal{R} = \mathcal{R}(L)$, and let \mathcal{S} denote the closure of the hull of SLE_6 from A_4 to $[A_1, A_2] \cup [A_2, A_3]$ in \mathcal{R} , as in Theorem 3.1. Let \mathcal{B} denote the trace of a Brownian

excursion in \mathcal{R} ; we will call its starting point x_s and its terminal point x_e . Consider the events

$$\begin{aligned} E_1 &= \{x_s \in [A_1, A_4] \text{ and } x_e \in [A_2, A_3]\}, \\ E_2 &= \{\mathcal{S} \cap [A_1, A_2] = \emptyset\}, \\ E_3 &= E_1 \cap E_2 \cap \{\mathcal{S} \cap \mathcal{B} = \emptyset\}. \end{aligned}$$

When E_2 holds, let \mathcal{L}_S denote the π -extremal distance between the vertical edges of \mathcal{R} in $\mathcal{R} \setminus \mathcal{S}$ (that is, in the quadrilateral “below” \mathcal{S}). Otherwise, let $\mathcal{L}_S = \infty$.

When E_1 holds, let \mathcal{R}_B^+ be the component of $\mathcal{R} \setminus \mathcal{B}$ above \mathcal{B} , and let \mathcal{R}_B^- be the component of $\mathcal{R} \setminus \mathcal{B}$ below \mathcal{B} . Let \mathcal{L}_B^- (resp. \mathcal{L}_B^+) denote the π -extremal distance between the vertical edges of \mathcal{R} in \mathcal{R}_B^- (resp. in \mathcal{R}_B^+), as before. When E_3 holds, let \mathcal{L}_{SB} denote the π -extremal distance between the vertical edges of \mathcal{R} in $\mathcal{R}_B^+ \setminus \mathcal{S}$ (that is, in the quadrilateral “below \mathcal{S} and above \mathcal{B} ”).

Let $\lambda > 0$. We are interested in the asymptotic behavior of

$$f(L) = \mathbf{E}[1_{E_3} \exp(-\lambda \mathcal{L}_{SB})]$$

when $L \rightarrow \infty$. By first taking expectations with respect to B (with the measure $\mu_{\mathcal{R}}$), and using the restriction property (Corollary 2.4) for the domains \mathcal{R} and \mathcal{R}_B^+ , it follows that as $L \rightarrow \infty$,

$$\begin{aligned} f(L) &= \mathbf{E}_B[\mathbf{E}_S[\exp(-\lambda \mathcal{L}_{SB})]] \\ &= \mathbf{E}_B[\exp(-u(\lambda) \mathcal{L}_B^+ + O(1))] \quad (\text{by Theorem 3.1 and restriction to } \mathcal{R}_B^+) \\ &= \exp(-\tilde{\xi}(1, u(\lambda))L + o(L)) \quad (\text{by (4.3)}). \end{aligned}$$

On the other hand, we may first take expectation with respect to \mathcal{S} . Given \mathcal{S} , the law of \mathcal{L}_{SB} is the same as that of \mathcal{L}_B^- , by complete conformal invariance of the excursion measure (which is the analogue of the restriction property to the excursion measure, see [28]). Hence, as $L \rightarrow \infty$,

$$\begin{aligned} f(L) &= \mathbf{E}_S[\mathbf{E}_B[1_{E_3} \exp(-\lambda \mathcal{L}_{SB})]] \\ &= \mathbf{E}_S[\mathbf{E}_B[1_{E_3} \exp(-\lambda \mathcal{L}_B^-)]] \\ &= \mathbf{E}_B[\mathbf{E}_S[1_{E_3} \exp(-\lambda \mathcal{L}_B^-)]] \\ &= \mathbf{E}_B[\mathbf{P}_S[E_3 | \mathcal{L}_B^+] \exp(-\lambda \mathcal{L}_B^-)] \\ &= \mathbf{E}_B[\exp(-\frac{1}{3} \mathcal{L}_B^+ + O(1)) \exp(-\lambda \mathcal{L}_B^-)] \quad (\text{by (3.4)}) \\ &= \exp(-\tilde{\xi}(\frac{1}{3}, 1, \lambda)L + o(L)) \quad (\text{by (4.2)}) \\ &= \exp(-\tilde{\xi}(1, \tilde{\xi}(\frac{1}{3}, \lambda))L + o(L)), \end{aligned}$$

by the cascade relations (4.1). Comparing with (4.5) gives $\tilde{\xi}(1, \tilde{\xi}(\frac{1}{3}, \lambda)) = \tilde{\xi}(1, u(\lambda))$. Finally,

$$\tilde{\xi}(\frac{1}{3}, \lambda) = u(\lambda)$$

follows, since $\lambda' \mapsto \tilde{\xi}(1, \lambda')$ is strictly increasing. □

5. Crossing exponents for critical percolation

It has been conjectured [42] that SLE_6 corresponds to the scaling limit of critical percolation clusters. As additional support for this conjecture, we now show that it implies the conjectured formula for the exponents corresponding to the probability that a long rectangle is crossed by p disjoint paths or clusters of critical percolation ([13], [8], [3]).

Let us first explain the conjectured relation between SLE_6 and critical percolation. Let $D \subset \mathbf{C}$ be a domain whose boundary $\partial D \subset \mathbf{C}$ is a simple closed curve. Let $a, b \in \partial D$ be distinct points. Let γ_1 be the counterclockwise arc on ∂D from a to b , and let γ_2 be the clockwise arc on ∂D from a to b . Let $\delta > 0$, and consider a fine hexagonal grid H in the plane with mesh δ ; that is, each face of the grid is a regular hexagon with edges of length δ , and each vertex has degree 3. For simplicity, assume that ∂D does not pass through a vertex of H , and that a and b do not lie on edges of H . Color each hexagon of H independently, black or white, with probability $\frac{1}{2}$. Then the union of the black hexagons forms one of the standard models for critical percolation (see Grimmett [14] for percolation background and references).

Given the random coloring, there is a unique path $\beta \subset \bar{D}$ that starts at a and ends at b , such that whenever β is not on γ_1 it has a black hexagon on its “right”, and whenever β is not on γ_2 it has a white hexagon on its “left”. This path is the boundary between the union of the white clusters in D touching γ_2 and the black clusters in D touching γ_1 . Let $f: D \rightarrow \mathbf{H}$ be a conformal homeomorphism such that $f(a) = 0$ and $f(b) = \infty$, and parameterize β in such a way that $A(f(\beta[0, t])) = t$. Let D_t be the component of $D \setminus \beta[0, t]$ that has b on its boundary, and let $K_t = D \setminus D_t$. The conjecture from [42] (stated a bit differently) is that as $\delta \rightarrow 0$ the process $(K_t, t \geq 0)$ converges to SLE_6 from a to b in D . In light of this conjecture, the Locality Theorem 2.2 and its corollaries are very natural.

Now consider an arc $I \subset \partial D$, which contains b but not a . Let b_1 and b_2 be the endpoints of I , labeled in such a way that the triplet a, b_1, b_2 is in counterclockwise order around D . Let $\gamma'_1 \subset \gamma_1$ be the counterclockwise arc from a to b_1 , and let $\gamma'_2 \subset \gamma_2$ be the clockwise arc from a to b_2 . Let T be the first time such that $\beta(t) \in I$, and set $S := \bigcup_{t < T} K_t$. Then the component α_1 of $\partial S \cap D$ joining γ'_1 to I is a crossing in \bar{B} from γ'_1 to I which is “maximal”, in the sense that any other crossing $\alpha \subset \bar{B}$ from γ'_1 to I is separated by α_1 from b_2 in D .

Let L be large, and recall the definition of the rectangle $\mathcal{R}=\mathcal{R}(L)$ with corners given by (3.1). Let $p\in\mathbf{N}_+$ and $\sigma=(\sigma_1, \dots, \sigma_p)\in\{\text{black, white}\}^p$. Consider the event $C_\sigma(\mathcal{R})$ that there are paths $\alpha_1, \dots, \alpha_p$ from $[A_4, A_1]$ to $[A_2, A_3]$ in \mathcal{R} such that each α_j is contained in the union of the hexagons of color σ_j , there is no hexagon which intersects more than one of these paths, and α_{j+1} separates α_j from $[A_1, A_2]$ in \mathcal{R} when $j=1, 2, \dots, p-1$.

Take α_1 to be the topmost crossing with color σ_1 , if such exists, let α_2 be the topmost crossing with color σ_2 which is below all the hexagons meeting α_1 , etc. Then $C_\sigma(\mathcal{R})$ holds if and only if these specific $\alpha_1, \dots, \alpha_p$ exist. Note that after we condition on α_1 , the hexagons “below” it are still independent and are black or white with probability $\frac{1}{2}$. Hence the following formula holds:

$$\mathbf{P}[C_\sigma(\mathcal{R})] = \mathbf{P}[\alpha_1 \text{ exists}] \mathbf{E}[\mathbf{P}[C_{\sigma_{+1}}(\mathcal{R}_{\alpha_1}) | \alpha_1] | \alpha_1 \text{ exists}],$$

where $\sigma_{+1}=(\sigma_2, \sigma_3, \dots, \sigma_p)$, \mathcal{R}_{α_1} is the union of the hexagons below α_1 , and $C_{\sigma_{+1}}(\mathcal{R}_{\alpha_1})$ is the event that there are multiple crossings with colors specified by σ_{+1} from $[A_4, A_1]$ to $[A_2, A_3]$ in \mathcal{R}_{α_1} .

It is clear that $\mathbf{P}[C_\sigma(\mathcal{R})]$ does not depend on the choice of the sequence σ , but only its length. Moreover, the conjectured conformal invariance (or the conjecture that SLE_6 is the scaling limit) implies that $\lim_{\delta\rightarrow 0} \mathbf{P}[C_\sigma(\mathcal{D})]$ depends on the quadrilateral \mathcal{D} only through its conformal modulus. Hence define

$$f_p(L) := \lim_{\delta\rightarrow 0} \mathbf{P}[C_\sigma(\mathcal{R}(L))], \quad \sigma \in \{\text{black, white}\}^p.$$

We also set $f_p(\infty):=0$ and $f_0(L):=1_{\{L<\infty\}}$.

Let S be the SLE_6 -hull from A_4 to $I:=[A_1, A_2]\cup[A_2, A_3]$ in $\mathcal{R}=\mathcal{R}(L)$, as defined in §3.1. Let \mathcal{R}_- be the component of $\mathcal{R}\setminus S$ which has A_1 on its boundary, and let \mathcal{L} denote the π -extremal length from $[A_4, A_1]$ to $[A_2, A_3]$ in \mathcal{R}_- . Note that $\mathcal{L}=\infty$ if $\bar{S}\cap[A_1, A_2]\neq\emptyset$. Then we have

$$f_p(L) = \mathbf{E}[f_{p-1}(\mathcal{L})], \quad p = 1, 2, \dots$$

To completely justify this step requires more work, which we omit, since this whole discussion depends on a conjecture anyway. The slight difficulty has to do with the fact that having a crossing of a closed rectangle is a closed condition, and the probability of a closed event can go up when taking a weak limit of measures. One simple way to deal with this is to note that when the continuous process has a crossing in the rectangle $\mathcal{R}(L+\varepsilon)$, every sufficiently close discrete approximation of it has a crossing of the rectangle $\mathcal{R}(L)$.

Consequently, induction and Theorem 3.1 give

$$f_p(L) = \exp(-(L+O(1))v_p), \quad L \rightarrow \infty, \tag{5.1}$$

where

$$v_p = u^{\circ p}(0) = \frac{1}{6}p(p+1), \quad (5.2)$$

as before. Here, the constant implicit in the $O(1)$ -notation may depend on p .

Note also that if

$$\sigma = (\text{white, black, white, black, } \dots, \text{white}) \in \{\text{black, white}\}^{2k-1},$$

then (in the discrete setting) the event $C_\sigma(\mathcal{R})$ is identical to the event that the rectangle \mathcal{R} is crossed from left to right by k disjoint white clusters.

The exponents (5.2) are those predicted in [8], [3].

References

- [1] AHLFORS, L. V., *Conformal Invariants: Topics in Geometric Function Theory*. McGraw-Hill, New York, 1973.
- [2] AIZENMAN, M., The geometry of critical percolation and conformal invariance, in *STAT-PHYS 19* (Xiamen, 1995), pp. 104–120. World Sci. Publishing, River Edge, NJ, 1996.
- [3] AIZENMAN, M., DUPLANTIER, B. & AHARONY, A., Path crossing exponents and the external perimeter in 2D percolation. *Phys. Rev. Lett.*, 83 (1999), 1359–1362.
- [4] BELAVIN, A. A., POLYAKOV, A. M. & ZAMOLODCHIKOV, A. B., Infinite conformal symmetry in two-dimensional quantum field theory. *Nuclear Phys. B*, 241 (1984), 333–380.
- [5] BURDZY, K. & LAWLER, G. F., Non-intersection exponents for Brownian paths. Part I: Existence and an invariance principle. *Probab. Theory Related Fields*, 84 (1990), 393–410.
- [6] CARDY, J. L., Conformal invariance and surface critical behavior. *Nuclear Phys. B*, 240 (1984), 514–532.
- [7] — Critical percolation in finite geometries. *J. Phys. A*, 25 (1992), L201–L206.
- [8] — The number of incipient spanning clusters in two-dimensional percolation. *J. Phys. A*, 31 (1998), L105.
- [9] CRANSTON, M. & MOUNTFORD, T., An extension of a result of Burdzy and Lawler. *Probab. Theory Related Fields*, 89 (1991), 487–502.
- [10] DUPLANTIER, B., Loop-erased self-avoiding walks in two dimensions: Exact critical exponents and winding numbers. *Phys. A*, 191 (1992), 516–522.
- [11] — Random walks and quantum gravity in two dimensions. *Phys. Rev. Lett.*, 81 (1998), 5489–5492.
- [12] DUPLANTIER, B. & KWON, K.-H., Conformal invariance and intersection of random walks. *Phys. Rev. Lett.*, 61 (1988), 2514–2517.
- [13] DUPLANTIER, B. & SALEUR, H., Exact determination of the percolation hull exponent in two dimensions. *Phys. Rev. Lett.*, 58 (1987), 2325–2328.
- [14] GRIMMETT, G., *Percolation*. Springer-Verlag, New York, 1989.
- [15] IKEDA, N. & WATANABE, S., *Stochastic Differential Equations and Diffusion Processes*, 2nd edition. North-Holland Math. Library, 24. North-Holland, Amsterdam, 1989.
- [16] KENYON, R., Conformal invariance of domino tiling. *Ann. Probab.*, 28 (2000), 759–795.
- [17] — The asymptotic determinant of the discrete Laplacian. *Acta Math.*, 185 (2000), 239–286.
- [18] — Long-range properties of spanning trees. Probabilistic techniques in equilibrium and nonequilibrium statistical physics. *J. Math. Phys.*, 41 (2000), 1338–1363.

- [19] LANGLANDS, R., POULIOT, P. & SAINT-AUBIN, Y., Conformal invariance in two-dimensional percolation. *Bull. Amer. Math. Soc. (N.S.)*, 30 (1994), 1–61.
- [20] LAWLER, G. F., *Intersections of Random Walks*. Birkhäuser Boston, Boston, MA, 1991.
- [21] — Hausdorff dimension of cut points for Brownian motion. *Electron. J. Probab.*, 1:2 (1996), 1–20 (electronic).
- [22] — The dimension of the frontier of planar Brownian motion. *Electron. Comm. Probab.*, 1:5 (1996), 29–47 (electronic).
- [23] — The frontier of a Brownian path is multifractal. Preprint, 1997.
- [24] LAWLER, G. F. & PUCKETTE, E. E., The intersection exponent for simple random walk. *Combin. Probab. Comput.*, 9 (2000), 441–464.
- [25] LAWLER, G. F., SCHRAMM, O. & WERNER, W., Values of Brownian intersection exponents, II: Plane exponents. *Acta Math.*, 187 (2001), 275–308.
<http://arxiv.org/abs/math.PR/0003156>.
- [26] — Values of Brownian intersection exponents, III: Two-sided exponents. To appear in *Ann. Inst. H. Poincaré Probab. Statist.* <http://arxiv.org/abs/math.PR/0005294>.
- [27] — Analyticity of intersection exponents for planar Brownian motion. To appear in *Acta Math.*, 188 (2002). <http://arxiv.org/abs/math.PR/0005295>.
- [28] LAWLER, G. F. & WERNER, W., Intersection exponents for planar Brownian motion. *Ann. Probab.*, 27 (1999), 1601–1642.
- [29] — Universality for conformally invariant intersection exponents. *J. Eur. Math. Soc. (JEMS)*, 2 (2000), 291–328.
- [30] LEBEDEV, N. N., *Special Functions and Their Applications*. Dover, New York, 1972.
- [31] LEHTO, O. & VIRTANEN, K. I., *Quasiconformal Mappings in the Plane*, 2nd edition. Grundlehren Math. Wiss., 126. Springer-Verlag, New York–Heidelberg, 1973.
- [32] LÖWNER, K., Untersuchungen über schlichte konforme Abbildungen des Einheitskreises, I. *Math. Ann.*, 89 (1923), 103–121.
- [33] MADRAS, N. & SLADE, G., *The Self-Avoiding Walk*. Birkhäuser Boston, Boston, MA, 1993.
- [34] MAJUMDAR, S. N., Exact fractal dimension of the loop-erased random walk in two dimensions. *Phys. Rev. Lett.*, 68 (1992), 2329–2331.
- [35] MANDELBROT, B. B., *The Fractal Geometry of Nature*. Freeman, San Francisco, CA, 1982.
- [36] MARSHALL, D. E. & ROHDE, S., The Loewner differential equation and slit mappings. Preprint.
- [37] POMMERENKE, CH., On the Loewner differential equation. *Michigan Math. J.*, 13 (1966), 435–443.
- [38] — *Boundary Behaviour of Conformal Maps*. Grundlehren Math. Wiss., 299. Springer-Verlag, Berlin, 1992.
- [39] REVUZ, D. & YOR, M., *Continuous Martingales and Brownian Motion*. Grundlehren Math. Wiss., 293. Springer-Verlag, Berlin, 1991.
- [40] ROHDE, S. & SCHRAMM, O., Basic properties of SLE. Preprint.
- [41] RUDIN, W., *Real and Complex Analysis*, 3rd edition. McGraw-Hill, New York, 1987.
- [42] SCHRAMM, O., Scaling limits of loop-erased random walks and uniform spanning trees. *Israel J. Math.*, 118 (2000), 221–288.

GREGORY F. LAWLER
Department of Mathematics
Duke University
Box 90320
Durham, NC 27708-0320
U.S.A.
jose@math.duke.edu

ODED SCHRAMM
Microsoft Research
1, Microsoft Way
Redmond, WA 98052
U.S.A.
schramm@microsoft.com

WENDELIN WERNER
Département de Mathématiques
Université Paris-Sud
Bât. 425
FR-91405 Orsay Cedex
France
wendelin.werner@math.u-psud.fr

Received January 3, 2000