CORRECTION NOTES

CORRECTION TO "A REMARK ON SEQUENTIAL DISCRIMINATION"

By David A. Freedman

University of California, Berkeley

I stated incomplete conditions for Theorem 2 in [1]. Dr. B. Weiss pointed this out to me. To correct the error, replace the first two sentences in Section 4 (Ann. Math. Statist. 38 1666-1670) with the following.

"For Theorem 2, suppose that each α_n is finitely generated. Suppose further that $A_1 \supset A_2 \supset \cdots$ and A_n an atom of α_n for all n imply $\bigcap_n A_n$ is non empty. Then any pairwise disjoint sequence of sets in α whose union is α is empty from some time on, so there is no distinction between finite and countable additivity."

The first full paragraph on page 1668 should be corrected in a similar way, so the first sentence reads:

"If each α_n is the σ -field generated by a countable partition; if $A_1 \supset A_2 \supset \cdots$ and A_n an atom of α_n for all n imply $\bigcap_n A_n$ is non empty; and each $P \cdot \Pi$ is countably additive: the situation is similar."

CORRECTION TO "PROBABILITY DENSITIES WITH GIVEN MARGINALS"

By S. Kullback

The George Washington University

On page 1243 of the *Ann. Math. Statist.* **39,** reference [3], change 315–324 to 179–188.

CORRECTION TO "A NOTE ON CONSERVATIVE CONFIDENCE INTERVALS FOR THE MEAN OF A MULTIVARIATE NORMAL"

By Alastair Scott

The London School of Economics

The statement after Lemma 2 on page 279 (Ann. Math. Statist. 38) that the proof of Lemma 2 can be generalized to m > 2 is not true, so that Lemma 2 and Theorem 2 are only proved for n = 2. A proof of Theorem 2 in a special case is contained in a forthcoming paper by Z. Sidak.