

From random partitions to fractional Brownian sheets

OLIVIER DURIEU¹ and YIZAO WANG²

¹*Laboratoire de Mathématiques et Physique Théorique, UMR-CNRS 7350, Fédération Denis Poisson, FR-CNRS 2964, Université de Tours, Parc de Grandmont, 37200 Tours, France.*

E-mail: olivier.durieu@lmpt.univ-tours.fr

²*Department of Mathematical Sciences, University of Cincinnati, 2815 Commons Way, Cincinnati, OH 45221-0025, USA. E-mail: yizao.wang@uc.edu*

We propose discrete random-field models that are based on random partitions of \mathbb{N}^2 . The covariance structure of each random field is determined by the underlying random partition. Functional central limit theorems are established for the proposed models, and fractional Brownian sheets, with full range of Hurst indices, arise in the limit. Our models could be viewed as discrete analogues of fractional Brownian sheets, in the same spirit that the simple random walk is the discrete analogue of the Brownian motion.

Keywords: fractional Brownian motion; fractional Brownian sheet; invariance principle; long-range dependence; random field; random partition; regular variation

1. Introduction

In this paper, we propose random-field models that are based on random partitions, and show that their partial-sum random fields scale to fractional Brownian sheets. Our motivation came from three recent papers, one by Hammond and Sheffield [12] and two by the authors and collaborator [3,9], where it was shown that fractional Brownian motions and some operator-scaling Gaussian random fields (that can be viewed as random-field generalizations of fractional Brownian motions, see [4]) may arise as the scaling limits of certain stochastic models, the dependence structure of which is essentially determined by certain random partitions of $\mathbb{N} = \{1, 2, \dots\}$. We start by briefly recalling the results in one dimension.

The two papers [9,12] established functional central limit theorems for fractional Brownian motions based on two different random partitions. In each model, there is an underlying random partition of the integers $\{1, \dots, n\}$, and conditioning on the random partition, ± 1 -valued random spins X_1, \dots, X_n are assigned, in certain ways to be specified later. The advantage of taking random spins is that in this way, the covariances of the partial sums are determined by the underlying random partitions. By appropriately choosing the random partition and the assignment rule of random spins, the partial sum $S_n = X_1 + \dots + X_n$ scales to a fractional Brownian motion as $n \rightarrow \infty$ in the form of

$$\left\{ \frac{S_{[nt]}}{n^H L(n)} \right\}_{t \in [0,1]} \Rightarrow \{B_t^H\}_{t \in [0,1]} \quad (1.1)$$

in $D([0, 1])$ as $n \rightarrow \infty$, where L is a slowly varying function at infinity and \mathbb{B}^H on the right-hand side above denotes the fractional Brownian motion with Hurst index $H \in (0, 1)$, a centered Gaussian process with covariance function

$$\text{Cov}(\mathbb{B}_s^H, \mathbb{B}_t^H) = \frac{1}{2}(t^{2H} + s^{2H} - |t - s|^{2H}), \quad s, t \geq 0.$$

Throughout, \Rightarrow stands for convergence in distribution and $D([0, 1])$ for the space of càdlàg functions equipped with the Skorohod topology [5].

The models in [9,12] are different in both the underlying random partitions and the ways of assigning ± 1 spins, and they lead to different ranges of Hurst index: $H \in (0, 1/2)$ in [9] and $H \in (1/2, 1)$ in [12]. The partial sum S_n can be interpreted as a correlated random walk and provides a simple discrete counterpart to the fractional Brownian motion, in the same spirit that the simple random walk can be viewed as the discrete counterpart of the Brownian motion.

In view of the non-standard normalization $n^H L(n)$ in (1.1) instead of $n^{1/2}$ for S_n of the i.i.d. random variables, such models are sometimes referred to as having *long-range dependence* [26,30]. Moreover, the fractional Brownian motion in the limit characterizes the non-negligible dependence at macroscopic scale of the discrete model when $H \neq 1/2$ (recall that $\mathbb{B}^{1/2}$ is a standard Brownian motion). Such limit theorems are of special interest for the study of long-range dependence, as they often reveal different types of dynamics underlying certain common long-range dependence phenomena. Namely, drastically different models may lead to the same stochastic process with long-range dependence, and fractional Brownian motions often show up in such limit theorems. Fractional Brownian motions, first considered by Kolmogorov [18] and studied rigorously by Mandelbrot and Van Ness [21], are arguably the most important stochastic processes in the investigation of long-range dependence: it is well known now that fractional Brownian motions arise in limit theorems on models from various areas, including finance [17], telecommunications [23], interacting particle systems [25], aggregation of correlated random walks [10], just to mention a few.

Results in [9,12] provide a new class of examples for long-range dependence: they may arise in the presence of certain random partitions. Such a point of view, to the best of our knowledge, has been rarely explored before. Our motivation is to demonstrate that the random-partition mechanism behind the long-range dependence phenomena in the aforementioned papers remains at work in a natural random-field setup. The generalization of aforementioned one-dimensional random partitions to high dimensions, however, is far from being unique. A first attempt has been successfully worked out in [3], where certain operator-scaling Gaussian random fields appear in the limit (see Remark 3.4).

Here we continue to explore other possibilities of random-field extensions. In particular, we shall propose three random-field extensions of the one-dimensional models, and show that the partial sums of proposed models scale to fractional Brownian sheets. Our limit theorems shall cover the full range of Hurst indices for the fractional Brownian sheets. This is in sharp contrast to the previous random-field model investigated earlier in [3], where the limit random fields are most of the time not fractional Brownian sheets. This reflects the fact that the random partitions considered here are essentially different from the ones considered in [3], and hence our models and limit theorems here complement the ones therein.

Recall that a fractional Brownian sheet with Hurst index $\mathbf{H} = (H_1, H_2) \in (0, 1)^2$ is a multi-parameter zero-mean Gaussian process $\{\mathbb{B}_t^{\mathbf{H}}\}_{t \in \mathbb{R}_+^2}$ with covariance

$$\text{Cov}(\mathbb{B}_s^{\mathbf{H}}, \mathbb{B}_t^{\mathbf{H}}) = \prod_{q=1}^2 \frac{1}{2} (t_q^{2H_q} + s_q^{2H_q} - |t_q - s_q|^{2H_q}), \quad \mathbf{t}, \mathbf{s} \in \mathbb{R}_+^2.$$

Fractional Brownian sheets are random-field generalizations of fractional Brownian motions proposed by Kamont [15]. These are centered Gaussian processes that are operator-scaling (generalization of self-similarity to random fields, see e.g. [4]) and with stationary rectangular increments. In the special case $H_1 = H_2 = 1/2$, the fractional Brownian sheet becomes the standard Brownian sheet, the random-field generalization of Brownian motion. For other Hurst indices, fractional Brownian sheets exhibit anisotropic long-range dependence. Representation and path properties of these random fields have been extensively investigated. See for example the recent survey by Xiao [33]. Stochastic partial differential equations driven by fractional Brownian sheets have also been studied (see, e.g., [13,24]). At the same time, fractional Brownian sheets are not the only operator-scaling random fields with stationary rectangular increments. There are other random fields with long-range dependence which could also be viewed as generalization of fractional Brownian motions. Limit theorems for fractional Brownian sheets and other Gaussian random fields with long-range dependence, however, have not been as much developed as for fractional Brownian motions. Recent developments in this direction include for examples limit theorems for linear random fields [19,32], for set-indexed fields [2], and for aggregated models [29,31].

Now we describe our models, which are extensions of the one-dimensional models in [9,12] to two dimensions, in more details. For these one-dimensional models, we first introduce a random partition of \mathbb{N} into different components, where each component may have possibly an infinite number of elements. Next, given a random decomposition $\{\mathcal{C}_k\}_{k \in \mathbb{N}}$ of \mathbb{N} , for each component \mathcal{C} we sample $X_{\mathcal{C}} = \{X_i\}_{i \in \mathcal{C}}$ according to a specific *assignment rule*, applied in an independent manner to all components $\{X_{\mathcal{C}_k}\}_{k \in \mathbb{N}}$. For these models, each X_i takes the values ± 1 only. We consider two possible assignment rules:

Identical assignment rule. Assign the same values for all X_i in the same component. The identical value is either 1 or -1 , with equal probabilities.

Alternating assignment rule. Assign ± 1 values in an alternating manner with respect to the natural order on \mathbb{N} , for X_i in the same component. Given a component, there are two such ways of assigning ± 1 values, and one of them is chosen with probability $1/2$. For example, given a component $\mathcal{C} = \{1, 2, 5\}$, the alternating assignment rule assigns $(1, -1, 1)$ or $(-1, 1, -1)$ with equal probabilities to (X_1, X_2, X_5) .

In particular, the Hammond–Sheffield model in [12] is based on a random partition of \mathbb{N} induced by a random forest with infinite components, each being an infinite tree, and the identical assignment rule (the random forest induces actually a random partition of \mathbb{Z}). The model in [9] is based on an exchangeable random partition on \mathbb{N} [27] induced by a certain infinite urn scheme and the alternating assignment rule. It is a modification of a model originally investigated in Karlin [16], and hence we refer to the model as the *randomized Karlin model*. The two models will be recalled in full detail in later sections. Note that this framework of building stationary




Figure 1. A component from a product of partitions. Left: component $\{1, 2, 4, 5, 6\} \times \{1, 2, 4, 6, 7\}$. Middle: alternating assignment rule of 1 (black) and -1 (white) values. Right: mixture of identical assignment rule in horizontal direction and alternating assignment rule in vertical direction.

sequences based on random partitions and assignment rules also includes the example of independent ± 1 spins, of which the partial-sum process is well known to scale to a Brownian motion. To achieve this, it suffices to take the finest partition of \mathbb{N} , that is, each component corresponds exactly to one element from \mathbb{N} , and then apply either assignment rule (the two are the same in this case).

Our random-field models are based on random partitions of \mathbb{N}^2 obtained as the product of two independent random partitions of \mathbb{N} . Namely, let $\mathcal{C}^{(q)} = \{\mathcal{C}_i^{(q)}\}_{i \in \mathbb{N}}$, $q = 1, 2$, be two partitions of \mathbb{N} . Let $\mathcal{C} = \mathcal{C}^{(1)} \times \mathcal{C}^{(2)}$ denote the partition of \mathbb{N}^2 whose components are the Cartesian products $\mathcal{C}_i^{(1)} \times \mathcal{C}_j^{(2)}$ for all $i, j \in \mathbb{N}$ (e.g., $\{1, 2\} \times \{1, 3\} = \{(1, 1), (1, 3), (2, 1), (2, 3)\}$ is a subset of \mathbb{N}^2). Once the random partition is given, one of the two assignment rules is applied in each direction. Figure 1 illustrates the product of partitions (left), the alternating assignment rule (middle), and an assignment rule of mixed type (right).

We shall investigate the partial-sum random fields of three ± 1 -valued models, each converging to fractional Brownian sheets in a different regime in terms of the Hurst indices. The contributions of the paper are summarized here.

- (i) In Section 2, we propose a generalization of the randomized Karlin model and show that the partial-sum random field scales to a fractional Brownian sheet with $\mathbf{H} \in (0, 1/2)^2$.
- (ii) In Section 3, we propose a generalization of the Hammond–Sheffield model and show that the partial-sum random field scales to a fractional Brownian sheet with $\mathbf{H} \in (1/2, 1)^2$.
- (iii) In Section 4, we propose a model that can be viewed as a combination of the Hammond–Sheffield model and the randomized Karlin model, and show that the partial-sum random field scales to a fractional Brownian sheet with $\mathbf{H} \in (1/2, 1) \times (0, 1/2)$.

More specifically, our main results Theorems 2.2, 3.3, and 4.1 are limit theorems in the form of

$$\frac{1}{Z_{\mathbf{H}}(\mathbf{n})} \left\{ \sum_{i \in [1, \lfloor \mathbf{n} \cdot t \rfloor]} X_i \right\}_{t \in [0, 1]^2} \Rightarrow \{\mathbb{B}_t^{\mathbf{H}}\}_{t \in [0, 1]^2}$$

in $D([0, 1]^2)$ as $\min(n_1, n_2) \rightarrow \infty$, where $Z_{\mathbf{H}}(\mathbf{n})$ is an appropriate normalization depending on the model (and hence on \mathbf{H}). Our models can thus be viewed as discrete counterparts of fractional Brownian sheets. Throughout, for any element $\mathbf{a} \in \mathbb{R}^2$, we write $\mathbf{a} = (a_1, a_2)$ and for any $\mathbf{a}, \mathbf{b} \in \mathbb{Z}^2$, we write $[\mathbf{a}, \mathbf{b}] = ([a_1, b_1] \times [a_2, b_2]) \cap \mathbb{Z}^2$ the set of points in the rectangle $[a_1, b_1] \times [a_2, b_2]$ with integer coordinates. We also use the notation $[\mathbf{n} \cdot \mathbf{t}] = ([n_1 t_1], [n_2 t_2])$ where $[\cdot]$ stands for the integer part.

The proofs for the two-dimensional randomized Karlin model and the other two models are completely different. For the randomized Karlin model, conditioning on the partition structure, the partial sums become sums of i.i.d. random variables. For the other two models, the proof is based on a martingale central limit theorem due to McLeish [22], already used for the one-dimensional Hammond–Sheffield model. However, the proofs for two-dimensional models are much more demanding than their one-dimensional counterpart, as in general, the martingale central limit theorem does not work as well for random fields as for stationary sequences, as pointed out by Bolthausen [7] already in the 80s. Indeed, for the one-dimensional Hammond–Sheffield model, the normalized partial sum $S_n = X_1 + \dots + X_n$ can be expressed as

$$\frac{S_n}{b_n} = \frac{1}{b_n} \sum_{j \in \mathbb{Z}} b_{n,j} X_j^*$$

for a stationary sequence of martingale differences $\{X_j^*\}_{j \in \mathbb{Z}}$ and some coefficients $b_{n,j}$, with $b_n^2 = \sum_{j \in \mathbb{Z}} b_{n,j}^2$. This is a remarkable representation at the heart of the proof; see [3] (the proof in [12] did not use directly this convenient presentation, but applied nevertheless an argument by martingale approximation). Then, to prove the weak convergence, by McLeish’s central limit theorem, the key step is to show

$$\lim_{n \rightarrow \infty} \frac{1}{b_n^2} \sum_{j \in \mathbb{Z}} b_{n,j}^2 (X_j^*)^2 = \text{Var}(X_0^*) \quad \text{in probability.} \tag{1.2}$$

This requires already an involved argument in one dimension; see [12], Lemma 3.2, and [3], Lemma 7.

In two dimensions, the situation becomes even more complicated as now the partial sum $S_n = \sum_{i \in [1, n]} X_i$ is expressed (Proposition 3.6 below) as

$$\frac{S_n}{b_n} = \frac{1}{b_{n_1}^{(1)}} \sum_{j \in \mathbb{Z}} b_{n_1, j}^{(1)} U_{j, n_2},$$

where $\{U_{j, n_2}\}_{j \in \mathbb{Z}}$ is a stationary martingale-difference sequence with respect to the filtration corresponding to the first direction. The new difficulty of the random-field models comes from the fact that the martingale differences now also depend on n_2 and the dependence structure of the random partition in the second direction. To overcome the new difficulty, at the core of our proofs for the counterpart of (1.2) is a decoupling argument. See Section 3.4.

We conclude the Introduction with a few remarks.

Remark 1.1. If one searches for a similar random field that scales to a fractional Brownian sheet with Hurst index $1/2$ in one direction, one can modify the proposed model by taking instead the finest partition (each integer consists of a component) in that direction. Such a model and its analysis are much easier. The details are omitted.

Remark 1.2. It will become clear that our constructions are not limited to two dimensions only. Our limit theorems could also be extended accordingly to high dimensions, where the limit random fields cover fractional Brownian sheets with all legitimate indices. For high-dimensional models and the corresponding limit theorems, the analysis can be carried out by an induction argument, but will be notationally heavy. Therefore, in this paper we focus on two dimensions, and only discuss the high dimension in Remark 4.4 at the end.

Remark 1.3. Our application of martingale central limit theorem is of a different nature from and much more complicated than the one for the other random-field extension in [3] (see Remark 3.4). There the partial sum can be expressed as a linear random field in the form of

$$\frac{S_n}{b_n} = \frac{1}{b_n} \sum_{j \in \mathbb{Z}^d} b_{n,j} X_j^*, \quad \mathbf{n} \in \mathbb{N}^d, \tag{1.3}$$

with a stationary sequence of martingale differences $\{X_j^*\}_{j \in \mathbb{Z}^d}$ in the lexicographical order for all $d \in \mathbb{N}$, whence the analysis becomes *dimension-free*. To the best of our knowledge, the model in [3] is one of the very rare examples in the literature where a one-dimensional sequence of stationary martingale differences can be elegantly embedded into the presentation of the partial sums of a high-dimensional random field. In general, embedding with respect to the lexicographical order could be formidable [8], and the simple representation (1.3) seems rather a coincidence.

2. Randomized Karlin model

In this section, we introduce the two-dimensional randomized Karlin model, and show that the partial-sum random field scales to a fractional Brownian sheet with Hurst index $H \in (0, 1/2)^2$.

2.1. One-dimensional model

We first recall the one-dimensional randomized Karlin model [9,16]. Let $\{Y_n\}_{n \in \mathbb{N}}$ be i.i.d. random variables with common distribution μ on \mathbb{N} . They induce a partition Π_∞ of \mathbb{N} by setting in the same equivalent class (component), denoted by $i \sim j$, if and only if $Y_i = Y_j$. Intuitively, imagine that we throw balls consecutively and independently into boxes labeled by \mathbb{N} , and set $Y_n = \ell$ if the n th ball falls into the box with label ℓ . This event occurs with probability $p_\ell = \mu(\{\ell\})$, and $i \sim j$ if and only if the balls at round i and j fall in the same box. The partition obtained this way is an infinite exchangeable random partition of \mathbb{N} , sometimes referred to as the partition generated by random samplings, or the paintbox partition [27]. Many estimates of this random partition that we apply here can be found in [11,16].

Throughout, we assume that $\{p_\ell\}_{\ell \in \mathbb{N}}$ is a non-increasing sequence (this can always be assumed because the attached value of each label is irrelevant) and that for some $\alpha \in (0, 1)$,

$$v(x) = \max\{\ell \geq 1 : p_\ell \geq 1/x\} \sim x^\alpha L(x) \quad \text{as } x \rightarrow \infty, \tag{2.1}$$

where L is a slowly varying function at ∞ . Without loss of generality, L is assumed to be bounded away from 0 and ∞ on every compact set in $(0, \infty)$. For example, the condition (2.1) is satisfied (with $L \equiv 1$) when

$$p_k \sim k^{-1/\alpha} \quad \text{as } k \rightarrow \infty.$$

The law of (X_1, \dots, X_n) given the partition Π_n of the set $\{1, \dots, n\}$ induced by Y_1, \dots, Y_n is then determined by the alternating assignment rule. To express the alternating assignment rule explicitly, we introduce

$$Y_{n,\ell} = \sum_{i=1}^n \mathbf{1}_{\{Y_i=\ell\}}, \quad \ell \in \mathbb{N}, \tag{2.2}$$

representing the number of balls in the box ℓ after the first n sampling. Then, the law of (X_1, \dots, X_n) given the partition can be equivalently determined by letting $\{\epsilon_n\}_{n \in \mathbb{N}}$ be i.i.d. random variables, independent of Π_∞ , with common distribution $\mathbb{P}(\epsilon_1 = -1) = \mathbb{P}(\epsilon_1 = 1) = 1/2$ and setting for each $n \in \mathbb{N}$,

$$X_n = \epsilon_\ell (-1)^{Y_{n,\ell}+1} \quad \text{if } Y_n = \ell. \tag{2.3}$$

Originally, Karlin [16] obtained a central limit theorem for the non-randomized model, with $\epsilon_\ell \equiv 1$. The functional central limit theorem of the partial-sum process $S_n = X_1 + \dots + X_n$ was established in [9].

Later in Section 4, we shall need a functional central limit theorem for a slightly more general version of the one-dimensional Karlin model. We say that $\{X_n\}_{n \in \mathbb{N}}$ is a *generalized one-dimensional Karlin model*, if instead of (2.3) we have

$$X_n = Z_\ell \epsilon_\ell (-1)^{Y_{n,\ell}+1} \quad \text{if } Y_n = \ell,$$

for i.i.d. random variables $\{Z_\ell\}_{\ell \in \mathbb{N}}$ with some common distribution ν , independent from Y and ϵ . In this way, given the partition induced by $\{Y_n\}_{n \in \mathbb{N}}$, $X_i = X_j$ if $i \sim j$, and otherwise X_i and X_j are independent and identically distributed as $Z_1 \epsilon_1$.

Proposition 2.1. *For the generalized one-dimensional randomized Karlin model with μ satisfying (2.1) with $\alpha \in (0, 1)$ and a slowly varying function L , for a distribution ν with bounded support, we have*

$$\left\{ \frac{S_{\lfloor nt \rfloor}}{n^{\alpha/2} L(n)^{1/2}} \right\}_{t \in [0,1]} \Rightarrow \sigma_\alpha \{ \mathbb{B}_t^{\alpha/2} \}_{t \in [0,1]}$$

in $D([0, 1])$ as $n \rightarrow \infty$, with

$$\sigma_\alpha^2 = \Gamma(1 - \alpha) 2^{\alpha-1} \text{Var}(X_1).$$

The proof of this result is omitted here, as it can be obtained by following the same strategy as in the proof of Theorem 2.2, the functional central limit theorem for two-dimensional randomized Karlin model to be introduced below. This result could also follow from a multivariate functional central limit theorem for the one-dimensional randomized Karlin model established in [9], Theorem 2.2, Corollary 2.8, where the limit corresponds to a decomposition of fractional Brownian motion into a bi-fractional Brownian motion and another smooth self-similar Gaussian process due to Lei and Nualart [20]. In [9], only the randomized one-dimensional Karlin model was addressed, although the same proof applies to the generalized model with ν having bounded support, too.

2.2. Two-dimensional model and main result

We next describe the two-dimensional randomized Karlin model. For each $q = 1, 2$, consider $Y^{(q)} = \{Y_n^{(q)}\}_{n \in \mathbb{N}}$ as i.i.d. sampling from a certain probability measure μ_q satisfying (2.1) with $\alpha_q \in (0, 1)$ and a slowly varying function L_q . Assume that $Y^{(1)}, Y^{(2)}$ are independent. Then, each $Y^{(q)}$ induces an infinite exchangeable random partition on \mathbb{N} and for each n , let $Y_{n,\ell}^{(q)}$ be the corresponding statistics as in (2.2) before. Write $Y_n = (Y_{n_1}^{(1)}, Y_{n_2}^{(2)})$ and for every pair $\mathbf{n}, \mathbf{m} \in \mathbb{N}^2$, set

$$\mathbf{n} \sim \mathbf{m} \quad \text{if } Y_{\mathbf{n}} = Y_{\mathbf{m}}.$$

In this way, equivalent subclasses (components) of \mathbb{N}^2 are indexed by labels $\ell \in \mathbb{N}^2$. This gives the random partition of \mathbb{N}^2 as the product of the two partitions determined by $Y^{(1)}$ and $Y^{(2)}$. Given the partition induced by $\{Y_i\}_{i \in [1, \mathbf{n}]}$, the law of $\{X_i\}_{i \in [1, \mathbf{n}]}$ is determined by the alternating assignment rule in both directions. This is equivalent to set, letting $\{\epsilon_\ell\}_{\ell \in \mathbb{N}^2}$ be i.i.d. random variables taking values in $\{-1, 1\}$ with equal probabilities,

$$X_{\mathbf{n}} = \epsilon_\ell \prod_{q=1}^2 (-1)^{Y_{n_q, \ell_q}^{(q)} + 1} \quad \text{if } Y_{\mathbf{n}} = \ell.$$

The so-obtained random field $\{X_{\mathbf{n}}\}_{\mathbf{n} \in \mathbb{N}^2}$ is referred to as the two-dimensional randomized Karlin model.

With a little abuse of language, for $\mathbf{n} \in \mathbb{N}^2$, we refer to $\{Y_i\}_{i \in [1, \mathbf{n}]}$ as the *first \mathbf{n} samplings*. We write $\mathbf{n} \rightarrow \infty$ for $\min(n_1, n_2) \rightarrow \infty$ and we write $\mathbf{n}^\alpha = \prod_{q=1}^2 n_q^{\alpha_q}$ and $\mathbf{L}(\mathbf{n}) = \prod_{q=1}^2 L_q(n_q)$.

The main result of this section is the following.

Theorem 2.2. *For the two-dimensional randomized Karlin model with μ_q satisfying (2.1) with $\alpha_q \in (0, 1)$ and slowly varying functions L_q for $q = 1, 2$, we have*

$$\left\{ \frac{S_{\lfloor \mathbf{n} \cdot t \rfloor}}{|\mathbf{n}|^{\alpha/2} \mathbf{L}(\mathbf{n})^{1/2}} \right\}_{t \in [0, 1]^2} \Rightarrow \sigma_\alpha \{ \mathbb{B}_t^{\alpha/2} \}_{t \in [0, 1]^2}$$

in $D([0, 1]^2)$ as $\mathbf{n} \rightarrow \infty$, with $\sigma_\alpha^2 = \prod_{q=1}^2 \Gamma(1 - \alpha_q) 2^{\alpha_q - 1}$.

2.3. Auxiliary estimates

Here we provide some useful estimates on the one-dimensional randomized Karlin model that we shall use in the proof of Theorem 2.2. Recall $Y_{n,\ell}$ in (2.2) and let

$$K_n = \sum_{\ell=1}^{\infty} \mathbf{1}_{\{Y_{n,\ell} > 0\}} \quad \text{and} \quad K_{n,r} = \sum_{\ell=1}^{\infty} \mathbf{1}_{\{Y_{n,\ell} = r\}} \quad \text{for all } r \in \mathbb{N},$$

denote the number of occupied boxes and number of boxes occupied with r balls, respectively, after n samplings. The statistics of K_n and $K_{n,r}$ (independent from ϵ_ℓ) have been studied in [16] already, where a similar model with ϵ_ℓ replaced by constant 1 was investigated. We summarize some results on K_n and $K_{n,r}$ below that will be needed later. In the sequel, Γ denotes the gamma function and for $r \geq 1$ and $\alpha \in (0, 1)$, we write

$$p_\alpha(r) = \frac{\alpha(1-\alpha) \cdots (r-1-\alpha)}{r!}.$$

Observe that $\sum_{r=1}^{\infty} p_\alpha(r) = 1$ and $\sum_{r=1}^{\infty} p_\alpha(2r-1) = 2^{\alpha-1}$.

Lemma 2.3. *Under the assumption (2.1), we have*

$$\lim_{n \rightarrow \infty} \frac{K_n}{n^\alpha L(n)} = \Gamma(1-\alpha), \tag{2.4}$$

$$\lim_{n \rightarrow \infty} \frac{K_{n,r}}{n^\alpha L(n)} = \Gamma(1-\alpha)p_\alpha(r), \tag{2.5}$$

$$\lim_{n \rightarrow \infty} \frac{\sum_{r=1}^{\infty} K_{n,2r-1}}{n^\alpha L(n)} = \Gamma(1-\alpha)2^{\alpha-1}, \tag{2.6}$$

where the convergences hold almost surely and also in L^p for all $p > 0$.

Proof. (i) For the almost sure convergence in the three limits above, see [11], Corollary 21 and discussion after Proposition 2, and [16], Theorem 9.

(ii) To prove the L^p convergence, we prove (2.4) holds in L^p for $p > 0$. This and the facts that $0 \leq K_{n,r} \leq K_n$ and $0 \leq \sum_{r \geq 1} K_{n,2r-1} \leq K_n$ then imply the L^p convergence in (2.5) and (2.6). For (2.4), it suffices to prove the uniform integrability of the sequence $(K_n^p / (n^\alpha L(n))^p)_{n \geq 1}$. This follows, writing $\Phi_n = \mathbb{E}K_n$, from the asymptotic equivalence (see [11], Proposition 17)

$$\Phi_n \sim \Gamma(1-\alpha)n^\alpha L(n) \quad \text{as } n \rightarrow \infty, \tag{2.7}$$

the fact that for every $m \in \mathbb{N}$, there exist a constant C_m , such that for all $n \in \mathbb{N}$,

$$\mathbb{E}K_n^m \leq C_m \Phi_n^m, \tag{2.8}$$

and then an application of the de la Vallée Poussin criterion for uniform integrability: $(K_n / (n^\alpha L(n)))_{n \in \mathbb{N}}$ is bounded in L^m for $m > p$. To see (2.8), we need the following lemma.

Lemma 2.4. For $n \in \mathbb{N}$, for all $k \in \mathbb{N}$ and $\ell_1, \dots, \ell_k \in \mathbb{N}$ distinct,

$$\mathbb{P}(Y_{n,\ell_1} > 0, \dots, Y_{n,\ell_k} > 0) \leq \prod_{q=1}^k \mathbb{P}(Y_{n,\ell_q} > 0).$$

Proof. To prove the desired result, it suffices to show

$$\mathbb{P}(Y_{n,\ell_k} > 0 | Y_{n,\ell_1} > 0, \dots, Y_{n,\ell_{k-1}} > 0) \leq \mathbb{P}(Y_{n,\ell_k} > 0), \tag{2.9}$$

for all $k \geq 2$ and $\ell_1, \dots, \ell_k \in \mathbb{N}$ distinct. Observe that

$$\begin{aligned} & \mathbb{P}(Y_{n,\ell_k} > 0 | Y_{n,\ell_1} > 0, \dots, Y_{n,\ell_{k-1}} > 0) \\ &= 1 - \mathbb{P}(Y_{n,\ell_k} = 0) \frac{\mathbb{P}(Y_{n,\ell_1} > 0, \dots, Y_{n,\ell_{k-1}} > 0 | Y_{n,\ell_k} = 0)}{\mathbb{P}(Y_{n,\ell_1} > 0, \dots, Y_{n,\ell_{k-1}} > 0)}. \end{aligned}$$

The ratio after $\mathbb{P}(Y_{n,\ell_k} = 0)$ is larger than one, and this yields (2.9) and hence the desired result. To see this, let $\{Y_n^*\}_{n \in \mathbb{N}}$ be another collection of i.i.d. random variables, taking values $i \in \mathbb{N} \setminus \{\ell_k\}$ with probability $p_i^* = p_i / (1 - p_{\ell_k})$. Then, the ratio above equals

$$\frac{\mathbb{P}(Y_{n,\ell_1}^* > 0, \dots, Y_{n,\ell_{k-1}}^* > 0)}{\mathbb{P}(Y_{n,\ell_1} > 0, \dots, Y_{n,\ell_{k-1}} > 0)} \geq 1. \quad \square$$

Now to obtain (2.8), it suffices to observe that

$$\begin{aligned} \mathbb{E}K_n^m &= \sum_{\ell_1, \dots, \ell_m} \mathbb{P}(Y_{n,\ell_1} > 0, \dots, Y_{n,\ell_m} > 0) \\ &\leq \sum_{k=1}^m C_{k,m} \sum_{\substack{\ell_1, \dots, \ell_k \\ \ell_i \neq \ell_j, i \neq j}} \prod_{q=1}^k \mathbb{P}(Y_{n,\ell_q} > 0) \leq \sum_{k=1}^m C_{k,m} \Phi_n^k, \end{aligned}$$

for some constants $C_{k,m} > 0$, and recall that $\Phi_n \uparrow \infty$ as $n \rightarrow \infty$. □

We also need to work with partitions generated between two times m and n , that is, the partitions generated by Y_{m+1}, \dots, Y_n . For this purpose, we introduce $Y_{m,n,\ell}^* = \sum_{i=m+1}^n \mathbf{1}_{\{Y_i = \ell\}}$,

$$K_{m,n}^* = \sum_{\ell=1}^{\infty} \mathbf{1}_{\{Y_{m,n,\ell}^* > 0\}} \quad \text{and} \quad K_{m,n,r}^* = \sum_{\ell=1}^{\infty} \mathbf{1}_{\{Y_{m,n,\ell}^* = r\}}.$$

We need the following lemma.

Lemma 2.5. *Under the assumption (2.1), with probability one for all $s, t \in [0, 1]$, $s < t$, the following limits hold:*

$$\lim_{n \rightarrow \infty} \frac{K_{\lfloor ns \rfloor, \lfloor nt \rfloor}^*}{n^\alpha L(n)} = (t - s)^\alpha \Gamma(1 - \alpha), \tag{2.10}$$

$$\lim_{n \rightarrow \infty} \frac{\sum_{i=r}^\infty K_{\lfloor ns \rfloor, \lfloor nt \rfloor, i}^*}{n^\alpha L(n)} = (t - s)^\alpha \Gamma(1 - \alpha) \sum_{i=r}^\infty p_\alpha(i), \tag{2.11}$$

$$\lim_{n \rightarrow \infty} \frac{K_{\lfloor ns \rfloor, \lfloor nt \rfloor, r}^*}{n^\alpha L(n)} = (t - s)^\alpha \Gamma(1 - \alpha) p_\alpha(r), \tag{2.12}$$

$$\lim_{n \rightarrow \infty} \frac{\sum_{i=1}^\infty K_{\lfloor ns \rfloor, \lfloor nt \rfloor, 2i-1}^*}{n^\alpha L(n)} = (t - s)^\alpha \Gamma(1 - \alpha) 2^{\alpha-1}. \tag{2.13}$$

Proof. To prove the desired results, it suffices to establish them for fixed s and t ; the results then hold for a countable dense set of $[0, 1]$ with probability one, and by continuity for all $s, t \in [0, 1]$ with probability one.

Observe that

$$K_{\lfloor ns \rfloor, \lfloor nt \rfloor}^* \stackrel{d}{=} K_{\lfloor nt \rfloor - \lfloor ns \rfloor} \quad \text{and} \quad K_{\lfloor ns \rfloor, \lfloor nt \rfloor, r}^* \stackrel{d}{=} K_{\lfloor nt \rfloor - \lfloor ns \rfloor, r} \quad \text{for all } r \in \mathbb{N}, \tag{2.14}$$

where “ $\stackrel{d}{=}$ ” stands for equality in distribution. By Lemma 2.3, it follows immediately that all convergences hold in probability. To strengthen to the almost sure sense, we apply a monotonicity argument as in [11], Proposition 2.

From now on, we fix $s, t \in [0, 1]$, $s < t$. We first prove (2.10). Let $V_n = \text{Var } K_n$ and, as before, $\Phi_n = \mathbb{E}K_n$. By [11], Lemma 1 and Proposition 17,

$$V_n \sim \Gamma(1 - \alpha)(2^\alpha - 1)n^\alpha L(n) \quad \text{as } n \rightarrow \infty.$$

Therefore, for $n_m = \lfloor m^{2/\alpha} \rfloor$, by (2.14) and the Borel–Cantelli lemma, we have

$$\lim_{n \rightarrow \infty} \frac{K_{\lfloor n_m s \rfloor, \lfloor n_m t \rfloor}^*}{\Phi_{\lfloor n_m t \rfloor - \lfloor n_m s \rfloor}} = 1 \quad \text{almost surely.}$$

Thus, by (2.7),

$$\lim_{m \rightarrow \infty} \frac{K_{\lfloor n_m s \rfloor, \lfloor n_m t \rfloor}^*}{n_m^\alpha L(n_m)} = (t - s)^\alpha \Gamma(1 - \alpha) \quad \text{almost surely.}$$

Furthermore, for m large enough, we have $\lfloor n_m s \rfloor < \lfloor n_{m+1} s \rfloor < \lfloor n_m t \rfloor < \lfloor n_{m+1} t \rfloor$, and since Φ_n is increasing,

$$\frac{K_{\lfloor n_{m+1} s \rfloor, \lfloor n_m t \rfloor}^*}{\Phi_{\lfloor n_{m+1} t \rfloor - \lfloor n_m s \rfloor}} \leq \frac{K_{\lfloor n_m s \rfloor, \lfloor n_m t \rfloor}^*}{\Phi_{\lfloor n_m t \rfloor - \lfloor n_m s \rfloor}} \leq \frac{K_{\lfloor n_m s \rfloor, \lfloor n_{m+1} t \rfloor}^*}{\Phi_{\lfloor n_m t \rfloor - \lfloor n_{m+1} s \rfloor}} \quad \text{for all } n_m \leq n \leq n_{m+1}.$$

Since

$$\lim_{m \rightarrow \infty} \frac{\Phi_{\lfloor n_{m+1}t \rfloor - \lfloor n_m s \rfloor}}{\Phi_{\lfloor n_m t \rfloor - \lfloor n_{m+1} s \rfloor}} = 1,$$

it follows that (2.10) holds with probability one. The same argument holds for (2.11), which implies (2.12). At last, (2.13) follows from (2.11) and (2.12). \square

2.4. Proof of Theorem 2.2

The main idea behind the proof is that conditioning on the underlying partitions, the partial sum S_n can be represented as a sum of independent ± 1 -valued random variables, for which limit theorems follow immediately. We illustrate this idea by first proving a central limit theorem of the model. We let $\mathcal{N}(0, \sigma^2)$ denote the Gaussian distribution with mean zero and variance σ^2 .

Proposition 2.6. *For the two-dimensional randomized Karlin model, if μ_q satisfies (2.1) with $\alpha_q \in (0, 1)$ and slowly varying functions L_q for $q = 1, 2$, then*

$$\frac{S_n}{|n|^{\alpha/2} L(n)^{1/2}} \Rightarrow \mathcal{N}(0, \sigma_\alpha^2),$$

as $n \rightarrow \infty$, with $\sigma_\alpha^2 = \prod_{q=1}^2 \Gamma(1 - \alpha_q) 2^{\alpha_q - 1}$.

Proof. Let $S_{n,\ell}$ be the sum over all the spins X_i associated to the box ℓ :

$$S_{n,\ell} = \sum_{i \in [1,n]} X_i \mathbf{1}_{\{Y_i = \ell\}}.$$

Because of the alternating assignment rule, $S_{n,\ell} \in \{-1, 0, 1\}$, and $S_{n,\ell} = 0$ if and only if the number of variables X_i associated to the box ℓ after n samplings is even. We are therefore interested in the number of boxes having an odd number of balls after n samplings. To give an expression of this number, to be denoted by \tilde{K}_n below, we first remark that the number of boxes with an odd number of balls from samplings $Y^{(q)}$ equals

$$\sum_{\ell=1}^{\infty} \mathbf{1}_{\{Y_{n,\ell}^{(q)} \text{ odd}\}} = \sum_{i=1}^{\infty} K_{n,2i-1}^{(q)},$$

where $K_{n,i}^{(q)} = \sum_{\ell=1}^{\infty} \mathbf{1}_{\{Y_{n,\ell}^{(q)} = i\}}$. It follows that

$$\tilde{K}_n = \prod_{q=1}^2 \left(\sum_{i_q=1}^{\infty} K_{n_q, 2i_q - 1}^{(q)} \right). \tag{2.15}$$

Therefore,

$$S_n | \tilde{K}_n \stackrel{d}{=} \sum_{i=1}^{\tilde{K}_n} \epsilon'_i, \tag{2.16}$$

where the left-hand side is understood as the conditional distribution of S_n given \tilde{K}_n , and on the right-hand side $\{\epsilon'_i\}_{i \in \mathbb{N}}$ are i.i.d. copies of ϵ_1 . Introduce furthermore the σ -field $\mathcal{Y} = \sigma(Y^{(1)}, Y^{(2)})$, and observe that \tilde{K}_n is \mathcal{Y} measurable. Now, by Lemma 2.3,

$$\lim_{n \rightarrow \infty} \frac{\text{Var}(S_n | \mathcal{Y})}{|n|^{\alpha} L(n)} = \lim_{n \rightarrow \infty} \prod_{q=1}^2 \frac{\sum_{i_q=1}^{\infty} K_{n_q, 2i_q-1}^{(q)}}{n_q^{\alpha_q} L_q(n_q)} = \prod_{q=1}^2 \sigma_{\alpha_q}^2 = \sigma_{\alpha}^2 \quad \text{a.s.}$$

Therefore, we obtain the conditional central limit theorem

$$\frac{S_n}{|n|^{\alpha/2} L(n)^{1/2}} \Big| \mathcal{Y} \Rightarrow \mathcal{N}(0, \sigma_{\alpha}^2) \quad \text{as } n \rightarrow \infty,$$

and the desired annealed version follows. □

To prove Theorem 2.2, we prove the convergence of finite-dimensional distributions and tightness separately.

Proof of convergence of finite-dimensional distributions. For $m \in \mathbb{N}$ fixed, consider $\lambda_1, \dots, \lambda_m \in \mathbb{R}$, and $\mathbf{t}^{(r)} = (t_1^{(r)}, t_2^{(r)}) \in [0, 1]^2$ for $r = 1, \dots, m$. Writing

$$\mathbf{n}_t = (\lfloor n_1 t_1 \rfloor, \lfloor n_2 t_2 \rfloor) \quad \text{for all } \mathbf{t} \in [0, 1]^2, \mathbf{n} \in \mathbb{N}^2, \tag{2.17}$$

we set

$$\widehat{S}_n = \sum_{r=1}^m \lambda_r S_{n_{t^{(r)}}}, \quad \mathbf{n} \in \mathbb{N}^2.$$

Similarly as before and using the Cramér–Wold device [14], Corollary 4.5, to show the convergence of finite-dimensional distributions, it suffices to show the following conditional central limit theorem:

$$\frac{\widehat{S}_n}{\sigma_{\alpha} |n|^{\alpha/2} L(n)^{1/2}} \Big| \mathcal{Y} \Rightarrow \sum_{r=1}^m \lambda_r \mathbb{B}_{t^{(r)}}^{\alpha/2},$$

as $n \rightarrow \infty$. For this purpose, we first remark that given \mathcal{Y} , \widehat{S}_n is the sum of \tilde{K}_n independent random variables corresponding to the \tilde{K}_n boxes that have at least one ball from the first n samplings, and that each such random variable is bounded by $|\lambda_1| + \dots + |\lambda_m|$ uniformly. At the same time, we know that $\tilde{K}_n \rightarrow \infty$ almost surely, as $n \rightarrow \infty$. Therefore, to establish the conditional central limit theorem it remains to show that the variance

$$\text{Var}(\widehat{S}_n | \mathcal{Y}) = \sum_{r=1}^m \sum_{r'=1}^m \lambda_r \lambda_{r'} \mathbb{E}_{\mathcal{Y}}(S_{n_{t^{(r)}}} S_{n_{t^{(r')}}})$$

converges to the corresponding one of the fractional Brownian sheet as $n \rightarrow \infty$, after normalization. Here and in the sequel, we write $\mathbb{E}_{\mathcal{Y}}(\cdot) = \mathbb{E}(\cdot | \mathcal{Y})$. This part is established in Lemma 2.7. \square

Lemma 2.7. *With the same notation as in (2.17), for all $n \in \mathbb{N}^2$, $t, s \in [0, 1]^2$,*

$$\lim_{n \rightarrow \infty} \frac{\mathbb{E}_{\mathcal{Y}}(S_{n_t} S_{n_s})}{\sigma_{\alpha}^2 |n|^{\alpha} L(n)} = \prod_{q=1}^2 \frac{1}{2} (|t_q|^{\alpha_q} + |s_q|^{\alpha_q} - |t_q - s_q|^{\alpha_q}).$$

Proof. We first consider the case of the one-dimensional model described in Section 2.1. We write, for $n \in \mathbb{N}$, $t, s \in [0, 1]$,

$$\mathbb{E}_{\mathcal{Y}}(S_{n_t} S_{n_s}) = \frac{1}{2} \mathbb{E}_{\mathcal{Y}}[S_{n_t}^2 + S_{n_s}^2 - (S_{n_t} - S_{n_s})^2], \tag{2.18}$$

where $S_n = \sum_{i=1}^n X_i$, $n_t = \lfloor nt \rfloor$, and $n_s = \lfloor ns \rfloor$. We saw in the proof of Proposition 2.6 that $\mathbb{E}_{\mathcal{Y}} S_n^2 = \sum_{i=1}^{\infty} K_{n,2i-1}$, and thus Lemma 2.3 yields that, almost surely,

$$\lim_{n \rightarrow \infty} \frac{\mathbb{E}_{\mathcal{Y}} S_{n_t}^2}{n^{\alpha} L(n)} = t^{\alpha} \sigma_{\alpha}^2. \tag{2.19}$$

For $n > n'$, by a similar argument as in the proof of Proposition 2.6, we see that

$$S_n - S_{n'} | \mathcal{Y} \stackrel{d}{=} \sum_{i=1}^{\tilde{K}_{n,n'}^*} \epsilon'_i \quad \text{with} \quad \tilde{K}_{n,n'}^* = \sum_{i=1}^{\infty} K_{n,n',2i-1}^*,$$

where $\{\epsilon'_i\}$ are i.i.d. copies of ϵ_1 . In this way,

$$\mathbb{E}_{\mathcal{Y}}(S_{n_t} - S_{n_s})^2 = \sum_{i=1}^{\infty} K_{n_t, n_s, 2i-1}^*,$$

and by (2.13),

$$\lim_{n \rightarrow \infty} \frac{\mathbb{E}_{\mathcal{Y}}(S_{n_t} - S_{n_s})^2}{n^{\alpha} L(n)} = |t - s|^{\alpha} \sigma_{\alpha}^2 \quad \text{almost surely.}$$

Combining this, (2.18) and (2.19), we have thus proved

$$\lim_{n \rightarrow \infty} \frac{\mathbb{E}_{\mathcal{Y}}(S_{n_t} S_{n_s})}{\sigma_{\alpha}^2 |n|^{\alpha} L(n)} = \frac{1}{2} (|t|^{\alpha} + |s|^{\alpha} - |t - s|^{\alpha}).$$

For the two-dimensional model, we start by introducing a different model. Let $\{\tilde{\epsilon}_k^{(q)}\}_{k \in \mathbb{N}}$, $q = 1, 2$, be i.i.d. random variables taking values ± 1 with equal probabilities and set $\tilde{\epsilon}_n = \prod_{q=1}^2 \tilde{\epsilon}_{n_q}^{(q)}$.

Now, assign

$$\tilde{X}_n = \prod_{q=1}^2 \tilde{\epsilon}_{\ell_q}^{(q)} (-1)^{Y_{n_q, \ell_q}^{(q)}} \quad \text{if } Y_n = \ell, \tag{2.20}$$

and set $\tilde{S}_n = \sum_{i \in [1, n]} \tilde{X}_i$. Although $\{\tilde{X}_n\}_{n \in \mathbb{N}^2}$ is different from $\{X_n\}_{n \in \mathbb{N}^2}$, observe that for all $i, j \in \mathbb{N}^2$,

$$(X_i, X_j) | \mathcal{Y} \stackrel{d}{=} (\tilde{X}_i, \tilde{X}_j) | \mathcal{Y}.$$

Indeed, this follows from the fact that $(\epsilon_\ell, \epsilon_{\ell'}) \stackrel{d}{=} (\tilde{\epsilon}_\ell, \tilde{\epsilon}_{\ell'})$, for all $\ell, \ell' \in \mathbb{N}^2$. (Note that $\{\epsilon_\ell\}_{\ell \in \mathbb{N}^d}$ and $\{\tilde{\epsilon}_\ell\}_{\ell \in \mathbb{N}^d}$ do not have the same joint distributions, although the fact that they have the same bivariate distributions serves our purpose.)

It follows that

$$\mathbb{E}_{\mathcal{Y}}(S_{n_t} S_{n_s}) = \mathbb{E}_{\mathcal{Y}}(\tilde{S}_{n_t} \tilde{S}_{n_s}).$$

However, $\mathbb{E}_{\mathcal{Y}}(\tilde{S}_{n_t} \tilde{S}_{n_s})$ is much easier to compute. From (2.20), \tilde{X}_i can be written as

$$\tilde{X}_i = \prod_{q=1}^2 \tilde{X}_{i_q}^{(q)} \quad \text{if } Y_i = \ell \text{ with } \tilde{X}_{i_q}^{(q)} = \tilde{\epsilon}_{\ell_q}^{(q)} (-1)^{Y_{i_q, \ell_q}^{(q)} + 1}, q = 1, 2.$$

In this way, one can write

$$\tilde{S}_n = \sum_{i \in [1, n]} \prod_{q=1}^2 \tilde{X}_{i_q}^{(q)} = \prod_{q=1}^2 \sum_{i_q=1}^{n_q} \tilde{X}_{i_q}^{(q)} = \prod_{q=1}^2 \tilde{S}_{n_q}^{(q)} \quad \text{with } \tilde{S}_n^{(q)} = \sum_{i=1}^n \tilde{X}_i^{(q)}.$$

Observe that $\{\tilde{X}_i^{(1)}\}_{i \in \mathbb{N}}$ and $\{\tilde{X}_i^{(2)}\}_{i \in \mathbb{N}}$ are independent and each $\tilde{S}_n^{(q)}$ is the partial sum of a one-dimensional Karlin model with parameter α_q . Therefore,

$$\begin{aligned} \frac{\mathbb{E}_{\mathcal{Y}}(\tilde{S}_{n_t} \tilde{S}_{n_s})}{|n|^{\alpha} L(n)} &= \prod_{q=1}^2 \frac{\mathbb{E}_{\mathcal{Y}}(\tilde{S}_{\lfloor n_q t_q \rfloor}^{(q)} \tilde{S}_{\lfloor n_q s_q \rfloor}^{(q)})}{n_q^{\alpha_q} L_q(n_q)} \\ &\rightarrow \sigma_{\alpha}^2 \prod_{q=1}^d \frac{1}{2} (|t_q|^{\alpha_q} + |s_q|^{\alpha_q} - |t_q - s_q|^{\alpha_q}) \quad \text{as } n \rightarrow \infty. \end{aligned} \quad \square$$

Proof of tightness. Applying a criterion of Bickel and Wichura [1], it suffices to establish for some $p > 0, \gamma > 1$,

$$\mathbb{E} \left(\frac{|S_m|}{|n|^{\alpha/2}} \right)^p \leq C \prod_{q=1}^2 \left(\frac{m_q}{n_q} \right)^{\gamma} \quad \text{for all } m, n \in \mathbb{N}^2, m \leq n. \tag{2.21}$$

To do so, pick $p > \max(2/\alpha_1, 2/\alpha_2)$. Recall that, given \mathcal{Y} , S_m is the sum of \tilde{K}_m independent copies of ϵ_1 . We infer

$$\mathbb{E}|S_m|^p = \mathbb{E}\mathbb{E}_{\mathcal{Y}}(|S_m|^p) \leq \mathbb{E}[C_p \mathbb{E}_{\mathcal{Y}} \tilde{K}_m^{p/2}] = C_p \mathbb{E} \tilde{K}_m^{p/2},$$

where we used Burkholder’s inequality, and C_p is a positive constant depending only on p . The expectation on the right-hand side above is then bounded from above by, recalling (2.15),

$$\mathbb{E} \tilde{K}_m^{p/2} = \prod_{q=1}^2 \mathbb{E} \left(\sum_{i_q=1}^{\infty} K_{m_q, 2i_q-1}^{(q)} \right)^{p/2} \leq \prod_{q=1}^2 \mathbb{E} (K_{m_q}^{(q)})^{p/2}.$$

Now, for each q , the expectation can be uniformly bounded by $C_q m_q^{\alpha_q p/2} L_q(m_q)^{p/2}$ for some constant $C_q > 0$ by Lemma 2.3. Therefore,

$$\mathbb{E} \left(\frac{|S_m|}{|n|^{\alpha/2}} \right)^p \leq C_p \prod_{q=1}^2 C_q \left(\frac{m_q}{n_q} \right)^{\gamma'} \left(\frac{L_q(m_q)}{L_q(n_q)} \right)^{p/2}$$

with $\gamma' = \min(\alpha_1, \alpha_2)p/2 > 1$. To conclude, we choose $\delta > 0$ such that $\gamma = \gamma' - \delta p/2 > 1$ and we apply Potter’s theorem (see [6], Theorem 1.5.6) to bound from above $L_q(m_q)/L_q(n_q)$ by $C(m_q/n_q)^{-\delta}$. The inequality (2.21) follows and we have thus proved the tightness. \square

3. Hammond–Sheffield model

In this section, we introduce the two-dimensional Hammond–Sheffield model and show that the partial-sum random field scales to a fractional Brownian sheet with Hurst index in $(1/2, 1)^2$.

3.1. One-dimensional model

We first recall the model in one dimension. Let μ be a probability measure on \mathbb{N} satisfying

$$\mu(\{n, n + 1, \dots\}) \sim n^{-\alpha} L(n)$$

with $\alpha \in (0, 1/2)$ and L a slowly varying function at infinity. Let $\{J_i\}_{i \in \mathbb{Z}}$ be i.i.d. random variables with distribution μ and consider the random graph $G = G(V, E)$ with vertex set $V = \mathbb{Z}$ and edge set $E = \{(i, i - J_i)\}_{i \in \mathbb{Z}}$. In words, for each vertex $i \in \mathbb{Z}$, a random jump J_i is sampled independently from μ and the vertex i is connected to the vertex $i - J_i$. For each vertex i , the largest connected subgraph of G containing i is a tree with an infinite number of vertices. Each such tree is referred to as a component of \mathbb{Z} . It was shown in [12] that for $\alpha \in (0, 1/2)$, the random graph G almost surely has infinitely many components, each being an infinite tree. The random forest G obtained this way induces a random partition of \mathbb{Z} , so that i and j are in the

same component, denoted by $i \sim j$, if and only if they are in the same tree. In the sequel, it is convenient to work with *ancestral lines* defined as the random sets

$$A_i = \{j \in \mathbb{Z} : \exists j = j_0 < j_1 < \dots < j_k = i, \text{ s.t. } (j_{\ell-1}, j_\ell) \in E, \ell = 1, \dots, k\} \cup \{i\},$$

for all $i \in \mathbb{Z}$. So, $i \sim j$ if and only if $A_i \cap A_j \neq \emptyset$.

We now apply the identical assignment rule. This entails that marginally $\mathbb{P}(X_i = -1) = \mathbb{P}(X_i = 1) = 1/2$, and conditioning on $\mathcal{G} = \sigma\{J_i, i \in \mathbb{Z}\}$, $X_i = X_j$ if $A_i \cap A_j \neq \emptyset$, and X_{i_1}, \dots, X_{i_k} are independent for any i_1, \dots, i_k such that A_{i_1}, \dots, A_{i_k} are mutually disjoint. The one-dimensional Hammond–Sheffield model is the stationary process $\{X_i\}_{i \in \mathbb{Z}}$ constructed this way.

The following notations and results from [12] will be used in our two-dimensional model. Let, for $k \in \mathbb{Z}$,

$$q_k = \mathbb{P}(0 \in A_k),$$

so $q_k = 0$ for $k < 0$. It is proved in [12], Lemma 3.1, that with the choice of μ above and $\alpha \in (0, 1/2)$, $\sum_{k=0}^\infty q_k^2 < \infty$, and for $S_n = \sum_{i=1}^n X_i$,

$$\text{Var}(S_n) \sim \frac{C_\alpha}{\sum_{k \geq 0} q_k^2} n^{2\alpha+1} L(n)^{-2} \quad \text{as } n \rightarrow \infty, \tag{3.1}$$

with

$$C_\alpha = (2\alpha(2\alpha + 1)\Gamma(1 - \alpha)^2\Gamma(2\alpha) \cos(\pi\alpha))^{-1} \tag{3.2}$$

(note that the constant in [12] contains a misprint that is corrected here).

We shall however need a slightly more general version when working with the two-dimensional model later. We say that $\{X_i\}_{i \in \mathbb{Z}}$ is a *generalized one-dimensional Hammond–Sheffield model* with distribution ν on \mathbb{R} if it is built using ν as the common marginal distribution instead of the symmetric law on $\{-1, 1\}$. That is, the underlying random partition is the same as before and, conditioning on the random partition, the identical assignment rule is applied ($X_i = X_j$ if $i \sim j$ and X_i and X_j are independent otherwise) with each X_i having the same marginal distribution ν .

Proposition 3.1. *For the generalized one-dimensional Hammond–Sheffield model with a centered distribution ν with bounded support,*

$$\left\{ \frac{S_{\lfloor nt \rfloor}}{n^H L(n)^{-1}} \right\}_{t \in [0,1]} \Rightarrow \left(\frac{C_\alpha}{\sum_{k \geq 0} q_k^2} \right)^{1/2} \{B_t^H\}_{t \in [0,1]}$$

in $D([0, 1])$ with $H = \alpha + 1/2$.

Remark 3.2. The results in [12] concern only ν supported on $\{-1, 1\}$. The relaxation of ν to bounded law does not affect most of the proof, which is based on a martingale central limit theorem. The boundedness is sufficient for certain ergodicity of the sequence of martingale differences ([12], Lemma 3.2, and [3], Lemma 7), and the rest of the proof would remain unchanged.

As the proof for the two-dimensional model will follow the same strategy but is much more involved, we therefore skip the proof of Proposition 3.1 here.

3.2. Two-dimensional model and main result

We now generalize Hammond–Sheffield model to two dimensions. Again the first step is to construct a random partition of \mathbb{Z}^2 . This random partition is taken as the product of independent random partitions from one-dimensional Hammond–Sheffield models, each with jump distribution μ_r , $r = 1, 2$ respectively, satisfying

$$\mu_r(\{n, n + 1, \dots\}) \sim n^{-\alpha_r} L_r(n) \quad \text{as } n \rightarrow \infty \tag{3.3}$$

with $\alpha_r \in (0, 1/2)$ and slowly varying function L_r at infinity. For $r = 1, 2$, let $\{A_i^{(r)}\}_{i \in \mathbb{Z}}$ be the ancestral lines corresponding to each random partition. In particular, $\{A_i^{(1)}\}_{i \in \mathbb{Z}}$ and $\{A_i^{(2)}\}_{i \in \mathbb{Z}}$ are independent. We then introduce the *ancestral lattices* A_i , $i \in \mathbb{Z}^2$, as

$$A_i = \{\mathbf{j} \in \mathbb{Z}^2 \mid j_1 \in A_{i_1}^{(1)}, j_2 \in A_{i_2}^{(2)}\} = A_{i_1}^{(1)} \times A_{i_2}^{(2)}.$$

For the partition of \mathbb{Z}^2 obtained by product, we have $i \sim j$ if and only if $A_i \cap A_j \neq \emptyset$. Once the random partition is given, the identical assignment rule is applied. That is, given $\{A_i\}_{i \in \mathbb{Z}^2}$, $X_i = X_j$ if $A_i \cap A_j \neq \emptyset$, and if A_{i_1}, \dots, A_{i_k} are mutually disjoint, X_{i_1}, \dots, X_{i_k} are i.i.d. with common distribution the uniform law on $\{-1, 1\}$. The so-constructed $\{X_i\}_{i \in \mathbb{Z}^2}$ is referred to as the two-dimensional Hammond–Sheffield model in the sequel.

We write for $\mathbf{n} \in \mathbb{Z}^2$, $q_{\mathbf{n}} = \mathbb{P}(\mathbf{0} \in A_{\mathbf{n}})$, and $q_n^{(r)} = \mathbb{P}(0 \in A_n^{(r)})$, $r = 1, 2$. Because of independence,

$$q_{\mathbf{n}} = q_{n_1}^{(1)} q_{n_2}^{(2)},$$

and then $\sum_{\mathbf{n} \in \mathbb{Z}^2} q_{\mathbf{n}}^2 < \infty$ when $(\alpha_1, \alpha_2) \in (0, 1/2)^2$. The main result of this section is the following functional central limit theorem, where as before $\mathbf{n}^\alpha = \prod_{r=1}^2 n_r^{\alpha_r}$ and $\mathbf{L}(\mathbf{n}) = \prod_{r=1}^2 L_r(n_r)$.

Theorem 3.3. *For the two-dimensional Hammond–Sheffield model, suppose that (3.3) holds with $\alpha_1, \alpha_2 \in (0, 1/2)$ and slowly varying functions L_1, L_2 respectively. For $S_{\mathbf{n}} = \sum_{i \in [1, \mathbf{n}]} X_i$, we have*

$$\left\{ \frac{S_{\lfloor \mathbf{n} \cdot t \rfloor}}{|\mathbf{n}|^{\mathbf{H}} \mathbf{L}(\mathbf{n})^{-1}} \right\}_{t \in [0, 1]^2} \Rightarrow \sigma_{\alpha} \{ \mathbb{B}_t^{\mathbf{H}} \}_{t \in [0, 1]^2}$$

in $D([0, 1]^2)$, as $\mathbf{n} \rightarrow \infty$, where $\mathbb{B}^{\mathbf{H}}$ is a fractional Brownian sheet with Hurst index \mathbf{H} with $H_r = \alpha_r + 1/2$, $r = 1, 2$, and

$$\sigma_{\alpha}^2 = \frac{C_{\alpha_1} C_{\alpha_2}}{\sum_{\mathbf{n} \in \mathbb{Z}^2} q_{\mathbf{n}}^2}$$

with C_{α} defined in (3.2).

Remark 3.4. Another natural extension of the Hammond–Sheffield model has been addressed in [3], where the random graph indexed by \mathbb{Z} in the original model is generalized to high dimensions by having i.i.d. jumps attached to vertices indexed by \mathbb{Z}^d and allowing each jump to take values in \mathbb{N}^d . With appropriate choice of the law of the jumps, the limit random fields therein are of different types from fractional Brownian sheets most of the time (even when fractional Brownian sheets arise in the limit, they are degenerate in the sense that at least one of the Hurst indices is either 1/2 or 1 [3], Section 5.2), and a so-called *scaling-transition* phenomenon [28, 29] occurs. In particular, the partial sum of interest therein is still over a rectangular region that increases to infinity, although for the same model (i.e., with fixed law of the jumps) various limits arise, depending on the relative growing rate of each direction of the increasing rectangle.

The rest of this section is devoted to the proof of Theorem 3.3. The strategy is to express the partial sum of the variable X_i as a weighted sum of martingale differences in the first direction and to apply a theorem of McLeish [22] for triangular arrays of martingale differences. The hard part lies in the analysis of the second direction, where we shall apply results for the generalized one-dimensional Hammond–Sheffield model.

3.3. Representation via martingale differences

Introduce for each $m \in \mathbb{Z}$, the σ -algebra of the past in the first direction $\mathcal{F}_m^{(1)} = \sigma\{X_i : i_1 < m, i_2 \in \mathbb{Z}\}$ and the operators

$$\mathcal{P}_m^{(1)}(\cdot) = \mathbb{E}(\cdot | \mathcal{F}_{m+1}^{(1)}) - \mathbb{E}(\cdot | \mathcal{F}_m^{(1)}), \quad m \in \mathbb{Z}.$$

Observe that $\mathcal{P}_m^{(1)}(Y) \in \mathcal{F}_{m+1}^{(1)}$ and $\mathbb{E}(\mathcal{P}_m^{(1)}(Y) | \mathcal{F}_m^{(1)}) = 0$ for any bounded random variable Y . Introduce

$$X_j^* = \mathcal{P}_{j_1}^{(1)}(X_j) = X_j - \mathbb{E}(X_j | \mathcal{F}_{j_1}^{(1)}), \quad \mathbf{j} \in \mathbb{Z}^2. \tag{3.4}$$

By definition, for all $j_2 \in \mathbb{Z}$, $\{X_j^*\}_{j_1 \in \mathbb{Z}}$ is a martingale-difference sequence with respect to the filtration $\{\mathcal{F}_{j_1}^{(1)}\}_{j_1 \in \mathbb{Z}}$. Denoting by $\{J_j^{(1)}\}_{j \in \mathbb{Z}}$ the random jumps in the first direction and observing that for all $\mathbf{j} \in \mathbb{Z}^2$,

$$X_j = \sum_{k \geq 1} \mathbf{1}_{\{J_{j_1}^{(1)} = k\}} X_{(j_1 - k, j_2)},$$

we obtain another representation of X_j^* as

$$X_j^* = X_j - \sum_{k \geq 1} p_k^{(1)} X_{(j_1 - k, j_2)}, \quad \mathbf{j} \in \mathbb{Z}^2, \tag{3.5}$$

where $p_k^{(1)} = \mu_1(\{k\})$, $k \in \mathbb{N}$. Recall that $q_n = 0$ if $\min(n_1, n_2) < 0$. We have the following results.

Lemma 3.5. (i) For all $m \in \mathbb{Z}$, $\mathbf{n} \in \mathbb{Z}^2$, $\mathcal{P}_m^{(1)}(X_{\mathbf{n}}) = q_{n_1-m}^{(1)} X_{(m,n_2)}^*$.

(ii) For all $\mathbf{n} \in \mathbb{Z}^2$,

$$X_{\mathbf{n}} = \sum_{m \leq n_1} q_{n_1-m}^{(1)} X_{(m,n_2)}^*,$$

where the sum converges in L^2 . Furthermore $\text{Var}(X_{\mathbf{0}}^*) = (\sum_{k \geq 0} (q_k^{(1)})^2)^{-1}$.

Proof. (i) For $m \in \mathbb{Z}$, write

$$X_{\mathbf{n}} = X_{\mathbf{n}} \mathbf{1}_{\{m \in A_{n_1}^{(1)}\}} + X_{\mathbf{n}} \mathbf{1}_{\{m \notin A_{n_1}^{(1)}\}}.$$

Observe that

$$X_{\mathbf{n}} \mathbf{1}_{\{m \in A_{n_1}^{(1)}\}} = X_{(m,n_2)} \mathbf{1}_{\{m \in A_{n_1}^{(1)}\}}$$

and $\{m \in A_{n_1}^{(1)}\}$ is independent of $\mathcal{F}_{m+1}^{(1)}$. It then follows that

$$\mathcal{P}_m^{(1)}(X_{\mathbf{n}} \mathbf{1}_{\{m \in A_{n_1}^{(1)}\}}) = \mathcal{P}_m^{(1)}(X_{(m,n_2)}) \mathbb{P}(m \in A_{n_1}^{(1)}) = X_{(m,n_2)}^* q_{n_1-m}^{(1)}.$$

On the other hand,

$$\mathbb{E}(X_{\mathbf{n}} \mathbf{1}_{\{m \notin A_{n_1}^{(1)}\}} | \mathcal{F}_{m+1}^{(1)}) = \mathbb{E}(X_{\mathbf{n}} \mathbf{1}_{\{m \notin A_{n_1}^{(1)}\}} | \mathcal{F}_m^{(1)}),$$

and thus

$$\mathcal{P}_m^{(1)}(X_{\mathbf{n}} \mathbf{1}_{\{m \notin A_{n_1}^{(1)}\}}) = 0.$$

(ii) By stationarity, it suffices to prove this for $X_{\mathbf{0}}$. For $n \in \mathbb{N}$, write

$$\sum_{m=-n}^0 \mathcal{P}_m^{(1)}(X_{\mathbf{0}}) = \sum_{m=-n}^0 q_m^{(1)} X_{(m,0)}^*.$$

Since $\{X_{(m,0)}^*\}_{m \in \mathbb{Z}}$ is a stationary martingale-difference sequence, we have $\mathbb{E}X_{(m,0)}^* = 0$ and $\mathbb{E}(X_{(m,0)}^* X_{(n,0)}^*) = 0$ if $m \neq n$. Then,

$$\mathbb{E} \left(\sum_{m=-n}^0 \mathcal{P}_m^{(1)}(X_{\mathbf{0}}) \right)^2 = \text{Var}(X_{\mathbf{0}}^*) \sum_{m=0}^n (q_m^{(1)})^2 \rightarrow \text{Var}(X_{\mathbf{0}}^*) \sum_{m \geq 0} (q_m^{(1)})^2 \quad \text{as } n \rightarrow \infty. \quad (3.6)$$

Here the assumption $\alpha_1 \in (0, 1/2)$ entails that $\sum_{m \geq 0} (q_m^{(1)})^2 < \infty$.

On the other hand, let $J_0^{(1)} \in \mathbb{N}$ denote the random jump at 0 in the first direction. One can write, in view of (3.5),

$$X_{\mathbf{0}}^* = \sum_{\ell \in \mathbb{N}} (\mathbf{1}_{\{J_0^{(1)} = \ell\}} - p_{\ell}^{(1)}) X_{(-\ell,0)}.$$

Thus,

$$\begin{aligned} \text{Var}(X_{\mathbf{0}}^*) &= \sum_{\ell, \ell' \in \mathbb{N}} \mathbb{P}(A_{-\ell}^{(1)} \cap A_{-\ell'}^{(1)} \neq \emptyset) \mathbb{E}[(\mathbf{1}_{\{J_0^{(1)}=\ell\}} - p_\ell^{(1)})(\mathbf{1}_{\{J_0^{(1)}=\ell'\}} - p_{\ell'}^{(1)})] \\ &= \sum_{\ell, \ell' \in \mathbb{N}} \mathbb{P}(A_{-\ell}^{(1)} \cap A_{-\ell'}^{(1)} \neq \emptyset) (-p_\ell^{(1)} p_{\ell'}^{(1)} + p_\ell^{(1)} \mathbf{1}_{\{\ell=\ell'\}}) \\ &= 1 - \sum_{\ell, \ell' \in \mathbb{N}} p_\ell^{(1)} p_{\ell'}^{(1)} \mathbb{P}(A_{-\ell}^{(1)} \cap A_{-\ell'}^{(1)} \neq \emptyset) = 1 - \mathbb{P}(A_0^{(1)} \cap \tilde{A}_0^{(1)} \neq \{0\}), \end{aligned}$$

where $\tilde{A}_0^{(1)}$ is an independent copy of $A_0^{(1)}$. Therefore,

$$\text{Var}(X_{\mathbf{0}}^*) = \mathbb{P}(A_0^{(1)} \cap \tilde{A}_0^{(1)} = \{0\}).$$

Combining with (3.6), we have

$$\lim_{n \rightarrow \infty} \mathbb{E} \left(\sum_{m=-n}^0 \mathcal{P}_m^{(1)}(X_{\mathbf{0}}) \right)^2 = \mathbb{P}(A_0^{(1)} \cap \tilde{A}_0^{(1)} = \{0\}) \sum_{m \geq 0} (q_m^{(1)})^2.$$

We thus have the convergence in L^2 by the fact that $\mathbb{E}X_{\mathbf{0}}^2 = 1$ and

$$\mathbb{P}(A_0^{(1)} \cap \tilde{A}_0^{(1)} = \{0\}) = \frac{1}{\sum_{m \geq 0} (q_m^{(1)})^2}. \tag{3.7}$$

Indeed, observe that

$$\sum_{m \geq 0} (q_m^{(1)})^2 = \sum_{m \geq 0} \mathbb{P}(-m \in A_0^{(1)}, -m \in \tilde{A}_0^{(1)}) = \mathbb{E}|A_0^{(1)} \cap \tilde{A}_0^{(1)}|,$$

and remark that $|A_0^{(1)} \cap \tilde{A}_0^{(1)}|$, the cardinality of intersection of the two independent ancestral lines, is a geometric random variable with rate $\theta = \mathbb{P}(A_0^{(1)} \cap \tilde{A}_0^{(1)} = \{0\})$. Thus $\mathbb{E}|A_0^{(1)} \cap \tilde{A}_0^{(1)}| = 1/\theta$, which proves (3.7). \square

Introduce $b_{n,j}^{(1)} = \sum_{i=1}^n q_{i-j}^{(1)}$, $n \in \mathbb{N}$, $j \in \mathbb{Z}$. From the preceding lemma, we have for all $\mathbf{n} \in \mathbb{N}^2$,

$$S_{\mathbf{n}} = \sum_{i \in [1, \mathbf{n}]} X_i = \sum_{j_1 \in \mathbb{Z}} b_{n_1, j_1}^{(1)} \sum_{j_2=1}^{n_2} X_j^*.$$

Further, for each $n \in \mathbb{N}$, the sequence $(\sum_{j_2=1}^n X_j^*)_{j_1 \in \mathbb{Z}}$ is a martingale-difference sequence with respect to the filtration $\{\mathcal{F}_{j_1}^{(1)}\}_{j_1 \in \mathbb{Z}}$. Denoting $(b_n^{(1)})^2 = \sum_{j \in \mathbb{Z}} (b_{n,j}^{(1)})^2$, by (3.1), we obtain

$$(b_n^{(1)})^2 \sim C_{\alpha_1} n^{2\alpha_1+1} L_1(n)^{-2} \quad \text{as } n \rightarrow \infty. \tag{3.8}$$

Now introduce similarly $b_{n,j}^{(2)} = \sum_{i=1}^n q_{i-j}^{(2)}$ and $(b_n^{(2)})^2 = \sum_{j \in \mathbb{Z}} (b_{n,j}^{(2)})^2$, for $n \in \mathbb{N}$, $j \in \mathbb{Z}$. In summary, we have shown the following.

Proposition 3.6. *In the notation above,*

$$\frac{S_n}{b_n} = \frac{1}{b_{n_1}^{(1)}} \sum_{j_1 \in \mathbb{Z}} b_{n_1, j_1}^{(1)} U_{j_1, n_2} \tag{3.9}$$

with

$$U_{j_1, n_2} = \frac{1}{b_{n_2}^{(2)}} \sum_{j_2=1}^{n_2} X_j^* \tag{3.10}$$

and

$$b_n^2 = (b_{n_1}^{(1)})^2 (b_{n_2}^{(2)})^2 \sim C_{\alpha_1} C_{\alpha_2} n^{2\alpha+1} L(n)^{-2} \quad \text{as } n \rightarrow \infty. \tag{3.11}$$

Note that again, for each $n \in \mathbb{N}$, $\{U_{j,n}\}_{j \in \mathbb{Z}}$ is a stationary martingale-difference sequence with respect to the filtration $\{\mathcal{F}_j^{(1)}\}_{j \in \mathbb{Z}}$.

3.4. A central limit theorem

Instead of proving directly the convergence of finite-dimensional distributions, we prove the following central limit theorem first, in order to better illustrate the key ideas of the proof.

Proposition 3.7. *For the two-dimensional Hammond–Sheffield model, suppose (3.3) holds with $\alpha_1, \alpha_2 \in (0, 1/2)$ and slowly varying functions L_1, L_2 , respectively. We have*

$$\frac{S_n}{b_n} \Rightarrow \mathcal{N}(0, \sigma^2) \quad \text{as } n \rightarrow \infty,$$

where $\sigma^2 = (\sum_{k \geq 0} q_k^2)^{-1}$.

The rest of this subsection is devoted to the proof of this proposition. With the representation in (3.9), by McLeish’s martingale central limit theorem [22], it suffices to show

$$\sup_{n \in \mathbb{N}^2} \mathbb{E} \left(\sup_{j \in \mathbb{Z}} \left(\frac{b_{n_1, j}^{(1)}}{b_{n_1}^{(1)}} \right)^2 U_{j, n_2}^2 \right) < \infty, \tag{3.12}$$

$$\lim_{n \rightarrow \infty} \sup_{j \in \mathbb{Z}} \left(\frac{b_{n_1, j}^{(1)}}{b_{n_1}^{(1)}} \right)^2 U_{j, n_2}^2 = 0 \quad \text{in probability} \tag{3.13}$$

and

$$\lim_{n \rightarrow \infty} \sum_{j \in \mathbb{Z}} \left(\frac{b_{n_1, j}^{(1)}}{b_{n_1}^{(1)}} \right)^2 U_{j, n_2}^2 = \mathbb{E} X_0^{*2} \quad \text{in probability.} \tag{3.14}$$

We start with the following observation.

Lemma 3.8. *For $\alpha_1 \in (0, 1/2)$, we have*

$$\lim_{n \rightarrow \infty} \sup_{j \in \mathbb{Z}} \frac{b_{n,j}^{(1)}}{b_n^{(1)}} = 0.$$

Proof. By Lemma 8 in [3], it suffices to prove that $\sum_{j \in \mathbb{Z}} (b_{n,j}^{(1)})^2 - (b_{n,j+1}^{(1)})^2 = o((b_n^{(1)})^2)$, which follows from $\sum_{j \in \mathbb{Z}} (b_{n,j}^{(1)} - b_{n,j+1}^{(1)})^2 = o((b_n^{(1)})^2)$ by the Cauchy–Schwarz inequality. To see the latter, observe that

$$\sum_{j \in \mathbb{Z}} (b_{n,j}^{(1)} - b_{n,j+1}^{(1)})^2 = \sum_{j \in \mathbb{Z}} (q_{n-j}^{(1)} - q_{-j}^{(1)})^2 \leq 2 \sum_{j \in \mathbb{Z}} (q_j^{(1)})^2 < \infty. \quad \square$$

We also need uniform bounds on the moments of $U_{j,n}$. To facilitate, we introduce a representation of $U_{j,n}$ as a weighted sum of martingale differences in the second direction. Let $\mathcal{F}_m^{(2)} = \sigma\{X_i | i_1 \in \mathbb{Z}, i_2 < m\}$ and $\mathcal{P}_m^{(2)}(\cdot) = \mathbb{E}(\cdot | \mathcal{F}_{m+1}^{(2)}) - \mathbb{E}(\cdot | \mathcal{F}_m^{(2)})$, $m \in \mathbb{Z}$. We set

$$X_n^{**} = \mathcal{P}_{n_2}^{(2)}(X_n^*) = X_n^* - \mathbb{E}(X_n^* | \mathcal{F}_{n_2}^{(2)}). \tag{3.15}$$

For all $n_1 \in \mathbb{Z}$, $(X_n^{**})_{n_2 \in \mathbb{Z}}$ is a martingale-difference sequence with respect to the filtration $(\mathcal{F}_{n_2}^{(2)})_{n_2 \in \mathbb{Z}}$. Proceeding as in Lemma 3.5, we obtain that for all $n \in \mathbb{N}^2$,

$$X_n^* = \sum_{m \leq n_2} \mathcal{P}_m^{(2)}(X_n^*) = \sum_{m \leq n_2} q_{n_2-m}^{(2)} X_{(n_1, m)}^{**},$$

where the sum converges in L^2 . We thus have, for all $n \in \mathbb{N}$, $j_1 \in \mathbb{Z}$,

$$U_{j_1, n} = \frac{1}{b_n^{(2)}} \sum_{j_2 \in \mathbb{Z}} b_{n, j_2}^{(2)} X_j^{**}. \tag{3.16}$$

Further,

$$\text{Var}(X_{\mathbf{0}}^{**}) = \frac{\text{Var}(X_{\mathbf{0}}^*)}{\sum_{k \geq 0} (q_k^{(2)})^2} = \frac{1}{\sum_{k \geq 0} q_k^2}.$$

Lemma 3.9. (i) *For all $n \in \mathbb{N}$, $\mathbb{E}U_{0,n}^2 = (\sum_{k \in \mathbb{Z}^2} q_k^2)^{-1} < \infty$.*

(ii) *For all $p \geq 1$, $\sup_{n \in \mathbb{N}} \mathbb{E}U_{0,n}^{2p} < \infty$.*

Proof. Part (i) is a direct consequence of (3.15): we have $\mathbb{E}(X_n^{**} X_m^{**}) = 0$ for $n \neq m$ and thus

$$\text{Var}(U_{0,n}) = \text{Var}(X_{\mathbf{0}}^{**}) = \frac{1}{\sum_{k \geq 0} q_k^2}.$$

For (ii), using that $(X_{(0,n)}^{**})_{n \in \mathbb{Z}}$ is a martingale-difference sequence, by Burkholder's inequality, writing $\|\cdot\|_p = (\mathbb{E}|\cdot|^p)^{1/p}$, for some constant C_p depending only on p ,

$$\begin{aligned} \|U_{0,n}\|_{2p} &\leq C_p \left\| \sum_{j \in \mathbb{Z}} \frac{(b_{n,j}^{(2)})^2}{(b_n^{(2)})^2} X_{(0,j)}^{**2} \right\|_p^{1/2} \\ &\leq C_p \left(\sum_{j \in \mathbb{Z}} \frac{(b_{n,j}^{(2)})^2}{(b_n^{(2)})^2} \|X_{\mathbf{0}}^{**2}\|_p \right)^{1/2} = C_p \|X_{\mathbf{0}}^{**}\|_{2p}. \end{aligned}$$

Then (ii) follows since $X_{\mathbf{0}}^{**}$ is bounded. □

Now, we establish the conditions of McLeish's theorem.

For (3.12), by the inequality $\sup_j |a_j| \leq \sum_j |a_j|$, the left-hand side is bounded by $\mathbb{E}U_{0,n}^2 = (\sum_{k \geq \mathbf{0}} q_k^2)^{-1} < \infty$ by Lemma 3.9(i).

For (3.13), for all $\varepsilon > 0$, one has

$$\begin{aligned} \mathbb{P}\left(\max_{j \in \mathbb{Z}} \left(\frac{b_{n_1,j}^{(1)}}{b_{n_1}^{(1)}}\right)^2 U_{j,n_2}^2 > \varepsilon\right) &\leq \sum_{j \in \mathbb{Z}} \mathbb{P}\left(\left(\frac{b_{n_1,j}^{(1)}}{b_{n_1}^{(1)}}\right)^2 U_{j,n_2}^2 > \varepsilon\right) \\ &\leq \sum_{j \in \mathbb{Z}} \left(\frac{b_{n_1,j}^{(1)}}{b_{n_1}^{(1)}}\right)^4 \frac{\mathbb{E}|U_{0,n_2}|^4}{\varepsilon^2}. \end{aligned}$$

Lemma 3.8 and Lemma 3.9(ii) then yield (3.13).

Condition (3.14) is much harder to establish. We shall prove the corresponding L^2 -convergence, which will follow from

$$\lim_{n \rightarrow \infty} \frac{1}{(b_{n_1}^{(1)})^4} \sum_{j_1, j_1' \in \mathbb{Z}} (b_{n_1, j_1}^{(1)})^2 (b_{n_1, j_1'}^{(1)})^2 \text{Cov}(U_{j_1, n_2}^2, U_{j_1', n_2}^2) = 0. \tag{3.17}$$

For this purpose, we first provide an approximation of X_j^* as follows. Introduce, for each integer $K \geq 1$, for all $j \in \mathbb{Z}^2$,

$$X_{j,K}^* = X_j - \sum_{k_1=1}^K p_k^{(1)} X_{(j_1-k, j_2)}.$$

Recalling (3.5), observe that

$$|X_{\mathbf{0}}^* - X_{\mathbf{0},K}^*| \leq \sum_{k=K+1}^{\infty} p_k^{(1)} \rightarrow 0 \quad \text{as } K \rightarrow \infty.$$

Then, define

$$U_{j_1,n,K} = \frac{1}{b_n^{(2)}} \sum_{j_2=1}^n X_{j_2,K}^* \quad \text{for } n, K \in \mathbb{N}, j_1 \in \mathbb{Z}. \tag{3.18}$$

Note that $\{U_{j,n,K}\}_{j \in \mathbb{Z}}$ for every $K, n \in \mathbb{N}$ is again a stationary martingale-difference sequence with respect to the filtration $\{\mathcal{F}_j^{(1)}\}_{j \in \mathbb{Z}}$. We shall need the following uniform bounds.

Lemma 3.10. (i) For all $p \geq 1$ and $K \geq 1$,

$$\sup_{n \in \mathbb{N}} \mathbb{E}|U_{0,n,K}|^{2p} < \infty.$$

(ii) For all $p \geq 1$,

$$\lim_{K \rightarrow \infty} \sup_{n \in \mathbb{N}} \mathbb{E}|U_{0,n} - U_{0,n,K}|^{2p} = 0.$$

Proof. This lemma can be established in the same way as for Lemma 3.9 before by proving that for all $p \geq 1$, there exists a finite constant C_p depending on p only, such that for all $n, K \in \mathbb{N}$,

$$\begin{aligned} \mathbb{E}|U_{0,n,K}|^{2p} &\leq C_p \mathbb{E}|X_{\mathbf{0},K}^*|^{2p}, \\ \mathbb{E}|U_{0,n} - U_{0,n,K}|^{2p} &\leq C_p \mathbb{E}|X_{\mathbf{0}}^* - X_{\mathbf{0},K}^*|^{2p}. \end{aligned} \quad \square$$

Now, to prove (3.17), we first show that for all $\varepsilon > 0$, one can choose $K \in \mathbb{N}$ large enough such that

$$|\text{Cov}(U_{0,n}^2, U_{j,n}^2) - \text{Cov}(U_{0,n,K}^2, U_{j,n,K}^2)| < \varepsilon \quad \text{for all } n, j \in \mathbb{N}. \tag{3.19}$$

To see this, we first bound

$$\begin{aligned} &|\mathbb{E}(U_{0,n}^2 U_{j,n}^2) - \mathbb{E}(U_{0,n,K}^2 U_{j,n,K}^2)| \\ &\leq \mathbb{E}|U_{0,n}^2 (U_{j,n}^2 - U_{j,n,K}^2)| + \mathbb{E}|(U_{0,n}^2 - U_{0,n,K}^2) U_{j,n,K}^2|. \end{aligned}$$

The first term on the right-hand side is bounded, applying the Cauchy–Schwarz inequality twice, by

$$(\mathbb{E}|U_{0,n}|^4)^{1/2} (\mathbb{E}|U_{j,n} + U_{j,n,K}|^4)^{1/4} (\mathbb{E}|U_{j,n} - U_{j,n,K}|^4)^{1/4}.$$

By Lemma 3.10, this expression converges to 0 uniformly in n , as $K \rightarrow \infty$. The second term can be treated similarly. Therefore (3.19) follows for K large enough and hence to show (3.17) it suffices to establish, for K large enough,

$$\lim_{n \rightarrow \infty} \frac{1}{(b_{n_1}^{(1)})^4} \sum_{j_1, j_1' \in \mathbb{Z}} (b_{n_1, j_1}^{(1)})^2 (b_{n_1, j_1'}^{(1)})^2 \text{Cov}(U_{j_1, n_2, K}^2, U_{j_1', n_2, K}^2) = 0. \tag{3.20}$$

For this purpose, we shall establish the following lemma.

Lemma 3.11. For all $K \in \mathbb{N}$, $\varepsilon > 0$, there exist integers $L_{K,\varepsilon}, N_{K,\varepsilon}$, such that

$$|\text{Cov}(U_{0,n,K}^2, U_{j,n,K}^2)| < \varepsilon \quad \text{for all } j > L_{K,\varepsilon}, n > N_{K,\varepsilon}.$$

Given this result, observe that the left-hand side of (3.20) without taking the limit is bounded by, writing $\sum_{j'_1 \in \mathbb{Z}} = \sum_{|j_1 - j'_1| \leq L_{K,\varepsilon}} + \sum_{|j_1 - j'_1| > L_{K,\varepsilon}}$ for each $j_1 \in \mathbb{Z}$,

$$\begin{aligned} & CL_{K,\varepsilon} \sum_{j_1 \in \mathbb{Z}} \left(\frac{b_{n_1, j_1}^{(1)}}{b_{n_1}^{(1)}} \right)^2 \sup_{j'_1 \in \mathbb{Z}} \left(\frac{b_{n_1, j'_1}^{(1)}}{b_{n_1}^{(1)}} \right)^2 + \frac{1}{(b_{n_1}^{(1)})^4} \sum_{j_1, j'_1 \in \mathbb{Z}} (b_{n_1, j_1}^{(1)})^2 (b_{n_1, j'_1}^{(1)})^2 \varepsilon \\ & \leq CL_{K,\varepsilon} \sup_{j'_1 \in \mathbb{Z}} \left(\frac{b_{n_1, j'_1}^{(1)}}{b_{n_1}^{(1)}} \right)^2 + \varepsilon, \end{aligned} \tag{3.21}$$

for all $n_2 > N_{K,\varepsilon}$. This and Lemma 3.8 give (3.20) and hence the third condition of McLeish’s central limit theorem (3.14). Therefore, the proof of Proposition 3.7 is completed. It remains to prove Lemma 3.11.

Proof of Lemma 3.11. Introduce, for each $K \in \mathbb{N}$, $j \in \mathbb{Z}$, the event

$$R_{j,K}^{(1)} = \left\{ \left(\bigcup_{i \in \{-K+1, \dots, 0\}} A_i^{(1)} \right) \cap \left(\bigcup_{i' \in \{j-K+1, \dots, j\}} A_{i'}^{(1)} \right) = \emptyset \right\}. \tag{3.22}$$

We have $\lim_{j \rightarrow \infty} \mathbb{P}(R_{j,K}^{(1)}) = 1$ for all K . This comes from

$$\begin{aligned} \mathbb{P}(A_0^{(1)} \cap A_j^{(1)} \neq \emptyset) & \leq \sum_{k \geq 0} \mathbb{P}(-k \in A_0^{(1)}, -k \in A_j^{(1)}) \\ & \leq \sum_{k \geq 0} q_k^{(1)} q_{j+k}^{(1)} \leq \left(\sum_{k \geq 0} (q_k^{(1)})^2 \right)^{1/2} \left(\sum_{k \geq 0} (q_{j+k}^{(1)})^2 \right)^{1/2}, \end{aligned}$$

and the fact that $\sum_{k \geq 0} (q_{j+k}^{(1)})^2 \rightarrow 0$ as $j \rightarrow \infty$. We now write

$$\mathbb{E}(U_{0,n,K}^2 U_{j,n,K}^2) = \mathbb{E}(U_{0,n,K}^2 U_{j,n,K}^2 \mathbf{1}_{R_{j,K}^{(1)}}) + \mathbb{E}(U_{0,n,K}^2 U_{j,n,K}^2 \mathbf{1}_{(R_{j,K}^{(1)})^c}). \tag{3.23}$$

The second term on the right-hand side, by applying the Cauchy–Schwarz inequality twice and Lemma 3.10, can be bounded uniformly in n by $C\mathbb{P}((R_{j,K}^{(1)})^c)^{1/2}$, which goes to zero as $j \rightarrow \infty$ (C is a positive constant). So, it suffices to show that the first term on the right-hand side above can be controlled to be arbitrarily close to $(\mathbb{E}U_{0,n,K}^2)^2$ for j, K large enough.

For this purpose, the key idea is to decouple the underlying partitions in the first direction $U_{0,n,K}$ and $U_{j,n,K}$. Otherwise, notice that the two are dependent for all choices of j and K . For the decoupling, first notice that the law of the partition in the first direction are determined by the law of those ancestral lines involved in the definition of $R_{j,K}^{(1)}$. To proceed we introduce a copy

of $\{A_j^{(1)}\}_{j \in \mathbb{Z}}$, denoted by $\{\tilde{A}_j^{(1)}\}_{j \in \mathbb{Z}}$, independent of the original two-dimensional Hammond–Sheffield model. Introduce the product partition \tilde{G} of \mathbb{Z}^2 as in the original model, but instead induced by $\{\tilde{A}_j^{(1)}\}_{j \in \mathbb{Z}}$ and $\{A_j^{(2)}\}_{j \in \mathbb{Z}}$. Then, define $\{\tilde{X}_j\}_{j \in \mathbb{Z}^2}$ as before on \tilde{G} by identical assignment rule. Define similarly \tilde{X}_i^* , $\tilde{X}_{i,K}^*$, $\tilde{U}_{j,n}$ and $\tilde{U}_{j,n,K}$ as before, based on \tilde{G} . These are identically distributed copies of the corresponding quantities of the original model. Define

$$\tilde{R}_{j,K}^{(1)} = \left\{ \left(\bigcup_{i \in \{-K+1, \dots, 0\}} A_i^{(1)} \right) \cap \left(\bigcup_{i' \in \{j-K+1, \dots, j\}} \tilde{A}_{i'}^{(1)} \right) = \emptyset \right\}. \tag{3.24}$$

We first remark that $\mathbb{P}(R_{j,K}^{(1)}) = \mathbb{P}(\tilde{R}_{j,K}^{(1)})$ for $j \geq K$ and

$$(U_{0,n,K}, U_{j,n,K}) | R_{j,K}^{(1)} \stackrel{d}{=} (U_{0,n,K}, \tilde{U}_{j,n,K}) | \tilde{R}_{j,K}^{(1)},$$

where each side is understood as the conditional distribution of a bivariate random vector. Therefore, we have

$$\begin{aligned} & \mathbb{E}(U_{0,n,K}^2 U_{j,n,K}^2 \mathbf{1}_{R_{j,K}^{(1)}}) \\ &= \mathbb{E}(U_{0,n,K}^2 \tilde{U}_{j,n,K}^2 \mathbf{1}_{\tilde{R}_{j,K}^{(1)}}) \\ &= \mathbb{E}(U_{0,n,K}^2 \tilde{U}_{j,n,K}^2) - \mathbb{E}(U_{0,n,K}^2 \tilde{U}_{j,n,K}^2 \mathbf{1}_{(\tilde{R}_{j,K}^{(1)})^c}) \\ &= \mathbb{E}(U_{0,n,K}^2 \tilde{U}_{0,n,K}^2) - \mathbb{E}(U_{0,n,K}^2 \tilde{U}_{j,n,K}^2 \mathbf{1}_{(\tilde{R}_{j,K}^{(1)})^c}), \end{aligned} \tag{3.25}$$

where in the last expression above, the second term is again bounded by $C\mathbb{P}((\tilde{R}_{j,K}^{(1)})^c)^{1/2}$, uniformly in n , for some positive constant C . To sum up, by (3.23) and (3.25) we arrive at the fact that there exists a constant $L_{K,\varepsilon}$ such that

$$|\mathbb{E}(U_{0,n,K}^2 U_{j,n,K}^2) - \mathbb{E}(U_{0,n,K}^2 \tilde{U}_{0,n,K}^2)| \leq \frac{\varepsilon}{2} \quad \text{for all } j > L_{K,\varepsilon}, n \in \mathbb{N}. \tag{3.26}$$

Finally, we will prove that

$$\lim_{n \rightarrow \infty} \text{Cov}(U_{n,0,K}^2, \tilde{U}_{n,0,K}^2) = 0. \tag{3.27}$$

This and (3.26) shall yield that there exists an integer $N_{K,\varepsilon}$ such that

$$|\mathbb{E}(U_{0,n,K}^2 \tilde{U}_{0,n,K}^2) - (\mathbb{E}U_{0,n,K}^2)^2| < \frac{\varepsilon}{2} \quad \text{for all } n \in N_{K,\varepsilon},$$

and complete the proof of the lemma.

It remains to show (3.27). We start by establishing a conditional central limit theorem for $U_{0,n,K}$, given the ancestral lines $A_{-K+1}^{(1)}, \dots, A_0^{(1)}$. We shall actually only need the random partition on $\{-K+1, \dots, 0\}$, denoted by $G_K^{(1)}$, induced by these ancestral lines. Recall the definition

of $U_{0,j,K}$ in (3.18). We have

$$U_{0,n,K} = \frac{1}{b_n^{(2)}} \sum_{j=1}^n X_{(0,j),K}^* = \frac{1}{b_n^{(2)}} \sum_{j_2=1}^n \left(X_{(0,j_2)} - \sum_{j_1=1}^K p_{j_1}^{(1)} X_{(-j_1,j_2)} \right) \equiv \frac{1}{b_n^{(2)}} \sum_{j=1}^n \mathbb{X}_j, \quad (3.28)$$

where we introduce $\mathbb{X}_j = X_{(0,j),K}^*$ to simplify the notation. Note that \mathbb{X} depends on K .

Here we need the crucial remark that, given $G_K^{(1)}$, $\{\mathbb{X}_i\}_{i \in \mathbb{Z}}$ is a generalized one-dimensional Hammond–Sheffield model, with a marginal law as the conditional law of $X_{\mathbf{0},K}^*$ given $G_K^{(1)}$, and hence with bounded support (Remark 3.2). To see this, the second expression of $U_{0,n,K}$ in (3.28) is more convenient: by definition of the two-dimensional model, it suffices to examine the partition of $\{-K + 1, \dots, 0\} \times \mathbb{N}$. Recall that the product partition is obtained by Cartesian products. It then follows that $\mathbb{X}_i \equiv \mathbb{X}_j$ if $i \sim j$ with respect to the random partition $G^{(2)}$ of the second direction of the model, and otherwise \mathbb{X}_i and \mathbb{X}_j are i.i.d. Note that this observation remains true if we condition on $G_K^{(1)}$ first; the marginal law will depend on $G_K^{(1)}$, but remains bounded. Then, Proposition 3.1 tells that

$$\left\{ \frac{1}{b_n^{(2)}} \sum_{i=1}^{\lfloor nt \rfloor} \mathbb{X}_i \right\}_{t \in [0,1]} \Big| G_K^{(1)} \Rightarrow \sigma_K \{ \mathbb{B}_t^H \}_{t \in [0,1]}, \quad (3.29)$$

where $H = \alpha_2 + 1/2$ and

$$\sigma_K^2 = \frac{\mathbb{E}(\mathbb{X}_1^2 | G_K^{(1)})}{\sum_{k \geq 0} (q_k^{(2)})^2} = \frac{\mathbb{E}(X_{\mathbf{0},K}^{*2} | G_K^{(1)})}{\sum_{k \geq 0} (q_k^{(2)})^2}.$$

See the [Appendix](#) for our notations for conditional limit theorems. We only need $t = 1$ to deal with $U_{0,n,K}$ in the central limit theorem here, but for the proof of finite-dimensional distributions later, we shall need the above conditional functional central limit theorem. In particular, (3.29) yields that

$$U_{0,n,K} | G_K^{(1)} \Rightarrow \sigma_K \cdot \mathcal{N}(0, 1).$$

Introduce similarly $\tilde{G}_K^{(1)}$ based on $\{\tilde{A}_{-K+1}^{(1)}, \dots, \tilde{A}_0^{(1)}\}$. By the same approach described above, we can show that

$$(U_{0,n,K}, \tilde{U}_{0,n,K}) | G_K^{(1)}, \tilde{G}_K^{(1)} \Rightarrow (\sigma_K Z, \tilde{\sigma}_K \tilde{Z}), \quad (3.30)$$

where Z and \tilde{Z} are two independent standard normal random variables and

$$\tilde{\sigma}_K^2 = \frac{\mathbb{E}(\tilde{X}_{\mathbf{0},K}^{*2} | \tilde{G}_K^{(1)})}{\sum_{k \geq 0} (q_k^{(2)})^2}.$$

To establish the joint convergence, by the Cramér–Wold device it suffices to consider, for all $a, b \in \mathbb{R}$,

$$aU_{0,n,K} + b\tilde{U}_{0,n,K} | G_K^{(1)}, \tilde{G}_K^{(1)} \stackrel{d}{=} \frac{1}{b_n^{(2)}} \sum_{i=1}^n (aX_{(0,i),K}^* + b\tilde{X}_{(0,i),K}^*) | G_K^{(1)}, \tilde{G}_K^{(1)},$$

where $\tilde{X}_{(0,i),K}^*$ is defined similarly as $X_{(0,i),K}^*$, and the two are assumed to be conditionally independent given $\{A_j^{(2)}\}_{j \in \mathbb{Z}}$. Again, given $G_K^{(1)}$ and $\tilde{G}_K^{(1)}$, the process

$$\{\bar{X}_i\}_{i \in \mathbb{N}} \equiv \{aX_{(0,i),K}^* + b\tilde{X}_{(0,i),K}^*\}_{i \in \mathbb{N}}$$

is a generalized one-dimensional Hammond–Sheffield model. The normalized partial sum $\sum_{i=1}^n \bar{X}_i / b_n^{(2)}$ then converges to a normal distribution, with variance equal to

$$\text{Var}(\bar{X}_1 | G_K^{(1)}, \tilde{G}_K^{(1)}) = \mathbb{E}[(aX_{(0,i),K}^* + b\tilde{X}_{(0,i),K}^*)^2 | G_K^{(1)}, \tilde{G}_K^{(1)}] = a^2\sigma_K^2 + b^2\tilde{\sigma}_K^2.$$

Hence, (3.30) follows as before by Proposition 3.1.

As a consequence of (3.30), we arrive at

$$(U_{0,n,K}, \tilde{U}_{0,n,K}) \Rightarrow (\sigma_K Z, \tilde{\sigma}_K \tilde{Z}) \quad \text{as } n \rightarrow \infty,$$

where now σ_K and $\tilde{\sigma}_K$ are random variables, and all four random variables on the right-hand side are independent. By the boundedness of $\sigma_K, \tilde{\sigma}_K$ and the uniform integrability of $U_{n,0,K}^4$ and $\tilde{U}_{n,0,K}^4$, it follows that

$$\lim_{n \rightarrow \infty} \text{Cov}(U_{n,0,K}^2, \tilde{U}_{n,0,K}^2) = \text{Cov}(\sigma_K^2 Z^2, \tilde{\sigma}_K^2 \tilde{Z}^2) = 0.$$

This completes the proof of (3.27). □

3.5. Proof of Theorem 3.3

Proof of convergence of finite-dimensional distributions. We use Cramér–Wold device. For $m \in \mathbb{N}$, let $\lambda_1, \dots, \lambda_m \in \mathbb{R}$ and $\mathbf{t}^{(1)}, \dots, \mathbf{t}^{(m)} \in [0, 1]^2$ be fixed. For $\mathbf{n} \in \mathbb{N}^2$, denote $\mathbf{n}_{\mathbf{t}^{(1)}}, \dots, \mathbf{n}_{\mathbf{t}^{(m)}} \in \mathbb{N}^2$ as before in (2.17) and to shorten the notation, the two coordinates of $\mathbf{n}_{\mathbf{t}^{(r)}}$ are denoted by $n_1(r) = \lfloor n_1 t_1^{(r)} \rfloor$ and $n_2(r) = \lfloor n_2 t_2^{(r)} \rfloor$ respectively, $\mathbf{t}^{(r)}$ being fixed. We have for all $\mathbf{n} \in \mathbb{N}^2$,

$$\frac{1}{b_n} \sum_{r=1}^m \lambda_r S_{\mathbf{n}_{\mathbf{t}^{(r)}}} = \frac{1}{b_{n_1}^{(1)}} \sum_{j_1 \in \mathbb{Z}} \sum_{r=1}^m \lambda_r b_{n_1(r), j_1}^{(1)} U_{j_1, n_2(r)},$$

where $U_{j_1, n_2(r)}$ is defined as in (3.10). One can show as before that

$$\left\{ \sum_{r=1}^m \lambda_r b_{n_1(r), j}^{(1)} U_{j, n_2(r)} \right\}_{j \in \mathbb{Z}}$$

is a martingale-difference sequence with respect to $\{\mathcal{F}_j^{(1)}\}_{j \in \mathbb{Z}}$. Therefore, we apply the central limit theorem of McLeish as in Section 3.4. The two conditions corresponding to (3.12) and (3.13) can be verified similarly as before. The third condition (3.14) becomes

$$\lim_{n_1 \rightarrow \infty} \frac{1}{(b_{n_1}^{(1)})^2} \sum_{j \in \mathbb{Z}} \left(\sum_{r=1}^m \lambda_r b_{n_1(r), j_1}^{(1)} U_{j, n_2(r)} \right)^2 = \frac{\text{Var}(\sum_{r=1}^m \lambda_r \mathbb{B}_{t^{(r)}}^{\mathbf{H}})}{(\sum_{k \geq \mathbf{0}} q_k^2)^2} \quad \text{in probability.}$$

This shall follow from

$$\begin{aligned} & \lim_{n_1 \rightarrow \infty} \frac{1}{(b_{n_1}^{(1)})^2} \sum_{j \in \mathbb{Z}} b_{n_1(r), j}^{(1)} b_{n_1(r'), j}^{(1)} U_{j, n_2(r)} U_{j, n_2(r')} \\ &= \frac{\text{Cov}(\mathbb{B}_{t^{(r)}}^{\mathbf{H}}, \mathbb{B}_{t^{(r')}}^{\mathbf{H}})}{(\sum_{k \geq \mathbf{0}} q_k^2)^2} \quad \text{in probability,} \end{aligned} \tag{3.31}$$

for all $r, r' \in \{1, \dots, m\}$. We do so again by computing the L^2 -convergence. Remark first that

$$\begin{aligned} & \mathbb{E} \left(\frac{1}{(b_{n_1}^{(1)})^2} \sum_{j \in \mathbb{Z}} b_{n_1(r), j}^{(1)} b_{n_1(r'), j}^{(1)} U_{j, n_2(r)} U_{j, n_2(r')} \right) \\ &= \frac{1}{(b_{n_1}^{(1)})^2} \sum_{j_1 \in \mathbb{Z}} b_{n_1(r), j_1}^{(1)} b_{n_1(r'), j_1}^{(1)} \frac{1}{(b_{n_2}^{(2)})^2} \sum_{j_2 \in \mathbb{Z}} b_{n_2(r), j_2}^{(2)} b_{n_2(r'), j_2}^{(2)} \text{Var}(X_{\mathbf{0}}^{**}) \\ &\sim \text{Cov}(\mathbb{B}_{t^{(r)}}^{\mathbf{H}}, \mathbb{B}_{t^{(r')}}^{\mathbf{H}}) \text{Var}(X_{\mathbf{0}}^{**}) \quad \text{as } n \rightarrow \infty, \end{aligned}$$

where $X_{\mathbf{0}}^{**}$ is defined as in (3.15) and the asymptotic follows from the identity

$$\begin{aligned} \sum_{j \in \mathbb{Z}} b_{n(r), j}^{(m)} b_{n(r'), j}^{(m)} &= \frac{1}{2} [(b_{n(r)}^{(m)})^2 + (b_{n(r')}^{(m)})^2 - |b_{n(r)}^{(m)} - b_{n(r')}^{(m)}|^2] \\ &= \frac{1}{2} [(b_{n(r)}^{(m)})^2 + (b_{n(r')}^{(m)})^2 - (b_{|n(r)-n(r')|}^{(m)})^2], \quad m = 1, 2, \end{aligned}$$

and (3.8). Therefore, to show (3.31), it suffices to prove, as a counterpart of (3.20),

$$\begin{aligned} & \lim_{n \rightarrow \infty} \frac{1}{(b_{n_1}^{(1)})^4} \sum_{j, j' \in \mathbb{Z}} b_{n_1(r), j}^{(1)} b_{n_1(r'), j}^{(1)} b_{n_1(r), j'}^{(1)} b_{n_1(r'), j'}^{(1)} \\ & \times \text{Cov}(U_{j, n_2(r), K} U_{j, n_2(r'), K}, U_{j', n_2(r), K} U_{j', n_2(r'), K}) = 0, \end{aligned}$$

which, as in (3.21), shall follow from the following lemma. The proof of convergence of finite-dimensional distributions is thus completed. \square

Lemma 3.12. *For all $K \in \mathbb{N}$, $\varepsilon > 0$, there exists $L_{K, \varepsilon}, N_{K, \varepsilon}$, such that*

$$|\text{Cov}(U_{0, n_2(r), K} U_{0, n_2(r'), K}, U_{j, n_2(r), K} U_{j, n_2(r'), K})| < \varepsilon \quad \text{for all } j > L_{K, \varepsilon}, n_2 > N_{K, \varepsilon}.$$

Proof. By the same idea as in the proof of Lemma 3.11, it suffices to show

$$\lim_{n_2 \rightarrow \infty} \text{Cov}(U_{0,n_2(r),K} U_{0,n_2(r'),K}, \tilde{U}_{0,n_2(r),K} \tilde{U}_{0,n_2(r'),K}) = 0.$$

As a consequence of (3.29), instead of (3.30) we now have

$$\begin{aligned} & (U_{0,n_2(r),K}, U_{0,n_2(r'),K}, \tilde{U}_{0,n_2(r),K}, \tilde{U}_{0,n_2(r'),K}) | G_K^{(1)}, \tilde{G}_K^{(1)} \\ & \Rightarrow (\sigma_K \mathbb{B}_{t_2^{(r)}}^{H_2}, \sigma_K \mathbb{B}_{t_2^{(r')}}^{H_2}, \tilde{\sigma}_K \tilde{\mathbb{B}}_{t_2^{(r)}}^{H_2}, \tilde{\sigma}_K \tilde{\mathbb{B}}_{t_2^{(r')}}^{H_2}), \end{aligned}$$

where \mathbb{B}^{H_2} and $\tilde{\mathbb{B}}^{H_2}$ are i.i.d. copies of fractional Brownian motion with Hurst index $H_2 = \alpha_2 + 1/2$, σ_K and $\tilde{\sigma}_K$ are as before. This completes the proof. □

Proof of tightness. Again, applying Bickel–Wichura’s criterion [1] and using (3.11), the tightness will follow from the existence of a real $\gamma > 1$ such that

$$\mathbb{E} \left(\frac{S_m}{b_n} \right)^2 \leq C \prod_{q=1}^2 \left(\frac{m_q}{n_q} \right)^\gamma \quad \text{for all } m \leq n.$$

Let $\delta > 0$ be such that $\gamma = 2 \min(\alpha_1, \alpha_2) + 1 - \delta > 1$. Using the representation of S_m as in (3.9) and the representation of $U_{0,n}$ as in (3.16), applying Burkholder’s inequality twice, we get

$$\mathbb{E} \left(\frac{S_m}{b_n} \right)^2 \leq C \left(\frac{b_{m_1}^{(1)}}{b_{n_1}^{(1)}} \right)^2 \left(\frac{b_{m_2}^{(2)}}{b_{n_2}^{(2)}} \right)^2 \|X_{\mathbf{0}}^{**}\|_2^2 \leq C \prod_{q=1}^2 \left(\frac{m_q}{n_q} \right)^{2\alpha_q+1} \frac{L_q(n_q)}{L_q(m_q)}.$$

We obtain the desired result by Potter’s bound $L_q(n_q)/L_q(m_q) \leq C(m_q/n_q)^{-\delta}$. □

4. Combining Hammond–Sheffield model and Karlin model

In this section, we combine a one-dimensional Hammond–Sheffield model and a one-dimensional randomized Karlin model together, and show that the combined model converges weakly to a fractional Brownian sheet with Hurst indices $H_1 \in (1/2, 1)$ and $H_2 \in (0, 1/2)$.

4.1. Model and main result

Consider two random partitions from the one-dimensional Hammond–Sheffield model and the randomized Karlin model, respectively. Assume the two random partitions are independent. Namely, let $G^{(1)} = G(E, V)$ be the underlying random forest structure of the Hammond–Sheffield model generated by a distribution μ_1 , and let $\{A_j^{(1)}\}_{j \in \mathbb{Z}}$ be the associated ancestral lines. Let $\{Y_j^{(2)}\}_{j \in \mathbb{N}}$ be i.i.d. random variables with common distribution μ_2 . Suppose $\mu_i, i = 1, 2$ are probability measures on \mathbb{N} satisfying (3.3) and (2.1) respectively with $\alpha_1 \in (0, 1/2)$ and

$\alpha_2 \in (0, 1)$. Assume $G^{(1)}$ and $\{Y_j^{(2)}\}_{j \in \mathbb{N}}$ are independent. Now, consider the product of the two random partitions. This is the random partition of $\mathbb{Z} \times \mathbb{N}$ determined by

$$i \sim j \text{ if and only if } A_{i_1}^{(1)} \cap A_{j_1}^{(1)} \neq \emptyset \text{ and } Y_{i_2}^{(2)} = Y_{j_2}^{(2)}.$$

Next, given the partition, we apply the identical assignment rule in the first direction, and the alternating assignment rule in the second (see Figure 1, right). Given a collection of components determined by $G^{(1)}$ and $\{Y_n^{(2)}\}_{n \in \mathbb{N}}$, we assign values $X_{\mathcal{C}} = \{X_j\}_{j \in \mathcal{C}}$ as follows. Let $\{\epsilon_{\mathcal{C}}\}_{\mathcal{C}}$ be a collection of i.i.d. random variables taking values in $\{-1, 1\}$ with equal probabilities, indexed by different components \mathcal{C} . For each \mathcal{C} fixed, express this as

$$\mathcal{C} = \mathcal{C}^{(1)} \times \{j_{\ell}^{(2)}\}_{\ell \in \mathbb{N}} \quad \text{with } 1 \leq j_1^{(2)} \leq j_2^{(2)} \leq \dots,$$

and set

$$X_j = (-1)^{\ell+1} \epsilon_{\mathcal{C}} \quad \text{for } j = (j_1, j_2) \in \mathcal{C}, j_2 = j_{\ell}^{(2)}.$$

The random field $\{X_j\}_{j \in \mathbb{Z} \times \mathbb{N}}$ constructed this way is referred to as the two-dimensional combined model. The main result of this section is the following invariance principle for $S_n = \sum_{i \in [1, n]} X_i$.

Theorem 4.1. *For the two-dimensional combined model with $\alpha_1 \in (0, 1/2)$, $\alpha_2 \in (0, 1)$, and slowly varying functions L_1, L_2 respectively,*

$$\left\{ \frac{S_{\lfloor n \cdot t \rfloor}}{n_1^{H_1} n_2^{H_2} L_1(n_1)^{-1} L_2(n_2)^{1/2}} \right\}_{t \in [0, 1]^2} \Rightarrow \sigma_{\alpha} \{ \mathbb{B}_t^H \}_{t \in [0, 1]^2}$$

in $D([0, 1]^2)$, as $n \rightarrow \infty$, with $H_1 = \alpha_1 + 1/2 \in (1/2, 1)$, $H_2 = \alpha_2/2 \in (0, 1/2)$, and

$$\sigma_{\alpha}^2 = \frac{C_{\alpha_1} \Gamma(1 - \alpha_2) 2^{\alpha_2 - 1}}{\sum_{j \geq 0} (q_j^{(1)})^2},$$

for C_{α_1} defined in (3.2) and $q_j^{(1)} = \mathbb{P}(0 \in A_j^{(1)})$, $j \in \mathbb{Z}$.

4.2. Proof of Theorem 4.1

The proof follows the same strategy as for the two-dimensional Hammond–Sheffield model in Sections 3.3 and 3.4. We first introduce the sequence $\{X_j^*\}_{j \in \mathbb{Z}}$ defined as in (3.4) by

$$X_j^* = X_j - \mathbb{E}(X_j | \mathcal{F}_{j_1}^{(1)}), \quad j \in \mathbb{Z},$$

where $\mathcal{F}_j^{(1)} = \sigma\{X_i | i_1 < j, i_2 \in \mathbb{N}\}$. Note that, to draw a parallel with Sections 3.3 and 3.4, we keep the same notation but the variables X_j^* here are different from the preceding section since the dependence in the second direction is given by a partition from the Karlin model.

Nevertheless, for any $j_2 \in \mathbb{N}$, the sequence $\{X_j\}_{j_1 \in \mathbb{Z}}$ is a martingale-difference sequence with respect to $\{\mathcal{F}_{j_1}^{(1)}\}_{j_1 \in \mathbb{Z}}$. So, Lemma 3.5 remains valid here (the proof is exactly the same) and we thus have

$$S_n = \sum_{j_1 \in \mathbb{Z}} b_{n_1, j_1}^{(1)} \sum_{j_2=1}^{n_2} X_j^*$$

with $b_{n,j}^{(1)} = \sum_{k=1}^n q_{k-j}^{(1)}$ defined as before. Recall from (3.8) that

$$(b_n^{(1)})^2 = \sum_{j \in \mathbb{Z}} (b_{n,j}^{(1)})^2 \sim C_{\alpha_1} n^{2\alpha_1+1} L_1(n)^{-2} \quad \text{as } n \rightarrow \infty.$$

Because of the alternating assignment rule in the second direction, we need to consider the number $\tilde{K}_n^{(2)} = \sum_{i=1}^{\infty} K_{n,2i-1}^{(2)}$ of odd-occupancy boxes, that is the number of values appearing an odd number of times among $\{Y_1^{(2)}, \dots, Y_n^{(2)}\}$. Recall from Lemma 2.3 that

$$(a_n^{(2)})^2 = \mathbb{E} \tilde{K}_n^{(2)} \sim \Gamma(1 - \alpha_2) 2^{\alpha_2-1} n^{\alpha_2} L_2(n) \quad \text{as } n \rightarrow \infty.$$

This time, for all $n \in \mathbb{N}^2$, we can write

$$\frac{S_n}{b_{n_1}^{(1)} a_{n_2}^{(2)}} = \frac{1}{b_{n_1}^{(1)}} \sum_{j \in \mathbb{Z}} b_{n_1, j_1}^{(1)} U_{j_1, n_2} \quad \text{with } U_{j_1, n_2} = \frac{1}{a_{n_2}^{(2)}} \sum_{i=1}^{n_2} X_j^*.$$

This is the counterpart of Proposition 3.6, representing the normalized partial sum of interest as a weighted linear process with stationary martingale-difference innovations.

We then introduce, for all $K \geq 1$, the approximations

$$X_{j,K}^* = X_j - \sum_{k=1}^K p_k^{(1)} X_{(j_1-k, j_2)} \quad \text{and} \quad U_{j_1, n, K} = \frac{1}{a_n^{(2)}} \sum_{j_2=1}^n X_{j,K}^*,$$

for all $j_1 \in \mathbb{Z}, j_2 \in \mathbb{N}, n \in \mathbb{N}$.

Proof of convergence of finite-dimensional distributions. This can be done as in Section 3.4 by the use of Cramér–Wold device and McLeish’s theorem [22]. For this purpose, we only need to show that the conclusions of Lemmas 3.9, 3.10, 3.11, and 3.12 are still valid with respect to the newly defined $X_j^*, X_{j,K}^*, U_{j_1, n_2}$ and $U_{j_1, n_2, K}$. For the sake of convenience, we restate Lemmas 3.9 and 3.10 in Lemma 4.2 below, and restate Lemma 3.11 in Lemma 4.3 below. The core arguments of Lemma 3.12 are all in Lemma 3.11 and we therefore omit the proof. \square

Lemma 4.2. (i) For all $n \in \mathbb{N}, \mathbb{E}U_{0,n}^2 = (\sum_{k \geq 0} q_k^2)^{-1} < \infty$.

(ii) For all $p \in \mathbb{N}$ and $K \in \mathbb{N}, \mathbb{E}U_{0,n}^{2p}$ and $\mathbb{E}U_{0,n,K}^{2p}$ are uniformly bounded.

(iii) For all $p \geq 1, \lim_{K \rightarrow \infty} \sup_{n \in \mathbb{N}} \mathbb{E}|U_{0,n} - U_{0,n,K}|^{2p} = 0$.

Proof. Denoting by $\mathcal{G}^{(1)}$ and $\mathcal{Y}^{(2)}$ the σ -fields generated by $G^{(1)}$ and $\{Y_j^{(2)}\}_{j \in \mathbb{N}}$ respectively, as for (2.16), we see that for all $n \in \mathbb{N}$, $j_1 \in \mathbb{N}$,

$$\sum_{j_2=1}^n X_j^* \Big| \mathcal{G}^{(1)}, \mathcal{Y}^{(2)} \stackrel{d}{=} \sum_{i=1}^{\tilde{K}_n^{(2)}} \epsilon'_i \Big| \mathcal{G}^{(1)},$$

where the random variables $\{\epsilon'_i\}_{i \in \mathbb{N}}$ are conditionally independent given $\mathcal{G}^{(1)}$, independent of $\mathcal{Y}^{(2)}$, and for all $i \in \mathbb{N}$, the conditional distribution $\epsilon'_i \Big| \mathcal{G}^{(1)}$ is the same as the conditional distribution $X_{\mathbf{0}}^* \Big| \mathcal{G}^{(1)}$. Note in the identity above, without the conditioning on $\mathcal{G}^{(1)}$, the $\{\epsilon'_i\}_{i \in \mathbb{N}}$ on the right-hand side are no longer independent. We can thus write

$$\begin{aligned} \mathbb{E}(U_{0,n}^2 \Big| \mathcal{G}^{(1)}, \mathcal{Y}^{(2)}) &= \frac{1}{(a_n^{(2)})^2} \mathbb{E} \left[\left(\sum_{j_2=1}^n X_j^* \right)^2 \Big| \mathcal{G}^{(1)}, \mathcal{Y}^{(2)} \right] \\ &= \frac{1}{(a_n^{(2)})^2} \mathbb{E} \left[\left(\sum_{i=1}^{\tilde{K}_n^{(2)}} \epsilon'_i \right)^2 \Big| \mathcal{G}^{(1)}, \mathcal{Y}^{(2)} \right] = \frac{\tilde{K}_n^{(2)}}{(a_n^{(2)})^2} \mathbb{E}(\epsilon_0'^2 \Big| \mathcal{G}^{(1)}). \end{aligned}$$

Thus

$$\mathbb{E}(U_{0,n}^2) = \text{Var}(X_{\mathbf{0}}^*) < \infty \quad \text{uniformly in } n.$$

This proves the first part.

For the second part, for all $p \geq 1$, by Burkholder's inequality we have

$$\mathbb{E}(U_{0,n}^{2p} \Big| \mathcal{G}^{(1)}, \mathcal{Y}^{(2)}) \leq C_p \left(\frac{\tilde{K}_n^{(2)}}{(a_n^{(2)})^2} \right)^p \mathbb{E}(X_{\mathbf{0}}^{*2p} \Big| \mathcal{G}^{(1)}).$$

Note that $\mathbb{E}(\tilde{K}_n^{(2)} / (a_n^{(2)})^2)^p$ is uniformly bounded by Lemma 2.3. Similarly,

$$\mathbb{E}(U_{0,n,K}^{2p} \Big| \mathcal{G}^{(1)}, \mathcal{Y}^{(2)}) \leq C_p \left(\frac{\tilde{K}_n^{(2)}}{(a_n^{(2)})^2} \right)^p \mathbb{E}(X_{\mathbf{0},K}^{*2p} \Big| \mathcal{G}^{(1)}).$$

For the third part, we have

$$\mathbb{E}(|U_{0,n,K} - U_{0,n}|^{2p} \Big| \mathcal{G}^{(1)}, \mathcal{Y}^{(2)}) \leq C_p \left(\frac{\tilde{K}_n^{(2)}}{(a_n^{(2)})^2} \right)^p \mathbb{E}(|X_{\mathbf{0},K}^* - X_{\mathbf{0}}^*|^{2p} \Big| \mathcal{G}^{(1)}) \rightarrow 0,$$

as $K \rightarrow \infty$. □

Lemma 4.3. For all $K \in \mathbb{N}$, $\varepsilon > 0$, there exists integers $L_{K,\varepsilon}$, $N_{K,\varepsilon}$, such that

$$|\text{Cov}(U_{0,n,K}^2, U_{j,n,K}^2)| < \varepsilon \quad \text{for all } j > L_{K,\varepsilon}, n > N_{K,\varepsilon}.$$

Proof. To proceed we introduce a copy of $\{A_j^{(1)}\}_{j \in \mathbb{Z}}$, denoted by $\{\tilde{A}_j^{(1)}\}_{j \in \mathbb{Z}}$, independent of the original model and we defined a new field $\{\tilde{X}_j\}_{j \in \mathbb{Z}}$ based on the combined model involving $\{\tilde{A}_j^{(1)}\}_{j \in \mathbb{Z}}$ and the same $\{Y_j^{(2)}\}_{j \in \mathbb{N}}$ as the original model. Then $\tilde{X}_j, \tilde{X}_j^*, \tilde{X}_{j,K}^*, \tilde{U}_{j,n}$, and $\tilde{U}_{j,n,K}$ are defined as the corresponding statistics of the combined model based on $\{\tilde{A}_j^{(1)}\}_{j \in \mathbb{Z}}$ and $\{Y_j^{(2)}\}_{j \in \mathbb{N}}$. In particular, these random variables are identically distributed as the variables $X_j, X_j^*, X_{j,K}^*, U_{0,n}$ and $U_{0,n,K}$, respectively, and they are conditionally independent from them given $\mathcal{Y}^{(2)}$. As in the proof of Lemma 3.11, observe that

$$(U_{0,n,K}, U_{j,n,K}) | R_{j,K}^{(1)} \stackrel{d}{=} (U_{0,n,K}, \tilde{U}_{j,n,K}) | \tilde{R}_{j,K}^{(1)},$$

for $R_{j,K}^{(1)}$ and $\tilde{R}_{j,K}^{(1)}$ defined as in (3.22) and (3.24). Therefore, we see that to prove the desired result it suffices to show that for all $K \geq 1$,

$$\lim_{n \rightarrow \infty} \text{Cov}(U_{0,n,K}^2, \tilde{U}_{0,n,K}^2) = 0,$$

corresponding to (3.27). Let $G_K^{(1)}$ be the random partition of $\{-K + 1, \dots, 0\}$ induced by $G^{(1)}$ and note that

$$U_{0,n,K} | G_K^{(1)} \stackrel{d}{=} \frac{1}{a_{n_2}^{(2)}} \sum_{i=1}^n X_{(0,i),K}^* | G_K^{(1)} \equiv \frac{1}{a_{n_2}^{(2)}} \sum_{i=1}^n \mathbb{X}_i | G_K^{(1)},$$

where again we write $\mathbb{X}_i = X_{(0,i),K}^*$ for the sake of simplicity. Here, conditionally given $G_K^{(1)}$, $\{\mathbb{X}_i\}_{i \in \mathbb{N}}$ is a generalized one-dimensional randomized Karlin model. Indeed, for $i \approx j$ with respect to $\{Y_j^{(2)}\}_{j \in \mathbb{N}}$, the random variables \mathbb{X}_i and \mathbb{X}_j are conditionally independent given $G_K^{(1)}$ and $\mathcal{Y}^{(2)}$, and for $i \sim j$, letting ℓ denote the number of integers in the component between i and j (say $i < j$ without loss of generality, so $\ell = \{k : i < k < j, Y_k^{(2)} = Y_i^{(2)} = Y_j^{(2)}\}$), we have $\mathbb{X}_i = (-1)^{\ell+1} \mathbb{X}_j$ given $G_K^{(1)}$ and $\mathcal{Y}^{(2)}$.

Similarly, let $\tilde{G}_K^{(1)}$ be the random partition of $\{-K + 1, \dots, 0\}$ induced by $\tilde{G}^{(1)}$. Then for all $a, b \in \mathbb{R}$,

$$aU_{0,n,K} + b\tilde{U}_{0,n,K} | G_K^{(1)}, \tilde{G}_K^{(1)} \stackrel{d}{=} \frac{1}{a_{n_2}^{(2)}} \sum_{i=1}^n (aX_{(0,i),K}^* + b\tilde{X}_{(0,i),K}^*) | G_K^{(1)}, \tilde{G}_K^{(1)},$$

where

$$\{\bar{\mathbb{X}}_i\}_{i \in \mathbb{N}} \equiv \{aX_{(0,i),K}^* + b\tilde{X}_{(0,i),K}^*\}_{i \in \mathbb{N}},$$

given $G_K^{(1)}$ and $\tilde{G}_K^{(1)}$, this time is a generalized one-dimensional randomized Karlin model. Since $\{\bar{\mathbb{X}}_i\}_{i \in \mathbb{N}}$ has bounded and centered marginal distribution, we can thus apply Proposition 2.1 for generalized one-dimensional randomized Karlin model. The variance of the limit normal distribution is then

$$\mathbb{E}(\bar{\mathbb{X}}_1^2 | G_K^{(1)}, \tilde{G}_K^{(1)}) = a^2 \text{Var}(X_{0,K}^* | G_K^{(1)}) + b^2 \text{Var}(\tilde{X}_{0,K}^* | \tilde{G}_K^{(1)}) = a^2 \sigma_K + b^2 \tilde{\sigma}_K.$$

It follows that, by the Cramér–Wold device,

$$(U_{0,n,K}, \tilde{U}_{0,n,K}) \Rightarrow (\sigma_K Z, \tilde{\sigma}_K \tilde{Z}),$$

where Z and \tilde{Z} are standard normal random variables and the four random variables are independent. To conclude, we deduce that

$$\lim_{n \rightarrow \infty} \text{Cov}(U_{0,n,K}^2, \tilde{U}_{0,n,K}^2) = \text{Cov}(\sigma_K^2 Z^2, \tilde{\sigma}_K^2 \tilde{Z}^2) = 0,$$

which imply the desired result. □

Proof of tightness. Again, we proceed using Bickel–Wichura’s criterion [1]. Observe that for all $n \in \mathbb{N}$, $\{U_{j,n}\}_{j \in \mathbb{Z}}$ is a martingale-difference sequence with respect to $\{\mathcal{F}_j^{(1)}\}_{j \in \mathbb{Z}}$. By Burkholder’s inequality, for all $p \geq 1$, for all $\mathbf{n}, \mathbf{m} \in \mathbb{N}^2$,

$$\begin{aligned} \mathbb{E} \left(\frac{S_{\mathbf{m}}}{b_{n_1}^{(1)} a_{n_2}^{(2)}} \right)^{2p} &\leq C_p \left(\frac{a_{m_2}^{(2)}}{b_{n_1}^{(1)} a_{n_2}^{(2)}} \right)^{2p} \mathbb{E} \left(\sum_{j \in \mathbb{Z}} (b_{m_1}^{(1)})^2 U_{j,m_2}^2 \right)^p \\ &\leq C_p \left(\frac{b_{m_1}^{(1)}}{b_{n_1}^{(1)}} \right)^{2p} \left(\frac{a_{m_2}^{(2)}}{a_{n_2}^{(2)}} \right)^{2p} \mathbb{E} U_{0,n}^{2p}. \end{aligned}$$

Using that $\mathbb{E} U_{0,n}^{2p}$ is bounded uniformly in n and that

$$\left(\frac{b_{m_1}^{(1)}}{b_{n_1}^{(1)}} \right)^{2p} \left(\frac{a_{m_2}^{(2)}}{a_{n_2}^{(2)}} \right)^{2p} \sim \frac{m_1^{H_1} L_1^{-1}(m_1)}{n_1^{H_1} L_1^{-1}(n_1)} \frac{m_2^{H_2} L_2(m_2)}{n_2^{H_2} L_2(n_2)} \quad \text{as } \mathbf{n} \rightarrow \infty,$$

we can conclude as for the other models, dealing with the slowly varying functions by using Potter’s bound. □

Remark 4.4. As we have seen, the proof follows the same structure as for the two-dimensional Hammond–Sheffield model. In fact, our models have their natural generalizations to high dimensions ($d \geq 2$), and the proof will follow the same strategy. The generalization of the model to high dimensions, based on independent random partitions and assignment rules in different directions, is intuitively obvious. However, it is notationally heavy to introduce. We only briefly explain how the proof would go. If in all directions the random partition is the same as the one in the one-dimensional Karlin model, then the same proof as Theorem 2.2, by first conditioning on the partition, shall work. If at least in one direction, say the first, the random partition and assignment rule are the ones of the one-dimensional Hammond–Sheffield model, then the same strategy as in two-dimensional Hammond–Sheffield model and the combined model shall work, by first writing

$$\frac{1}{b_{\mathbf{n}}} S_{n_1, \dots, n_d} = \frac{1}{b_{n_1}^{(1)}} \sum_{j_1 \in \mathbb{Z}} b_{j_1, n_1}^{(1)} U_{j_1, n_2, \dots, n_d},$$

with $\{U_{j_1, n_2, \dots, n_d}\}_{j_1 \in \mathbb{Z}}$ a stationary sequence of martingale differences. The analysis of this martingale-difference sequence shall need results for generalized $(d - 1)$ -dimensional models (to be defined properly first). To complete the details of this strategy would require an induction argument.

Appendix: Conditional convergence

We follow the notations of Kallenberg [14], Chapter 5. Let $(\Omega, \mathcal{A}, \mathbb{P})$ be a probability space, (S, \mathcal{S}) be a Borel space and (T, \mathcal{T}) be a measurable space. Let ξ, η be two random elements in S, T respectively. A regular conditional distribution of ξ given η is defined as a random measure ν of the form

$$\nu(\eta, B) = \mathbb{P}(\xi \in B | \sigma(\eta)) \quad \text{almost surely, } B \in \mathcal{S},$$

where ν is a probability kernel from (T, \mathcal{T}) to (S, \mathcal{S}) : $\nu(\cdot, B)$ is \mathcal{T} -measurable for all $B \in \mathcal{S}$, and $\nu(t, \cdot)$ is a probability measure on (S, \mathcal{S}) for all $t \in T$. Under the previous regularity assumption on the space (S, \mathcal{S}) and (T, \mathcal{T}) , such a probability kernel ν exists, and is unique almost everywhere $\mathbb{P} \circ \eta^{-1}$ [14], Theorem 5.3. Furthermore, for all measurable function f on $(S \times T, \mathcal{S} \times \mathcal{T})$, with $\mathbb{E}|f(\xi, \eta)| < \infty$,

$$\mathbb{E}(f(\xi, \eta) | \sigma(\eta)) = \int \nu(\eta, ds) f(s, \eta) \quad \text{almost surely.}$$

See for example [14], Theorem 5.4.

Some of our results are in the form of conditional (functional) limit theorems for the random field given underlying the random partition. The random partition, denoted by η here, and the random field $\{X_i\}_{i \in \mathbb{Z}^d}$, $d \in \mathbb{N}$ are defined on a common probability space $(\Omega, \mathcal{A}, \mathbb{P})$. Let $\{Z_n\}_{n \in \mathbb{N}}$ be a sequence of real-valued random variables (the normalized partial sum with appropriate normalization) in the same probability space. Then, let $\nu_n(\eta, \cdot)$ denote the regular conditional distribution of Z_n given η . With $\mathcal{G} = \sigma(\eta)$, we write for some \mathcal{G} -measurable random variable V (possibly a constant),

$$Z_n | \mathcal{G} \Rightarrow V \cdot \mathcal{N}(0, 1),$$

if $\nu_n(\eta(\omega), \cdot)$ as $n \rightarrow \infty$ converges to the standard normal distribution multiplied by $V(\omega)$ almost surely. That is, for all bounded continuous functions $h : \mathbb{R} \rightarrow \mathbb{R}$,

$$\lim_{n \rightarrow \infty} \int h(z) \nu_n(\eta(\omega), dz) = \int h(z) \frac{1}{\sqrt{2\pi}} e^{-z^2/2} dz \cdot V(\omega) \quad \text{almost surely.}$$

In this case we say that the conditional central limit theorem holds.

The conditional functional central limit theorem is interpreted in a similar way. Let $\mathcal{Z} = \{\mathcal{Z}(t)\}_{t \in T}$ and $\mathcal{Z}_n = \{Z_n(t)\}_{t \in T}$, $n \in \mathbb{N}$ with $T = [0, 1]^d$, $d \in \mathbb{N}$, be real-valued stochastic processes in $D(T)$ equipped with the Skorohod topology, defined in the same probability space. We write

$$\{\mathcal{Z}_n(t)\}_{t \in T} | \mathcal{G} \Rightarrow V \cdot \{\mathcal{Z}(t)\}_{t \in T},$$

if, letting $\nu_n(\eta, \cdot)$ denote this time the regular conditional distribution of \mathcal{Z}_n given η and $\mu_{\mathcal{Z}}$ denote the distribution of \mathcal{Z} , both as probability measures on $D(T)$, for all bounded and continuous function h from $D(T)$ to \mathbb{R} ,

$$\lim_{n \rightarrow \infty} \int_{D(T)} h(\zeta) \nu_n(\eta(\omega), d\zeta) = \int_{D(T)} h(\zeta) \mu_{\mathcal{Z}}(d\zeta) \cdot V(\omega) \quad \text{almost surely.}$$

Acknowledgements

The authors would like to thank two anonymous referees for their careful reading and helpful comments. YW's research was partially supported by the NSA grants H98230-14-1-0318 and H98230-16-1-0322, the ARO grant W911NF-17-1-0006, and Charles Phelps Taft Research Center at University of Cincinnati.

References

- [1] Bickel, P.J. and Wichura, M.J. (1971). Convergence criteria for multiparameter stochastic processes and some applications. *Ann. Math. Stat.* **42** 1656–1670. [MR0383482](#)
- [2] Biermé, H. and Durieu, O. (2014). Invariance principles for self-similar set-indexed random fields. *Trans. Amer. Math. Soc.* **366** 5963–5989. [MR3256190](#)
- [3] Biermé, H., Durieu, O. and Wang, Y. (2017). Invariance principles for operator-scaling Gaussian random fields. *Ann. Appl. Probab.* **27** 1190–1234. [MR3655864](#)
- [4] Biermé, H., Meerschaert, M.M. and Scheffler, H.-P. (2007). Operator scaling stable random fields. *Stochastic Process. Appl.* **117** 312–332. [MR2290879](#)
- [5] Billingsley, P. (1999). *Convergence of Probability Measures*, 2nd ed. New York: Wiley. [MR1700749](#)
- [6] Bingham, N.H., Goldie, C.M. and Teugels, J.L. (1987). *Regular Variation. Encyclopedia of Mathematics and Its Applications* **27**. Cambridge: Cambridge Univ. Press. [MR0898871](#)
- [7] Bolthausen, E. (1982). On the central limit theorem for stationary mixing random fields. *Ann. Probab.* **10** 1047–1050. [MR0672305](#)
- [8] Dedecker, J. (2001). Exponential inequalities and functional central limit theorems for a random fields. *ESAIM Probab. Stat.* **5** 77–104. [MR1875665](#)
- [9] Durieu, O. and Wang, Y. (2016). From infinite urn schemes to decompositions of self-similar Gaussian processes. *Electron. J. Probab.* **21** Paper No. 43, 23. [MR3530320](#)
- [10] Enriquez, N. (2004). A simple construction of the fractional Brownian motion. *Stochastic Process. Appl.* **109** 203–223. [MR2031768](#)
- [11] Gnedin, A., Hansen, B. and Pitman, J. (2007). Notes on the occupancy problem with infinitely many boxes: General asymptotics and power laws. *Probab. Surv.* **4** 146–171. [MR2318403](#)
- [12] Hammond, A. and Sheffield, S. (2013). Power law Pólya's urn and fractional Brownian motion. *Probab. Theory Related Fields* **157** 691–719. [MR3129801](#)
- [13] Hu, Y., Øksendal, B. and Zhang, T. (2000). Stochastic partial differential equations driven by multi-parameter fractional white noise. In *Stochastic Processes, Physics and Geometry: New Interplays, II (Leipzig, 1999)*. *CMS Conf. Proc.* **29** 327–337. Providence, RI: Amer. Math. Soc. [MR1803426](#)
- [14] Kallenberg, O. (1997). *Foundations of Modern Probability*. New York: Springer. [MR1464694](#)
- [15] Kamont, A. (1996). On the fractional anisotropic Wiener field. *Probab. Math. Statist.* **16** 85–98. [MR1407935](#)

- [16] Karlin, S. (1967). Central limit theorems for certain infinite urn schemes. *J. Math. Mech.* **17** 373–401. [MR0216548](#)
- [17] Klüppelberg, C. and Kühn, C. (2004). Fractional Brownian motion as a weak limit of Poisson shot noise processes – With applications to finance. *Stochastic Process. Appl.* **113** 333–351. [MR2087964](#)
- [18] Kolmogoroff, A.N. (1940). Wienersche Spiralen und einige andere interessante Kurven im Hilbertschen Raum. *Dokl. Akad. Nauk SSSR* **26** 115–118. [MR0003441](#)
- [19] Lavancier, F. (2007). Invariance principles for non-isotropic long memory random fields. *Stat. Inference Stoch. Process.* **10** 255–282. [MR2321311](#)
- [20] Lei, P. and Nualart, D. (2009). A decomposition of the bifractional Brownian motion and some applications. *Statist. Probab. Lett.* **79** 619–624. [MR2499385](#)
- [21] Mandelbrot, B.B. and Van Ness, J.W. (1968). Fractional Brownian motions, fractional noises and applications. *SIAM Rev.* **10** 422–437. [MR0242239](#)
- [22] McLeish, D.L. (1974). Dependent central limit theorems and invariance principles. *Ann. Probab.* **2** 620–628. [MR0358933](#)
- [23] Mikosch, T. and Samorodnitsky, G. (2007). Scaling limits for cumulative input processes. *Math. Oper. Res.* **32** 890–918. [MR2363203](#)
- [24] Øksendal, B. and Zhang, T. (2001). Multiparameter fractional Brownian motion and quasi-linear stochastic partial differential equations. *Stoch. Stoch. Rep.* **71** 141–163. [MR1922562](#)
- [25] Peligrad, M. and Sethuraman, S. (2008). On fractional Brownian motion limits in one dimensional nearest-neighbor symmetric simple exclusion. *ALEA Lat. Am. J. Probab. Math. Stat.* **4** 245–255. [MR2448774](#)
- [26] Pipiras, V. and Taqqu, M.S. (2017). *Long-Range Dependence and Self-Similarity*. *Cambridge Series in Statistical and Probabilistic Mathematics* **45**. Cambridge: Cambridge Univ. Press. [MR3729426](#)
- [27] Pitman, J. (2006). *Combinatorial Stochastic Processes*. *Lecture Notes in Math.* **1875**. Berlin: Springer. [MR2245368](#)
- [28] Puplinskaitė, D. and Surgailis, D. (2015). Scaling transition for long-range dependent Gaussian random fields. *Stochastic Process. Appl.* **125** 2256–2271. [MR3322863](#)
- [29] Puplinskaitė, D. and Surgailis, D. (2016). Aggregation of autoregressive random fields and anisotropic long-range dependence. *Bernoulli* **22** 2401–2441. [MR3498033](#)
- [30] Samorodnitsky, G. (2016). *Stochastic Processes and Long Range Dependence*. Cham: Springer. [MR3561100](#)
- [31] Shen, Y. and Wang, Y. (2017). Operator-scaling Gaussian random fields via aggregation. Preprint. Available at <https://arxiv.org/abs/1712.07082>.
- [32] Wang, Y. (2014). An invariance principle for fractional Brownian sheets. *J. Theoret. Probab.* **27** 1124–1139. [MR3278934](#)
- [33] Xiao, Y. (2009). Sample path properties of anisotropic Gaussian random fields. In *A Minicourse on Stochastic Partial Differential Equations*. *Lecture Notes in Math.* **1962** 145–212. Berlin: Springer. [MR2508776](#)

Received September 2017 and revised January 2018