

# Some properties of the Biggs-Smith geometry

Hendrik Van Maldeghem

Valerie Ver Gucht

## Abstract

We settle two problems posed by the first author in [2] concerning the Biggs-Smith geometry. The first is to determine the generating rank, the second is to prove that this geometry possesses exactly eighteen ovoids.

## 1 Introduction

The Biggs-Smith geometry is a bislim point-line geometry arising from the Biggs-Smith graph (for precise definitions, see below). It is introduced in [2] as one out of a family of ten highly symmetric bislim geometries with nice properties, and connected with interesting small groups — usually (almost) simple ones. In the general study performed in [2], there were two questions related to the Biggs-Smith geometry that remained unsolved in [2]. The first one is the determination of the generating rank. It was shown in [2] that this rank was either 19 or 20. The conjecture was the first value, since the universal embedding rank is also 19. In the present paper, we show this conjecture to be true. This reinforces the general conjecture that these ranks are the same whenever the geometries have a rich enough automorphism group — but no-one knows how rich it precisely has to be, and no clue for a proof has been found yet, as far as we know.

The second problem concerns ovoids. In [2], eighteen ovoids were constructed — to which we refer as *classical* here — and these were used to construct a real embedding of the Biggs-Smith geometry. The existence of another class of ovoids would probably give rise to a different embedding. This would be quite interesting since such a situation would be new. In the present paper, however, we show that there are no non-classical ovoids. This negative result is, in our opinion, worth recording, also for its proof, which makes use of the interplay between two constructions of the Biggs-Smith graph: the original one as defined by Biggs and Smith [1], and the recent one by the first author [2].

## 2 Definitions

A point-line geometry is a system  $\Gamma = (\mathcal{P}, \mathcal{L}, \mathbf{I})$  consisting of a point set  $\mathcal{P}$ , a line set  $\mathcal{L}$  and a symmetric incidence relation  $\mathbf{I}$  between  $\mathcal{P}$  and  $\mathcal{L}$  expressing precisely when a point is incident with a line. Usually we think of a line as the set of points incident with it and we accordingly use phrases like “a point is on a line”, “a line goes through a point”, etc. If all lines of  $\Gamma$  carry the same number  $s + 1$  of points and all points are incident with the same number  $t + 1$  of lines, then we say that  $\Gamma$  has *order*  $(s, t)$ . If  $s = 2$ , then we call  $\Gamma$  *slim*. If also  $t = 2$ , then we call  $\Gamma$  *bislim*.

An *isomorphism* (*monomorphism*) from the point-line geometry  $\Gamma$  to the point-line geometry  $\Gamma'$  is a bijection (injective map) from the point set of  $\Gamma$  to the point set of  $\Gamma'$  together with a bijection (injective map) from the line set of  $\Gamma$  to the line set of  $\Gamma'$  such that two elements of  $\Gamma$  are incident precisely when their images in  $\Gamma'$  are incident.

Let  $\Gamma = (\mathcal{P}, \mathcal{L}, \mathbf{I})$  be a point-line geometry and  $\mathbf{PG}(d, \mathbb{K})$  the  $d$ -dimensional projective space over the field  $\mathbb{K}$ . An *embedding of  $\Gamma$  in  $\mathbf{PG}(d, \mathbb{K})$*  is a monomorphism of  $\Gamma$  into the point-line geometry of  $\mathbf{PG}(d, \mathbb{K})$  such that the images of all points span  $\mathbf{PG}(d, \mathbb{K})$ . Usually, one identifies a point with its image in  $\mathbf{PG}(d, \mathbb{K})$ .

Let  $\Gamma$  be a slim geometry. If the positive integer  $d$  is maximal with respect to the property that  $\Gamma$  admits an embedding in  $\mathbf{PG}(d, 2)$ , then  $d + 1$  is called the *universal embedding rank of  $\Gamma$* . If  $d > 0$  does not exist, then we say that the universal embedding rank is zero. Also, we say that a subset of points of  $\Gamma$  *generates*  $\Gamma$  if the smallest slim subgeometry of  $\Gamma$  containing these points and with the property that, whenever two points of that subgeometry are collinear in  $\Gamma$ , they are also collinear in the subgeometry, coincides with  $\Gamma$ . The *generating rank of  $\Gamma$*  is the minimal number of points needed to generate  $\Gamma$ . Obviously the generating rank of  $\Gamma$  cannot be smaller than the universal embedding rank of  $\Gamma$ .

An *ovoid* of a point-line geometry  $\Gamma$  is a set of points with the property that each line is incident with exactly one point of the ovoid.

## 3 The Biggs-Smith geometry

The bislim point-line geometry **BS** is the *neighborhood geometry* arising from the non-bipartite Biggs-Smith graph on 102 vertices (see figure 1). This means that the points of **BS** are the vertices of the graph and the lines are a second copy of the same vertex set with the incidence relation induced by adjacency. The original construction of the Biggs-Smith graph uses 17 copies  $H_i$ ,  $i \in \mathbf{GF}(17)$  of an “H”-graph and tells one very explicitly how to join vertices of each of these copies (see figure 2).

So the vertices are  $p_0, \dots, p_{16}$ ,  $r_0, \dots, r_{16}$ , etc. The notation  $17/4$  at the  $q$ -vertex of the “H”-graph means that we join  $q_i$  with  $q_{i+4}$ . Similarly for the others.

We now repeat the construction of the Biggs-Smith graph and the geometry **BS** as given in [2].

Consider the projective line  $\mathbf{PG}(1, 17)$  over the field  $\mathbf{GF}(17)$ . A *harmonic triplet* consists of three pairs of points  $\{a, b\}, \{c, d\}, \{e, f\}$ , with  $a, b, \dots, f$  all distinct, such that




Figure 1: Biggs-Smith graph




Figure 2: “H”-graph

we have the following equality of cross ratios:  $(a, b; c, d) = (a, b; e, f) = (c, d; e, f) = -1$ . Three mutually disjoint harmonic triplets will be called a *trisection*. The group  $\mathbf{PSL}_2(17)$  acts in two orbits of size 102 on the set of harmonic triplets. Let  $\mathcal{P}$  be one of these orbits. Then  $\mathcal{P}$  is the vertex set of the Biggs-Smith graph, and two vertices are adjacent if the corresponding harmonic triplets have a pair of points in common, e.g.  $\{\{\infty, 0\}, \{1, -1\}, \{4, -4\}\}$  is adjacent to  $\{\{\infty, 0\}, \{2, -2\}, \{8, -8\}\}$ . The point set of  $\mathbf{BS}$  is also  $\mathcal{P}$ , and the trisections within  $\mathcal{P}$  form the line set.

It is trivial to see that the harmonic triplets for which some pair contains a fixed point of  $\mathbf{PG}(1, 17)$  form an ovoid of  $\mathbf{BS}$ . Such an ovoid will be called *classical*. Since there are eighteen points in  $\mathbf{PG}(1, 17)$ , there are eighteen classical ovoids.

Our Main Result reads:

**Main Result** *The geometry  $\mathbf{BS}$  has generating rank equal to 19. Consequently, the generating rank is equal to the universal embedding rank. Also,  $\mathbf{BS}$  does not contain ovoids different from the classical ones.*

## 4 Generating rank

**Theorem 4.1.** *The generating rank of  $\mathbf{BS}$  is equal to 19.*

*Proof.* Since the universal embedding rank of  $\mathbf{BS}$  is equal to 19, the generating rank is at least 19. We consider the construction of the Biggs-Smith graph with the “H”-graph.

We assert that  $\mathcal{G} := \{r_i, p_0, p_1, p_2 \mid i \in \mathbf{GF}(17) \setminus \{-5\}\}$  is a generating set for  $\mathbf{BS}$ . It suffices to prove that  $r_{-5}$  belongs to the slim subgeometry  $\mathcal{S}$  of  $\mathbf{BS}$  generated by the points of  $\mathcal{G}$ . Then this subgeometry coincides with  $\mathbf{BS}$ , as it is easy to see (and which, in fact, will also follow from our proof).

A vertex  $r_i$ ,  $i \in \mathbf{GF}(17)$  is adjacent to two other vertices  $r_{i-2}$  and  $r_{i+2}$  and one vertex  $y_i$ . Hence, the vertices  $y_i$  with  $i \in \mathbf{GF}(17) \setminus \{-3, -7\}$  belong to  $\mathcal{S}$ .

Each vertex  $p_i$ ,  $i \in \mathbf{GF}(17)$  is adjacent with two vertices  $p_{i+1}$  and  $p_{i-1}$  and with one vertex  $x_i$ . The latter has next to  $p_i$  still two other neighbors:  $q_i$  and  $y_i$ . Consequently, for  $i = 0, 1, 2$ , the point  $q_i$  belongs to  $\mathcal{S}$ . Also, we have that  $x_1$  belongs to the subgeometry  $\mathcal{S}$ .

A vertex  $s_i$ ,  $i \in \mathbf{GF}(17)$  is adjacent to two other vertices  $s_{i+8}$  and  $s_{i-8}$  and with one vertex  $y_i$ . The latter has two more neighbors:  $r_i$  and  $x_i$ . It's easy to see that the point  $s_1$  is a point of  $\mathcal{S}$ .

The vertex  $s_{-8}$  is adjacent to the vertices  $s_1$  and  $y_{-8}$ , representing points which belong to  $\mathcal{S}$ . Hence,  $s_0$  belongs to  $\mathcal{S}$ . Inductively, by reducing each value of  $i$  in the previous case by 1, we conclude that the vertices  $s_i$  with  $i \in \mathbf{GF}(17) \setminus \{1, 2, 3, 4, 5\}$  belong to the subgeometry  $\mathcal{S}$ .

For  $i$  different from  $-5, 1, 2, 3, 4, 5$ , the vertices  $s_i$  and  $r_i$  are in  $\mathcal{S}$ . Hence, the twelve vertices  $x_i$  with  $i \in \mathbf{GF}(17) \setminus \{-5, 2, 3, 4, 5\}$  belong to  $\mathcal{S}$ . (Observe that for  $i = 1$  we already have shown that  $x_1$  belongs to  $\mathcal{S}$ .)

Given a vertex  $q_i$ , this vertex is adjacent to two other vertices  $q_{i+4}$  and  $q_{i-4}$ . Also,  $q_i$  is adjacent to the vertex  $x_i$ . Hence,  $q_i \in \mathcal{S}$  for  $i = -8, -7, -6$ .

The vertex  $p_i$  is adjacent to two vertices  $p_{i-1}$  and  $p_{i+1}$  and to  $x_i$ . By starting with the vertex with  $i$ -coordinate equal to 0 and letting  $i$  successively diminish by one, we get that  $p_i \in \mathcal{S}$  for  $i = -1, -2, -3, -4, -5$ . Because of the fact that  $p_{-6}$  is adjacent to  $p_{-5}$  and to  $x_{-6}$  we have  $p_{-7} \in \mathcal{S}$ .

The vertex  $x_{-7}$  is adjacent to  $p_{-7}$  and to  $q_{-7}$ , hence  $y_{-7}$  belongs to  $\mathcal{S}$ .

The vertex  $r_{-7}$  is adjacent to  $y_{-7}$  and to  $r_8$ , hence the vertex  $r_{-5} \in \mathcal{S}$ . Since this vertex is the missing vertex, the generating rank of  $\mathbf{BS}$  is equal to 19. ■

## 5 Ovoids

**Theorem 5.1.** *The point-line geometry **BS** has 18 ovoids, each containing 34 points.*

*Proof.* Let  $\mathcal{O}$  be any ovoid of **BS**. We will show that  $\mathcal{O}$  is classical. Consider any point  $v$  of  $\mathcal{O}$ . Then, by considering  $v$  as a vertex of the Biggs-Smith graph, and then interpreting  $v$  as a line of **BS**, we see that  $v$  is adjacent to exactly one other vertex  $w$  that, viewed as a point, belongs to  $\mathcal{O}$ . Now, the harmonic triplets corresponding to  $v$  and  $w$  have a pair of points of  $\mathbf{PG}(1, 17)$  in common, and so they are contained in precisely two classical ovoids. Hence, in order to prove the theorem, it suffices to show that two adjacent vertices  $v$  and  $w$  are contained in precisely two ovoids of **BS**. By the transitivity of the automorphism group on pairs of adjacent vertices, we may take  $v = p_0$  and  $w = x_0$ .

We identify the points of  $\mathcal{O}$  with the corresponding vertices of the Biggs-Smith graph.

So suppose  $p_0, x_0 \in \mathcal{O}$ . It's easy to see that each copy of the "H"-graph contains exactly two vertices of  $\mathcal{O}$ : one vertex corresponding to  $\{p, q, y\}$  together with one vertex corresponding to  $\{r, s, x\}$ . Hence,  $r_0, y_0, q_0, s_0$  are not in  $\mathcal{O}$ . Also, the vertices  $x_1, p_2, x_{-1}, p_{-2}, p_1, p_{-1}, q_4, q_{-4}$  do not belong to  $\mathcal{O}$ . Next, we distinguish two cases based on  $H_1$ : either  $y_1 \in \mathcal{O}$  or  $q_1 \in \mathcal{O}$ .

If  $y_1$  belongs to  $\mathcal{O}$  then  $q_1$  does not. The vertices  $r_3, r_{-1}, s_{-8}$  and  $s_{-7}$  are not in  $\mathcal{O}$ . Since  $y_0$  and  $s_{-8}$  are not in  $\mathcal{O}$  it follows that  $s_8$  is an element of the ovoid. Consequently,  $r_8, x_8, s_7, y_{-1}$  are not in  $\mathcal{O}$ . The two vertices of  $H_{-1}$  belonging to the ovoid are  $q_{-1}$  and  $s_{-1}$ . Hence,  $x_3, q_7, x_{-5}, q_8, s_{-2}, y_7, y_8$  are not in  $\mathcal{O}$ . Since  $q_0$  and  $q_8$  do not belong to  $\mathcal{O}$ , we have that  $x_4$  is an element of  $\mathcal{O}$  which implies that  $r_4, s_4, p_3$  and  $p_5$  are not in  $\mathcal{O}$ . As  $x_3$  and  $p_2$  are not in the ovoid, the vertex  $p_4$  is element of  $\mathcal{O}$ . The vertices  $y_4, x_5$  and  $p_6$  are not in  $\mathcal{O}$ . But the vertex  $s_3$  is element of the ovoid. This means that  $s_2, y_{-6}$  and  $y_{-5}$  are not contained in the ovoid. Since  $r_0$  and  $r_4$  do not belong to  $\mathcal{O}$  the vertex  $y_2$  is element of the ovoid. Thus,  $q_2, s_{-6}$  do not belong to  $\mathcal{O}$ . Also, since  $x_8$  and  $q_4$  are not in  $\mathcal{O}$ , the vertex  $q_{-5}$  is in  $\mathcal{O}$ . Therefore,  $p_{-5}$  and  $q_3$  are not elements of the ovoid. The second vertex of  $H_3$ , next to  $s_3$ , belonging to the ovoid, is  $y_3$ , implying that  $r_1, r_5$  and  $s_{-5}$  are not contained in  $\mathcal{O}$ . It is easy to see that  $r_{-5}$  belongs to  $\mathcal{O}$ . Also,  $x_2$  is element of the ovoid since  $p_1$  and  $p_3$  are not. Hence,  $y_{-3}, y_{-7}, r_2, q_{-2}$  and  $q_6$  are not vertices of  $\mathcal{O}$ . We deduce that  $y_{-2}$  is in  $\mathcal{O}$  and consequently,  $s_6$  and  $r_{-4}$  are not in the ovoid. Since  $y_4$  and  $s_{-5}$  are not in  $\mathcal{O}$  the vertex  $s_{-4}$  is an element of the ovoid. Thus that  $x_{-4}, y_5$  and  $s_{-3}$  are not contained in the ovoid. Since  $x_{-4}$  and  $p_{-5}$  are not in  $\mathcal{O}$  the vertex  $p_{-3}$  is an element of the ovoid. It is also easy to see that  $q_5$  and  $s_5$  are contained in  $\mathcal{O}$ . It follows that  $q_{-3}, x_{-2}, x_{-8}$  and  $y_{-4}$  are not contained in the ovoid. Since  $x_{-2}$  and  $s_{-2}$  are not in  $\mathcal{O}$  it follows that  $r_{-2}$  belongs to  $\mathcal{O}$ . The vertices  $y_{-4}$  and  $q_{-4}$  are not contained in the ovoid, hence  $p_{-4}$  is a vertex in  $\mathcal{O}$ . So,  $\{r_{-6}, x_{-3}, p_{-6}\}$  is not a subset of  $\mathcal{O}$ . It is clear that  $r_{-3}$  is a vertex of  $\mathcal{O}$ , but  $r_{-7}$  is not. We already found four elements of  $H_{-6}$  not in  $\mathcal{O}$ , so the two remaining vertices,  $x_{-6}$  and  $q_{-6}$  do belong to the ovoid. Thus,  $x_7$  and  $p_{-7}$  are not in  $\mathcal{O}$ . An analogous reasoning holds for  $H_{-7}$ :  $x_{-7}$  and  $q_{-7}$  belong to  $\mathcal{O}$ , but  $x_6$  and  $p_{-8}$  do not. In the same way we deduce that  $r_{-8}, r_6$  and  $y_6$  are in  $\mathcal{O}$ . Furthermore,  $p_7$  and  $r_7$  are vertices contained in the ovoid, implying that  $y_{-8}$  is not in  $\mathcal{O}$  and hence  $q_{-8}$  is element of  $\mathcal{O}$ . Since  $y_8$  and  $q_8$  are both not in  $\mathcal{O}$ , it follows that  $p_8$  is a vertex of  $\mathcal{O}$ . The copy  $H_1$  has already one vertex contained in  $\mathcal{O}$  and four not contained in  $\mathcal{O}$ ,

thus the remaining vertex  $s_1$  is an element of  $\mathcal{O}$ . We now have examined all vertices of the Biggs-Smith graph and may conclude that the ovoid  $\mathcal{O}$  which contains the vertices  $p_0$ ,  $x_0$  and  $y_1$  is unique.

Now let us consider the second case where besides  $p_0$  and  $x_0$ , also  $q_1$  is an element of  $\mathcal{O}$ . The 4-arc transitivity of the Biggs-Smith graph implies that there is an automorphism taking the 4-arc  $(x_0, p_0, p_1, x_1, y_1)$  onto the 4-arc  $(x_0, p_0, p_1, x_1, q_1)$ . Hence there is a unique ovoid containing the vertices  $x_0$ ,  $p_0$  and  $q_1$ .

The theorem is proved. ■

## References

- [1] N. L. Biggs and D. H. Smith, On trivalent graphs, *Bull. Lond. Math. Soc.* **3** (1971), 127 – 131.
- [2] H. Van Maldeghem, Ten exceptional geometries from trivalent distance regular graphs, *Ann. Combin.* **6** (2002), 209 – 228.

Hendrik Van Maldeghem  
Department of Pure Mathematics and Computer Algebra  
Ghent University  
Galglaan 2, 9000 Gent  
BELGIUM  
hvm@cage.UGent.be

Valerie Ver Gucht  
Department of Applied Mathematics, Biometrics and Process Control  
Ghent University  
Coupure Links 653, 9000 Gent  
BELGIUM  
Valerie.VerGucht@biomath.UGent.be