

Affine Kac-Moody Algebras and Semi-Infinite Flag Manifolds

Boris L. Feigin¹ and Edward V. Frenkel^{2, *}

¹ 117218, Profsoyuznaya, 13/12, apt 91, Moscow, USSR

² 140410, Suvorova, 100, apt 27, Kolomna, Moscow district, USSR

Dedicated to Dmitry Borisovich Fuchs on his 50th birthday

Abstract. We study representations of affine Kac-Moody algebras from a geometric point of view. It is shown that Wakimoto modules introduced in [18], which are important in conformal field theory, correspond to certain sheaves on a semi-infinite flag manifold with support on its Schubert cells. This manifold is equipped with a remarkable semi-infinite structure, which is discussed; in particular, the semi-infinite homology of this manifold is computed. The Cousin-Grothendieck resolution of an invertible sheaf on a semi-infinite flag manifold gives a two-sided resolution of an irreducible representation of an affine algebra, consisting of Wakimoto modules. This is just the BRST complex. As a byproduct we compute the homology of an algebra of currents on the real line with values in a nilpotent Lie algebra.

1. Introduction

In [18, 19] we have introduced and studied a new class of representations of affine Kac-Moody algebras, the so-called *Wakimoto modules* [44]. These representations allow *bosonic realization*, the Sugawara energy-momentum tensor being quadratic in bosons. This gives a new bosonization rule for the Wess-Zumino-Witten (WZW) models. In [19] we explicitly constructed the intertwining operators between Wakimoto modules and chains (or primary fields) which are submodules of their homomorphisms, using vertex operators. Our results enable us to give an integral representation of the correlation functions in WZW models on the plane in spirit of [14] (it was done soon after [15, 27]). In [20] we have proposed the *two-sided Bernstein-Gelfand-Gelfand (BGG) resolution*, or BRST complex, of an irreducible representation of an affine Kac-Moody algebra, consisting of Wakimoto modules (recall that the usual BGG resolution [7, 26, 41] is one-sided and consists of Verma modules). According to Felder's work [23] (where a similar resolution is constructed over the Virasoro algebra), this

* Address after September 15, 1989: Mathematics Department, Harvard University, Cambridge, MA 02138, USA

enables us to compute the correlation functions in the WZW models on the torus, “extracting” an irreducible representation from Wakimoto modules with the use of our resolution.

Wakimoto modules also play an important role in the study of the functorial correspondence between affine Kac-Moody algebras and the Virasoro algebra (or W -algebras [20, 47]) and of the highest weight modules with central charge $-$ (dual Coxeter number) [19], Appendix B.

Thus we see that Wakimoto modules are very useful both in representation theory and in conformal field theory.

In this work we show that Wakimoto modules naturally appear in the representation theory of affine Kac-Moody algebras: *Verma modules, contragredient Verma modules and Wakimoto modules are the particular cases of the general construction*. Moreover, modules with similar properties emerge in the case of finite-dimensional simple Lie algebras. Analogous construction seems to exist for arbitrary Kac-Moody algebras.

The crucial point in our investigation is the correspondence between the highest weight representations of the Lie algebra and the sheaves of \mathcal{D} -modules (or constructible sheaves) on its flag manifold, *lisse* with respect to Schubert stratification [1, 2, 4, 10, 34].

As well-known, the contragredient Verma module with integral highest weight over a finite-dimensional Lie algebra corresponds to the constant sheaf supported on the big Schubert cell and to the sheaf of local cohomology of the appropriate invertible sheaf on a flag manifold with support on this cell. If we take the local cohomology of the invertible sheaf, with support on another cell, then we obtain the module, which we call the twisted Verma module. Twisted Verma modules with equal highest weights are labelled by elements of the Weyl group (in particular, the unit corresponds to the contragredient Verma module, and the element of maximal length $-$ to the Verma module). Their composition series quotients coincide but the composition series themselves are different.

The Cousin-Grothendieck resolution of a dominant invertible sheaf with respect to Schubert stratification of a flag manifold appears to be the contragredient BGG resolution [35], the Cousin-Grothendieck resolution of a twisted invertible sheaf gives the twisted BGG resolution, which consists of twisted Verma modules. In particular, the element of maximal length of the Weyl group corresponds to the usual BGG resolution.

The infinite-dimensional affine case is more interesting. Here again we have twisted Verma modules, but additional opportunities appear.

The affine Weyl group is infinite and there are no elements of maximal length. But let us consider a “limit element” of “semi-infinite length” of this group. The highest weight module corresponding to is a Wakimoto module. Thus, Wakimoto modules are intermediate between contragredient Verma modules and Verma modules (which correspond to the element of “infinite length”).

The sheaves, corresponding to Wakimoto modules, “live” on the limit *semi-infinite flag manifold*, which is the coset space of the affine Kac-Moody group by an appropriate subgroup. They are *lisse* with respect to the Schubert stratification. The usual flag manifold of an affine Kac-Moody algebra allows two stratifications: by cells of finite dimension or by cells of finite codimension

[40, 39]. The cells of the semi-infinite flag manifold are both of infinite dimension and of infinite codimension.

The semi-infinite flag manifold is endowed with a rather intriguing “semi-infinite structure.” For example, its Schubert cells represent the semi-infinite homology classes of this manifold. It is interesting that this homology may also be obtained using Floer’s theory [24]. There is a remarkable Morse function of Conley-Zehnder type [11] on this manifold, whose singular points are of semi-infinite indices and give the same semi-infinite homology. We believe that there exists a theory of semi-infinite manifolds, including semi-infinite sheaves and their cohomologies. Our flag manifolds seem to be the first examples of this theory.

A Wakimoto module corresponds to a constant sheaf, supported on a Schubert cell of the semi-infinite flag manifold and to a sheaf of local “semi-infinite cohomology” of the invertible sheaf with support on this cell. Cousin-Grothendieck resolution is a two-sided BGG resolution (a similar resolution may be obtained in a different way).

The Two-sided BGG resolution gives the semi-infinite analogue of Borel-Weil-Bott theorem. As a corollary we obtain an unexpected result about usual homology of the Lie algebras of currents on the real line with values in nilpotent subalgebras of the simple Lie algebra.

The paper is arranged as follows.

In Sect. 2 we treat the finite-dimensional counterpart of our constructions. This section illustrates the main ideas, which we apply later. In Sect. 3 we give the definition of Wakimoto modules over affine Kac-Moody algebras and account for their place in the representation theory of these algebras. Section 4 is devoted to semi-infinite flag manifolds. It clarifies the relations of Wakimoto modules with the geometry of these manifolds. In Sect. 5 we construct Wakimoto modules overcoming certain homological problems. In Sect. 6 we establish and prove two-sided BGG resolution and use it for computation of (co)homology. The concluding Sect. 7 is devoted to some examples and applications.

Appendix A contains the definition of semi-infinite (co)homology [16], used in this work. In Appendix B we sketch our results and conjectures [19] about the structure of highest weight modules with central charge $-$ (dual Coxeter number).

2. The Finite-Dimensional Case

2.1. Notations and Preliminaries [29, 5, 6]

Let G be a complex simple Lie group of rank n , \mathfrak{g} is its Lie algebra, $\mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$ is the Cartan decomposition of \mathfrak{g} , $\Delta = \Delta(G)$ is the root system of G , $\Lambda = \Lambda_G$ is the root lattice of G . $\Delta = \Delta_+ \cup \Delta_-$, where Δ_+ (Δ_-) is the set of positive (negative) roots. Denote by $\alpha_1, \dots, \alpha_n$ the set of simple roots, E_i, H_i, F_i – Cartan generators of \mathfrak{g} . Let (\cdot, \cdot) be the inner product in \mathfrak{h}^* , $S = S(G)$ is the Weyl group of G , w_0 is its maximal element.

Let $F = G/B$ be flag manifold of G , where B is the Borel subgroup of G , corresponding to the Lie algebra $\mathfrak{b} = \mathfrak{h} \oplus \mathfrak{n}_+$. As is well-known, F decomposes into Schubert cells F_s , where s runs the Weyl group S , which are the orbits of the

nilpotent group N_+ , corresponding to the Lie algebra n_+ . Namely, $F_s = N_s \cdot \bar{s}$, where \bar{s} is the image of $s \in G$ under the projection $G \rightarrow F$. The cell F_s is isomorphic to $sN_+s^{-1} \cap N_+$, $\text{codim } F_s = l(s)$ is the length of the element s of the Weyl group, defined as follows: $l(s) = \#\{\alpha \in \Delta_+ : s^{-1}\alpha \in \Delta_-\}$ (here and further $\#$ denotes the number of elements of the set).

The category \mathcal{O} of highest weight representations (in this paper highest weight representation means *any* representation from the category \mathcal{O}) of \mathfrak{g} decomposes into the subcategories \mathcal{O}_θ of representations, where the center of $\mathcal{U}(\mathfrak{g})$ acts by the character θ . If λ is the dominant integral weight, and $\theta(\lambda)$ is the corresponding central character, then $\mathcal{O}_{\theta(\lambda)}$ consists of the modules with highest weights of the form $s * \lambda = s(\lambda + \varrho) - \varrho$, $s \in S$, and ϱ is a half of the sum of positive roots of \mathfrak{g} .

2.2. Twisted Verma Modules

Let $\mu \in \mathfrak{h}^*$ and \mathbb{C}_μ be one-dimensional representation of $b_+ = \mathfrak{h} \oplus n_+$, determined as composition $b_+ \rightarrow \mathfrak{h} \xrightarrow{\mu} \mathbb{C}$. Denote by M_μ the Verma module over \mathfrak{g} with highest weight μ : $M_\mu = \mathcal{U}(\mathfrak{g}) \otimes_{\mathcal{U}(b_+)} \mathbb{C}_\mu$. Verma modules are characterized by the following properties:

- a) M_μ lies in the category \mathcal{O} ,
- b) $H_0(n_-, M_\mu) \simeq \mathbb{C}_\mu$ (as \mathfrak{h} -module), $H_i(n_-, M_\mu) = 0, i \neq 0$.

Let M_μ^* be the contragradient Verma module. It is characterized by the property a) and by the property

- b*) $H^0(n_+, M_\mu^*) \simeq \mathbb{C}_\mu$ (as \mathfrak{h} -module), $H^i(n_+, M_\mu^*) = 0, i \neq 0$.

Our aim is to introduce highest weight modules M_μ^w over \mathfrak{g} , where $w \in S$ which are intermediate between M_μ and M_μ^* in the following sense. Put $n_+^w = wn_+w^{-1}$. M_μ^w is characterized by the property a) and by the property

- b_w) $H^{l(w)}(n_+^w, M_\mu^w) \simeq \mathbb{C}_{\mu - w(\varrho) + \varrho}$ (as \mathfrak{h} -module),

$$H^i(n_+^w, M_\mu^w) = 0, \quad i \neq l(w).$$

In particular, $M_\mu^1 \simeq M_\mu^*$, $M_\mu^{w_0} = M_\mu$.

We call M_μ^w *twisted Verma modules*¹. They are highest weight modules with highest weight μ . M_μ^w is free over $n_+^w \cap n_-$ and dual to M_μ^w is free over $n_+^w \cap n_+$. Their composition series quotients coincide with those of the Verma module M_μ , but they are “glued” in a different way. If M_μ is irreducible then $M_\mu \simeq M_\mu^w$, but if it is not so, then the composition structures of $M_\mu^{w_1}$ and $M_\mu^{w_2}$ differ from each other.

We will construct twisted Verma modules.

But first, let us recall the correspondence between the highest weight \mathfrak{g} -modules from $\mathcal{O}_{\theta(\lambda)}$ (λ is a dominant integral weight) and certain sheaves on the flag manifold.

Any integral weight $\nu = w * \lambda$ defines the linear holomorphic complex bundle ξ_ν on F and invertible sheaf ξ_ν^* of its holomorphic sections. Let \mathcal{D}_ν be the sheaf of differential operators on the sheaf ξ_ν^* . We denote by $\xi = \xi_0$ the structural sheaf.

¹ Using shifted cohomology $H_s^i(n_+^w, \cdot)$ of n_+^w with respect to the decomposition $n_+^w = (n_+^w \cap n_+) \oplus (n_+^w \cap n_-)$ (see Appendix A) we can rewrite b_w) as follows: $H_s^0(n_+^w, M_\mu^w) \simeq \mathbb{C}_\mu$ (as \mathfrak{h} -module), $H_s^i(n_+^w, M_\mu^w) = 0, i \neq 0$

The following functor U_λ^w defines the equivalence between the derived category of the highest weight g -modules $D\mathcal{O}_{\theta(\lambda)}$ and of the derived category D_λ^w of the N_+ -invariant holonomic $\mathcal{D}_{w*\lambda}$ -modules on F [of amplitude $\leq l(w)$] [1]

$$U_\lambda^w : \mathcal{U} \in D_\lambda^w \rightarrow \mathbb{R}\Gamma(F; \mathcal{U}) \in D\mathcal{O}_{\theta(\lambda)}.$$

There are also contravariant functors from $D\mathcal{O}_{\theta(\lambda)}$ to the derived category Con of the category Con of constructible sheaves on F , lisse with respect to Schubert stratification.

Let M be g -module from $\mathcal{O}_{\theta(\lambda)}$. Denote by ξ the structural sheaf on the manifold G/N_- . The sheaf $\xi \otimes M$ is equivariant and equipped with the left action of g and the right action of h . Consider the complex of sheaves $C_*(g, \xi \otimes M)$ on F , where $C_*(g, \xi \otimes M)$ denotes the standard homological complex of g . This is the complex of sheaves with constructible cohomology, equipped with the action of h . So, it decomposes into direct sum:

$$C_*(g, \xi \otimes M) = \bigoplus_{\mu \in h^*} C_*^\mu(g, \xi \otimes M),$$

where $C_*^\mu(g, \xi \otimes M)$ denotes μ -eigenspace of h . The subcomplexes of sheaves $C_*^\mu(g, \xi \otimes M)$ are acyclic if $\mu \neq w*\lambda$. The subcomplex $C_*^{w*\lambda}(g, \xi \otimes M)$ on G/N_- defines the complex of sheaves $\tilde{C}_*^{w*\lambda}(M)$ on F and the complex of sheaves with constructible cohomologies $\text{Hom}_\xi(\tilde{C}_*^{w*\lambda}(M), \tilde{\xi}_{w*\lambda}) = \mathcal{V}_M$ on F , $\mathcal{V}_M \in \text{Con}$. For the point $\kappa \in F$ denote by b_κ the Lie algebra of the stabilizer of κ and by n_κ its radical. All subalgebras b_κ are conjugated and we identify $b_\kappa/n_\kappa = h_\kappa$ with h . The stalk of \mathcal{V}_M at κ is isomorphic to $C_*^{w*\lambda}(n_\kappa, M)$. This defines the functor $V_\lambda^w : D\mathcal{O}_{\theta(\lambda)} \rightarrow \text{Con}$.

For a given $\mathcal{V} \in \text{Der Con}$ we can construct the $\mathcal{D}_{w*\lambda}$ -module using the functor of local cohomology:

$$Z_\lambda^w : \mathcal{V} \in \text{Con} \rightarrow \mathbb{R} \text{Hom}_\xi(\mathcal{V}, \tilde{\xi}_{w*\lambda}) \in D_\lambda^w.$$

Three functors, defined above, give the following commutative diagram:

$$\begin{array}{ccc} & D\mathcal{O}_{\theta(\lambda)} & \\ \swarrow V_\lambda^w & & \nwarrow U_\lambda^w \\ \text{Con} & \xrightarrow{Z_\lambda^w} & D_\lambda^w \end{array}$$

Let \mathbf{C}_s be the constant sheaf with support on Schubert cell F_s . It is well-known that $Z_\lambda^w(\mathbf{C}_s) = \mathcal{H}_{F_s}^{l(s)}(F; \tilde{\xi}_{w*\lambda}) [l(s)]$ is the sheaf of local cohomology of $\tilde{\xi}_{w*\lambda}$ with support on F_s , and so the g -module, corresponding to \mathbf{C}_s is the space of local cohomology $H_{F_s}^{l(s)}(F; \tilde{\xi}_{w*\lambda})$. It follows from the definition, that $H_{F_s}^{l(s)}(F; \tilde{\xi}_{w*\lambda})$ is isomorphic to $M_{(sw)*\lambda}^s$ (in particular, $H_{F_1}^0(F; \tilde{\xi}_{w*\lambda}) \simeq M_{w*\lambda}^*$ [35], $H_{F_{w_0}}^{\dim F}(F; \tilde{\xi}_{w*\lambda}) \simeq M_{(w_0w)*\lambda}$).

Thus, we see that the twisted Verma module corresponds to the sheaf of local cohomology and to the constant sheaf with support on the Schubert cell. In particular, $V_\lambda^w(M_{(sw)*\lambda}^s) \simeq \mathbf{C}_s$




Fig. 1a, b. Generalized BGG resolutions. a Usual BGG resolution. b Twisted BGG resolution

2.3. Generalized BGG Resolution

Now we pass to the Cousin-Grothendieck resolution of the invertible sheaf $\xi_{w*\lambda}$. According to the Borel-Weil-Bott theorem [9] $H^{l(w)}(F; \xi_{w*\lambda}) \simeq L_\lambda$ -irreducible representation of g with highest weight λ and $H^i(F; \xi_{w*\lambda}) = 0, i \neq l(w)$.

The spectral sequence for the computation of $H^i(F; \xi_{w*\lambda})$ associated with the filtration of F by Schubert varieties degenerates in the first term and we obtain the Cousin-Grothendieck resolution $C_{w*\lambda}^*$, whose terms are isomorphic to

$$C_{w*\lambda}^j = \bigoplus_{\substack{s \in S \\ l(s) = j}} H_{F_s}^{l(s)}(\xi_{w*\lambda}) = \bigoplus_{\substack{s \in S \\ l(s) = j}} M_{(sw)*\lambda}^s.$$

The cohomologies of this resolution are trivial in all dimensions except the $l(w)^{\text{th}}$ and the $l(w)^{\text{th}}$ cohomology is isomorphic to L_λ .

The complex C_λ^* (investigated in [35]) is contragradient to the BGG resolution [7] and consists of contragradient Verma modules. The complex $C_{w_0*\lambda}^*$ coincides with BGG resolution and consists of Verma modules.

We call $C_{w*\lambda}^*$ the *generalized strong BGG resolution*. It consists of twisted Verma modules, and its $l(w)^{\text{th}}$ cohomology group is isomorphic to irreducible representation L_λ of g . The structures of the usual and one of the twisted BGG resolutions over sl_3 are shown in Fig. 1.

2.4. Algebraic Constructions

Now we will give algebraic constructions of twisted Verma modules and generalized BGG resolutions. They will be adapted in Sect. 5 for construction of Wakimoto modules.

Let Γ_w be the Heisenberg algebra with basis $a_\alpha, a_\alpha^*, \alpha \in w(\Delta_+) = \Delta_+^w$ and commutation relations $[a_\alpha, a_\beta] = [a_\alpha^*, a_\beta^*] = 0$, $[a_\alpha, a_\beta^*] = \delta_{\alpha, \beta}$. Let π_w be an irreducible representation of Γ_w with vacuum vector, annihilated by $a_\alpha, \alpha \in \Delta_+^w \cap \Delta_+$ and by $a_\alpha^*, \alpha \in \Delta_+^w \cap \Delta_-$. Denote by A_0 the algebra of operators on π_w , commuting with $a_\alpha^*, \alpha \in \Delta_+^w$ (it consists of all polynomials in $a_\alpha^*, \alpha \in \Delta_+^w$) and by A_1 – its normalizer. Evidently, A_1/A_0 is identified with Lie algebra Vect_w of vector fields on the formal neighbourhood \mathcal{U}_w of the cell $F_w \subset F$. Lie algebra g acts on \mathcal{U}_w and hence imbeds into Vect_w , the space of embeddings being parametrized by $H^1(g, A_0) \simeq h^*$. Fix $\mu \in h^*$ and the corresponding embedding. Then π_w is g -module with highest weight μ , which is isomorphic to M_μ^w . In particular, if $w = 1$, then M_μ^w is the module of the functions on the big cell of F , which is isomorphic to the contragradient Verma module M_μ^* , if $w = w_0$, then M_μ^w is the module of δ -functions on F , isomorphic to Verma module M_μ .

Now we will give a description of the *generalized weak BGG resolutions*.

Let Ω_w^* be the de Rham complex on the formal neighbourhood \mathcal{U}_w of the cell F_w . Consider the corresponding complex $\tilde{\Omega}_w^*$ of local cohomologies:

$$H_{F_w}^{l(w)}(\mathcal{U}_w, \Omega_w^0) \rightarrow H_{F_w}^{l(w)}(\mathcal{U}_w, \Omega_w^1) \rightarrow H_{F_w}^{l(w)}(\mathcal{U}_w, \Omega_w^2) \rightarrow \dots$$

Evidently, cohomologies of this complex are non-trivial only in dimension $l(w)$ and equal to \mathbb{C} . Consider the tensor product complex $L_\lambda \otimes \tilde{\Omega}_w^* = \Omega_w^*(\lambda)$. The center of $\mathcal{U}(g)$ acts on $\tilde{\Omega}_w^*(\lambda)$ and $\Omega_w^*(\lambda)$ decomposes into a direct sum of subcomplexes, corresponding to eigenvalues θ of the center: $\Omega_w^*(\lambda) = \bigoplus \Omega_w^*(\lambda)_\theta$. The subcomplexes $\Omega_w^*(\lambda)_\theta$ are acyclic if $\theta \neq \theta(\lambda)$. The subcomplex $\Omega_w^*(\lambda)_{\theta(\lambda)}$ is the generalized weak BGG resolution $\tilde{C}_{w*\lambda}^*$. For $w = 1$ this fact was proved in [7]. In other cases the proof is analogous. $\tilde{C}_{w*\lambda}^*$ is isomorphic to $C_{w*\lambda}^*$ only if $w = 1$ or $w = w_0$.

We give algebraic construction of this complex.

Let $T^+(\mathcal{U}_w)$ be the tangent bundle over \mathcal{U}_w with changed parity of fibers. The complex Ω_w is the restriction to g of the graded module over the Lie superalgebra of vector fields on $T^+(\mathcal{U}_w)$, which contains the canonical element-differential, commuting with g .

Let Γ_w^+ be the extension of Γ_w by odd generators $\varphi_\alpha, \varphi_\alpha^*, \alpha \in \Delta_+^w$ commuting with Γ_w and with the following anti-commutation relations:

$$[\varphi_\alpha, \varphi_\beta]_+ = [\varphi_\alpha^*, \varphi_\beta^*]_+ = 0, \quad [\varphi_\alpha, \varphi_\beta^*]_+ = \delta_{\alpha, \beta}.$$

Let π_w^+ be the irreducible representation of Γ_w^+ with vacuum vector, annihilated by $a_\alpha, \varphi_\alpha, \alpha \in \Delta_+^w \cap \Delta_+$ and by $a_\alpha^*, \varphi_\alpha^*, \alpha \in \Delta_+^w \cap \Delta_-$. Introduce grading on Γ_w^+ and π_w^+ , putting $\text{deg } a_\alpha = \text{deg } a_\alpha^* = 0$, $\text{deg } \varphi_\alpha = -1$, $\text{deg } \varphi_\alpha^* = 1$.

Denote by A_0^+ the algebra of operators on π_w^+ commuting with $a_\alpha, \varphi_\alpha, \alpha \in \Delta_+^w$, and let A_1^+ be its normalizer in $\text{End } \pi_w^+$. A_1^+/A_0^+ is identified with Lie superalgebra of vector fields on $T^+(\mathcal{U}_w)$, and since g acts on $T^+(\mathcal{U}_w)$, there is the embedding of g into this superalgebra.

There is a canonical element $d \in \Gamma_w^+$ of degree 1 such that $[d, g] = 0$, $[d, d]_+ = 0$. This equips π_w^+ with the structure of the complex of g -modules, which is isomorphic to $\tilde{\Omega}_w^*$. Note that if $w = 1$, then π_w^+ and $\tilde{\Omega}_w^*$ is nothing but the de Rham complex on the big cell of F .

Using this algebraic construction we can describe the generalized BGG resolution. Introduce modified length on Weyl group. Let

$$\varphi_s = \{\alpha \in \Delta_+ : s^{-1}\alpha \in \Delta_-\}$$

and let $\Delta_{(\pm)}^w = w(\Delta_{\pm}) \cap \Delta_+$. Put

$$l_w(s) = \#(\varphi_s \cap \Delta_{(+)}^w) - \#(\varphi_s \cap \Delta_{(-)}^w).$$

In particular, $l_1(s) = l(s)$, $l_{w_0}(s) = -l(s)$.

We have $\bar{C}_{w*\lambda}^{i+l(w)} = \bigoplus_{\substack{s \in S \\ l_w(s) = i}} M_{s*\lambda}^w$. Using this result we can compute the cohomology $H^*(n_+^w, L_\lambda)$. Namely,

$$H^{i+l(w)}(n_+^w, L_\lambda) = \bigoplus_{\substack{s \in S \\ l_w(s) = i}} \mathbf{C}_{s*\lambda - w(\varrho)}$$

as the h -module, or equally, $H_s^i(n_+^w, L_\lambda) = \bigoplus_{\substack{s \in S \\ l_w(s) = i}} \mathbf{C}_{s*\lambda}$. This is the Borel-Weil-Bott

[9] theorem.

In particular, $\dim H^i(n_+^w, L_\lambda) = \# \{s : l(s) = i\}$ for any w . We can compare it with the following result:

$$\dim H^{2i}(F; \mathbf{C}) = \# \{s : l(s) = i\}, \quad H^{2i+1}(F; \mathbf{C}) = 0.$$

We will generalize this result to semi-infinite flag manifolds in Sect. 4.

In conclusion we define certain functors in the category $\text{Der } \mathcal{O}_{\theta(\lambda)}$ connected with the Weyl group. Let $T_w^s = U_\lambda^s \circ Z_\lambda^s \circ V_\lambda^w$. We have $T_w^s \circ T_{s'}^w = T_s^s$. The functor T_w^s transforms the twisted Verma module $M_{(s'w)*\lambda}^w$ to $M_{(s's)*\lambda}^s$. So T_w^s transforms the generalized BGG resolution $C_{w*\lambda}^*$ to $C_{s*\lambda}^*$, and hence L_λ to $L_\lambda[l(s) - l(w)]$. Note, that Bruhat-Hecke correspondence N_w on

$$F \times F : N_w = \{(\kappa, \kappa') \in F \times F : (b_\kappa, b_{\kappa'}) \text{ are in relative position } w\}$$

gives the functor $T_1^w = U_\lambda^1 \circ N_{w*} \circ Z_\lambda^1 \circ V_\lambda^1$. Functors T_w^s seem to be closely related to Kazhdan-Lusztig theory [33]. It would be interesting to give an algebraic construction for them.

3. Wakimoto Modules: Definition

Now we pass to representations of affine Kac-Moody algebras. The affine Kac-Moody algebra Lg^A is the unique central extension of the algebra of currents $Lg = g \otimes \mathbf{C}((t))$. The commutation relations in Lg^A read

$$[A(m), B(l)] = [A, B](m+l) + m \cdot \delta_{m, -l} \langle A, B \rangle \cdot K,$$

where $A(m)$ denotes $A \otimes t^m \in Lg$, $A \in g$, K is the central element, \langle, \rangle is Killing form, normalized so that $\langle H_i, H_j \rangle = (\bar{\alpha}_i, \bar{\alpha}_j)$, where $\bar{\alpha}_i = 2\alpha_i / (\alpha_i, \alpha_i)$, $i = 1, \dots, n$.

Let $\hat{\Delta}$ be the root system of Lg^A , $\alpha_0, \alpha_1, \dots, \alpha_n$ - be the set of simple roots, $\hat{\Delta}_+(\hat{\Delta}_-)$ denotes the set of positive (negative) roots. The roots of Lg^A are divided into imaginary and real roots. Imaginary roots are of the form $l\delta$, where $l \in \mathbf{Z}$ and $\delta = \alpha_0$

$+\alpha_{\max}$, α_{\max} being maximal root of g . Real roots are of the form $l\delta + \alpha$, where $l \in \mathbb{Z}$, $\alpha \in \Delta$.

Cartan decomposition of Lg^A is given by $Lg^A = \hat{n}_- \oplus \hat{h} \oplus \hat{n}_+$, where

$$\hat{n}_- = n_- \otimes 1 \oplus g \otimes t^{-1} \mathbf{C}[t^{-1}], \quad \hat{h} = h \otimes 1 \oplus \mathbf{C}K,$$

$$\hat{n}_+ = n_+ \otimes 1 \oplus g \otimes t \mathbf{C}[[t]].$$

Any character χ of \hat{h} determines the character of the algebra $\hat{b}_+ = \hat{h} \oplus \hat{n}_+$, $\hat{b}_+ \rightarrow \hat{h}_+ \xrightarrow{x} \mathbf{C}$ [note, that $\chi = (\bar{\chi}, k)$, $\bar{\chi} \in h^*$, $k \in (\mathbf{C}K)^* \simeq \mathbf{C}$, k is called the central charge]. Let M_χ be Verma module $\mathcal{U}(Lg^A) \otimes_{\mathcal{U}(\hat{b}_+)} \mathbf{C}_\chi$, where \mathbf{C}_χ is one-dimensional representation of \hat{b}_+ , determined by χ .

The category $\hat{\mathcal{O}}$ of highest weight Lg^A -modules [13, 41] decomposes into the direct sum of subcategories $\hat{\mathcal{O}}_\theta$, where θ is the eigenvalue of the Casimir element of Lg^A [29, 30]. Denote by $\hat{\mathcal{O}}_{\theta(\chi)}$ the subcategory containing M_χ [$\theta(\chi) = (\chi + 2\rho, \chi)$, where $\rho \in h^*$, $\rho(\alpha_i) = 1$, $i = 0, \dots, n$]. In particular, if χ is the dominant integral, then $\hat{\mathcal{O}}_{\theta(\chi)}$ consists of modules with highest weights of the form $s * \chi = s(\chi + \rho) - \rho$, where s is an element of affine Weyl group $S_{\text{aff}} = S_{\text{aff}}(G)$.

The Verma module M_χ is characterized by the following properties (cf. Sect. 2):

- a) M_χ belongs to the category $\hat{\mathcal{O}}$,
- b) $H_0(\hat{n}_-, M_\chi) \cong \mathbf{C}_\chi$ (as \hat{h} -module), $H_i(\hat{n}_-, M_\chi) = 0$, $i \neq 0$.

Now let M_χ^* be a module, contragradient to M_χ . M_χ^* is characterized by the property a) and by the property

- b*) $H^0(\hat{n}_+, M_\chi^*) \simeq \mathbf{C}_\chi$ (as \hat{h} -module), $H^i(\hat{n}_+, M_\chi^*) = 0$, $i \neq 0$.

Following Sect. 2 we should consider the flag manifold $X = X(G) = LG^A/B^A$ [39, 40]. Here LG^A denotes the unique central extension of the group of all smooth maps $S^1 \rightarrow G$ and B^A denotes Lie subgroup of LG^A corresponding to Lie algebra $\hat{b} = \hat{n}_- \oplus \hat{h}$. We also need the dense submanifold $X_{\text{an}} = X_{\text{an}}(G)$ of X , $X_{\text{an}} = L_{\text{an}}G^A/B_{\text{an}}^A$, where $L_{\text{an}}G^A$ denotes the group of analytic maps $S^1 \rightarrow G$, and B_{an}^A denotes the corresponding subgroup. Flag manifold X decomposes into disjoint union of Schubert cells X_s , where s runs the affine Weyl group S_{aff} of G , which are the orbits of \hat{N}_+ (Lie group of \hat{n}_+). $X_s = \hat{N}_+ \cdot \bar{s}$, where \bar{s} is the image of $s \in LG^A$ in X . The Schubert cell X_s is isomorphic to $s\hat{N}_+s^{-1} \cap \hat{N}_+$, and is of finite codimension.

Let χ be integral dominant weight. Then we can introduce the functors $\hat{U}_\lambda^w, \hat{V}_\lambda^w, \hat{Z}_\lambda^w$, acting between the appropriate derived categories. It is possible to construct twisted Verma modules $M_{(s_w)*\chi}^s$ as local cohomology $H_{X_s}^{l(s)}(X, \xi_{w*\chi})$, where $\xi_{w*\chi}$ denotes the suitable invertible sheaf on X . Twisted Verma modules M_μ^w are generally characterized by the property

$$\text{b}_w) \quad H^{l(w)}(w\hat{n}_+w^{-1}, M_\mu^w) \simeq \mathbf{C}_{\mu - w(\rho) + \rho},$$

$$H^i(w\hat{n}_+w^{-1}, M_\mu^w) = 0, \quad i \neq l(w).$$

Here $l(w) = \# \{ \alpha \in \hat{\Delta}_+ : w^{-1}\alpha \in \hat{\Delta}_- \}$ is the length of $w \in S_{\text{aff}}$. In particular, M_μ^1 is the contragradient Verma module M_μ^* .

Twisted Verma modules correspond to constant sheaves, supported on the Schubert cell. It is possible to construct them algebraically and also to define affine Weyl twist functors T_w^s as in 2.4. Twisted Verma modules compose twisted BGG resolutions of irreducible representation as in 2.3.

So our program may be generalized in the affine case as well. But in contrast to the finite-dimensional case, the affine Weyl group is infinite and there are no maximal elements. That is why, starting with the contragradient Verma module M_μ^* we cannot pass to the Verma module M_μ “moving” on the Schubert cell, as in the finite-dimensional case (where M_μ^* corresponds to the cell of maximal dimension and M_μ to the cell of maximal codimension – to the point), in the affine case there are no Schubert cells of maximal codimension.

But we can try to take into consideration twisted Verma modules, which correspond to certain “limit elements” of the affine Weyl group (and also “limit” cells of flag manifold).

Recall that $S_{\text{aff}} \simeq S \ltimes \mathcal{A}$, where S is the finite Weyl group and \mathcal{A} is the root lattice.

The group S_{aff} acts on $Lg^{\mathcal{A}}$ as follows:

$$s \cdot t_a(X_\alpha \otimes t^l) = X_{s(\alpha)} \otimes t^{l+(a,\alpha)}, \quad \alpha \in \mathcal{A}, s \in S, a \in \mathcal{A}.$$

Choose an element $\gamma \in \mathcal{A}$ and transform the subalgebra \hat{n}_+ by its powers:

$$t_{m\gamma}(X_\alpha \otimes t^l) = X_\alpha \otimes t^{l+m(\alpha,\gamma)},$$

and so when $m \rightarrow \infty$ we obtain the limit subalgebra $\hat{n}_+(\gamma) = \lim t_{m\gamma}(\hat{n}_+)$:

$$\hat{n}_+(\gamma) = \bigoplus_{\substack{\alpha \in \mathcal{A} \\ (\alpha,\gamma) < 0}} \mathbb{C}X_\alpha \otimes \mathbb{C}((t)) \oplus \left(h \oplus \bigoplus_{\substack{\alpha \in \mathcal{A} \\ (\alpha,\gamma) = 0}} \mathbb{C}X_\alpha \right) \otimes t\mathbb{C}[[t]] \oplus \bigoplus_{\substack{\alpha \in \mathcal{A}_+ \\ (\alpha,\gamma) = 0}} \mathbb{C}X_\alpha \otimes 1.$$

Note that $p_\gamma = h \oplus \bigoplus_{\substack{\alpha \in \mathcal{A} \\ (\alpha,\gamma) \leq 0}} \mathbb{C}X_\alpha$ is the parabolic subalgebra of \mathfrak{g} .

The algebra $\hat{n}_+(\gamma)$ depends only on p_γ , so we denote $a_p = \hat{n}_+(\gamma)$, where $p = p_\gamma$. Let $p = r_p \oplus v_p$, where v_p is the reductive subgroup of p and r_p is the nilpotent radical of p . Then

$$a_p = r_p \otimes \mathbb{C}((t)) \oplus v_p \otimes t\mathbb{C}[[t]] \oplus (v_p \cap n_+) \otimes 1.$$

We have the twisted Cartan decomposition of $Lg^{\mathcal{A}}: Lg^{\mathcal{A}} = a_p \oplus \hat{h} \oplus a_p^*$, where $*$ denotes Cartan involution [29].

It is natural to consider the infinitely twisted Verma modules $W_{\chi,p}$, corresponding to a_p (and to $\lim t_{m\gamma}$), which are characterized by the property a) and by the property

$$\begin{aligned} \text{b}_p) \quad & H^{\infty/2+i}(a_p, W_{\chi,p}) = 0, \quad i \neq 0, \\ & H^{\infty/2}(a_p, W_{\chi,p}) \simeq \mathbb{C}_\chi \quad (\text{as } \hat{h}\text{-module}) \end{aligned}$$

where $H^{\infty/2+i}(a_p, \cdot)$ denotes semi-infinite cohomology of a_p with respect to decomposition $a_p = a_p^+ \oplus a_p^-$, $a_p^\pm = a_p \cap \hat{n}_\pm$ (see Appendix A).

We call $W_{\chi,p}$ *Wakimoto modules* (note, that in [18–20] we considered only $W_{\chi,b}$). Composition series quotients of $W_{\chi,p}$ and M_χ coincide, and if M_χ is irreducible, then M_χ is isomorphic to $W_{\chi,p}$. $W_{\chi,p}$ is free over a_p^- and dual to $W_{\chi,p}$ is free over a_p^+ .

One can define $W_{\chi,p}$ as the limit of $M_\chi^{t_{m\gamma}}$ in the sense of Jantzen filtration [30]. Let $J^i W_{\chi,p}$ and $J^i M_\chi^{t_{m\gamma}}$ be the i th terms of Jantzen filtration of $W_{\chi,p}$ and $M_\chi^{t_{m\gamma}}$. Then for any j there is m_0 such that $J^i W_{\chi,p} \simeq J^i M_\chi^{t_{m\gamma}}$ for $i = 1, \dots, j$, $m > m_0$.

According to our geometric approach, we should consider the *semi-infinite flag manifold* $X_p = LG^A/B_p^A = LG/B_p$, where B_p^A is the twisted Borel subgroup corresponding to Lie algebra $\hat{b}_p = a_p^* \oplus \hat{h}$ (twisted Borel subalgebra) and its dense submanifold $X_p^{\text{an}} = L_{\text{an}}G^A/B_p^{A,\text{an}} = L_{\text{an}}G/B_p^{\text{an}}$.

In the next section we will study the geometrical and topological properties of X_p, X_p^{an} . In particular, we will show that X_p^{an} decomposes into Schubert cells of semi-infinite dimensions (that is both of infinite dimension and of infinite codimension), which are labelled by the affine Weyl group. These cells are related to Wakimoto modules in the same way as cells on X are related to contragradient Verma modules (or twisted Verma modules): the constant sheaf on the cell corresponds to Wakimoto module, and the Wakimoto module is isomorphic to the local “semi-infinite” cohomology of the invertible sheaf on X_p . Cousin-Grothendieck resolution of the invertible sheaf (with respect to Schubert stratification) gives two-sided BGG resolution of the irreducible representation, whose terms are Wakimoto modules. Explicit algebraic constructions of Wakimoto modules and of two-sided BGG resolutions (in spirit of 2.4) are given in Sects. 5 and 6.

Recall that we have defined Wakimoto modules starting with the contragradient Verma module M_μ^* and taking the limit element on the affine Weyl group. It is also possible to start with Verma module M_μ . In this way we obtain co-twisted Verma modules, characterized by the property:

$$\begin{aligned} \text{b}^w) \quad H_{l(w)}(w\hat{n}_-w^{-1}, {}^wM_\mu) &\simeq \mathbf{C}_{\mu + \varrho - w(\varrho)} \quad (\text{as } h\text{-module}), \\ H_i(w\hat{n}_-w^{-1}, {}^wM_\mu) &= 0, \quad i \neq l(w). \end{aligned}$$

Evidently, ${}^wM_\mu \cong M_\mu^{w*}$.

These modules correspond to the sheaves on the “turned” flag manifold $X^+ = LG^A/B_+^A$, where B_+^A is the Lie group corresponding to \hat{b}_+ (or on its dense submanifold $X_+^{\text{an}} = L_{\text{an}}G/B_+^{A,\text{an}}$). These sheaves are supported on Schubert cells X_s^+ of $X^+ : X_s^+ = \hat{N}_+ \cdot \bar{s}$. The cell X_s^+ is isomorphic to $s\hat{N}_-s^{-1} \cap \hat{N}_+$ and is of finite dimension.

Taking the limit, as above, we obtain contragradient Wakimoto modules $W_{\chi, p}^*$, corresponding to limit elements of S_{aff} (and to limit cells of X^+). They are characterized by the property

$$\begin{aligned} \text{b}^p) \quad H_{\infty/2+i}(a_p^*, W_{\chi, p}^*) &= 0, \quad i \neq 0, \\ H_{\infty/2}(a_p^*, W_{\chi, p}^*) &\simeq \mathbf{C}_\chi \quad (\text{as } h\text{-module}), \end{aligned}$$

where $H_{\infty/2+i}(a_p^*, \cdot)$ denotes semi-infinite homology of a_p^* with respect to the decomposition: $a_p^* = a_p^{*+} \oplus a_p^{*-}$, $a_p^{*\pm} = a_p^* \cap \hat{n}_\pm$.

The corresponding flag manifold is $X_p^+ = LG^A/B_p^{A+}$, where B_p^{A+} is the Lie group corresponding to the Lie algebra $\hat{b}_p^+ = a_p \oplus \hat{h}$.

We see that the way starting with M_μ is dual to the way starting with M_μ^* . Possibly, there are other ways.

Wakimoto modules are intermediate between M_μ and M_μ^* . The cells of X_p, X_p^+ may be considered as infinitely far cells of the usual flag manifolds X and X^+ .

So we may consider manifolds X, X_p, X_p^+, X^+ as the pieces of the unified “flag manifold” of affine Kac-Moody algebra Lg^A , whose geometry is in one-to-one correspondence with representation theory of Lg^A .

4. Semi-Infinite Flag Manifolds

In this section we will study semi-infinite flag manifolds X_p and their submanifolds.

4.1. The Usual Cohomology of X_p

The group B_p^A is homotopically equivalent to the group \hat{H} (Lie group of \hat{h}) for any p . Hence LG^A/B_p^A is homotopically equivalent to $LG^A/\hat{H} \simeq LG/H$ and to $\varphi = \tilde{\mathcal{U}}/\tilde{T}$, where $\tilde{\mathcal{U}}$ is a compact form of LG and \tilde{T} is its maximal torus, acting on $\tilde{\mathcal{U}}$ as constant maps. So we obtain the following.

Proposition 1. *For any p the semi-infinite flag manifold X_p is homotopically equivalent to the manifold $X = LG^A/\hat{B}$. In particular, the cohomology rings of X_p and X coincide, and*

$$\dim H^{2i}(X_p, \mathbb{Z}) = \# \{s \in S_{\text{aff}} : l(s) = i\}, \quad H^{2i+1}(X_p, \mathbb{Z}) = 0.$$

Note, that $H^*(X, \mathbb{Z})$ was computed on [36, 31, ...].

4.2. Loop Spaces and Semi-Infinite Structure

Let R_p be the nilpotent radical of P (Lie group of p) and $N_p = P/R_p$ the reductive subgroup of P . V_p is the product of the semi-simple Lie group G_p and abelian group H_p .

Consider the space $L(P)$ of C^∞ -maps $S^1 \rightarrow F_p = G/P$. Note that $\pi_1(L(P)) \simeq H_2(F_p, \mathbb{Z})$ is isomorphic to the group of characters $P \rightarrow \mathbb{C}^*$ (or the quotient of lattices Λ_G/Λ_{G_p}). Denote by $\tilde{L}(P)$ the universal covering space of $L(P)$. This space is isomorphic to the space of C^∞ -maps $D^2 \rightarrow F_p$, where D^2 is a closed disk, up to the following equivalence: two maps $\gamma_1 : D^2 \rightarrow F_p$ and $\gamma_2 : D^2 \rightarrow F_p$ are identified, if they coincide on the border of D^2 and are homotopically equivalent in the class of such maps.

The natural map $X_p \rightarrow L(P)$ is a fibering, the fiber being the usual (not semi-infinite) flag manifold Y_p of the group LV_p^A . Y_p is the product of $X(G_p)$ and $X(H_p)$, where $X(H_p)$ – “flag manifold” of LH_p^A – is isomorphic to the product of the group of characters of P [or to $H_2(F_p, \mathbb{Z})$] and the vector space $h_p \otimes t\mathbb{C}[t]$ (h_p is the Lie algebra of H_p). Thus, X_p is homotopically equivalent to the bundle over $\tilde{L}(P)$ with fiber $X(G_p)$. Denote it by \tilde{X}_p .

Note that $\tilde{L}(P) \simeq LG/LP_0$, where LP_0 denotes the connected component of the unit of the group LP . We put $\tilde{L}_{\text{an}}(P) = L_{\text{an}}G/L_{\text{an}}P_0$.

The manifolds $L(P)$ and $L_{\text{an}}(P)$ are endowed with remarkable semi-infinite structure. At this moment we cannot give a strict definition of such objects as semi-infinite manifolds, sheaves on them and semi-infinite cohomology of sheaves. But we are convinced that the suitable theory does exist and that manifolds $\tilde{L}(P), \tilde{L}_{\text{an}}(P)$ are the first examples of this theory.

As an illustration we will compute the semi-infinite homology of these manifolds. We will propose two ways for computations.

The first one is an application of Floer’s theory [24] which is the semi-infinite analogue of the usual Morse theory [8]. In contrast to the usual theory in Floer’s theory the indices of the singular points of the Morse function are infinite, but the difference of indices of two points is finite, so we can define the relative index

putting the index of a certain marked point to be equal to 0. Singular points compose Morse complex graded by the relative index with a differential defined as usual and semi-infinite homologies are the homologies of this complex. In our case there is a remarkable Morse function of Conley-Zehnder type [11], the complex being non-trivial only in “even” dimensions (as in finite-dimensional case [8]), and so singular points represent homology classes of $\widetilde{L}(P)$.

The second is the decomposition of $\widetilde{L}_{\text{an}}(P)$ into Schubert cells of semi-infinite dimensions. It means that cells are both of infinite dimension and of infinite codimension, but they are commensurable, so that we can define the relative dimension of the cell which is finite. Dimensions of all cells are “even” and hence they represent semi-infinite homology classes of $\widetilde{L}_{\text{an}}(P)$. Note that this cellular decomposition is related to singular points of our Morse function.

The decomposition of $\widetilde{L}_{\text{an}}(P)$ into Schubert cells gives the decomposition of X_p into Schubert cells, which are connected with Wakimoto modules. It leads to the semi-infinite analogue of Cousin-Grothendieck resolution which gives two-sided BGG resolution.

4.3. Morse Function

As is well-known, $F_p = G/P = \mathcal{U}/T_p$ (where \mathcal{U} is a compact form of G , T_p is a compact form of P) is a symplectic manifold an orbit of the coadjoint representation of \mathcal{U} . Denote the symplectic form on F_p by ω_p . Let $\tau \in t_p$ (Lie algebra of T_p) be a regular element of t_p and $h_\tau(\kappa)$ be the hamiltonian of the vector field τ on F_p . It is a Morse function on F_p with singular points-images of the elements $\tilde{s} \in S(G)/S(G_p)$ in F_p [8].

Let us define the semi-infinite Morse function $H_\tau(\gamma): \widetilde{L}(P) \rightarrow \mathbb{C}$. For any $\gamma: D^2 \rightarrow F_p$, $\gamma \in \widetilde{L}(P)$ we put

$$H_\tau(\gamma) = \int_\gamma \omega_p - \int_{\partial\gamma} h_\tau d\varphi.$$

We call $H_\tau(\gamma)$ the Conley-Zehnder function [11].

Its singular points are such maps $\gamma: D^2 \rightarrow F_p$, which transfer the border of D^2 to $\tilde{s} \in F_p$ (and winds the disk D^2 on a certain element of $H_2(F_p; \mathbb{Z}) \simeq A_G/A_{G_p}$). So singular points of $H_\tau(\gamma)$ are labelled by $S_{\text{aff}}(G)/S_{\text{aff}}(G_p)$.

Let us compute the relative indices of singular points of $H_\tau(\gamma)$.

At first, recall how to compute the indices of singular points of $h_\tau(\kappa)$. Let $\Delta_p = \Delta(G) \setminus \Delta(G_p)$, $\Delta_p = \Delta_p^+ \cup \Delta_p^-$, $\Delta_p^\pm = \Delta_p \cap \Delta_\pm$. We put

$$l_p(\tilde{s}) = \# \{ \alpha \in \Delta_p^+ : \tilde{s}\alpha \in \Delta(G)_- \}$$

(this definition is correct because $s(\Delta_p^+) = \Delta_p^+$, if $s \in S(G_p)$). The index of the point \tilde{s} is equal to $\text{ind}(\tilde{s}) = 2l_p(\tilde{s})$.

Denote $\hat{\Delta}_p = \hat{\Delta}(G) \setminus \hat{\Delta}(G_p)$. For $\tilde{s} \in S_{\text{aff}}(G)/S_{\text{aff}}(G_p)$, let

$$\begin{aligned} \widetilde{l}_p(\tilde{s}) &= \# \{ \alpha \in \hat{\Delta}_p^+ : \alpha = l\delta + \beta, \beta \in \Delta_p^+; \tilde{s}\alpha \in \hat{\Delta}(G)_- \} \\ &\quad - \# \{ \alpha \in \hat{\Delta}_p^+ : \alpha = l\delta + \beta, \beta \in \Delta_p^-; \tilde{s}\alpha \in \hat{\Delta}(G)_- \}. \end{aligned}$$

This definition is correct, because for any $b \in A_{G_p}$ and $\alpha \in \Delta_p^+$ there is $\alpha' \in \Delta_p^- : (\alpha', b) = (\alpha, b)$.

Now if we put $\text{ind}(1)=0$, then

$$\text{ind}(\tilde{s}) = 2\overline{lt}_p(\tilde{s}), \quad \tilde{s} \in S_{\text{aff}}(G)/S_{\text{aff}}(G_p).$$

So semi-infinite homologies of $\widehat{L}(P)$ are contained in even dimensions $2i$ and generated over \mathbf{C} by elements of the set $M_p(i)$, where $M_p(i) = \{\tilde{s} \in S_{\text{aff}}(G)/S_{\text{aff}}(G_p) : \overline{lt}_p(\tilde{s}) = i\}$.

4.4. Cellular Decomposition

The cells of $\widehat{L}_{\text{an}}(P)$ are the orbits of the group \widehat{N}_+ . Let $\overline{\mathcal{U}}_s = \widehat{N}_+ \cdot \tilde{s}$, $\tilde{s} \in S_{\text{aff}}(G)/S_{\text{aff}}(G_p)$, \tilde{s} is embedded onto $\widehat{L}_{\text{an}}(P)$. Then $\overline{\mathcal{U}}_s$ is identified with $\widehat{N}_p \cap \tilde{s}\widehat{N}_+ \tilde{s}^{-1}$, where \widehat{N}_p is the subgroup of $L_{\text{an}}G^A$, which corresponds to Lie algebra $r_p \otimes \mathbf{C}((t))$ [note that $\widehat{N}_p \cap s\widehat{N}_+ s^{-1} = \widehat{N}_p \cap \widehat{N}_+$, $s \in S_{\text{aff}}(G_p)$]. So, putting $\dim \overline{\mathcal{U}}_1 = 0$ we obtain that relative dimension $\dim \overline{\mathcal{U}}_s = 2\overline{lt}_p(\tilde{s})$. It agrees with Sect. 4.3. Note that all cells are of “equal size” and the closure of $\overline{\mathcal{U}}_s$ consists of $\overline{\mathcal{U}}_s$, such that $\dim \overline{\mathcal{U}}_s < \dim \overline{\mathcal{U}}_s$.

We also obtain the decomposition of X_p^{an} into Schubert cells. Let $s \in X_p$ be the image of $s \in S_{\text{aff}}(G)$ in X_p^{an} . Then the orbits of s under the action of \widehat{N}_+ give this decomposition. Denote $\mathcal{U}_s = \widehat{N}_+ \cdot s \subset X_p^{\text{an}}$. Evidently, \mathcal{U}_s is the product of the appropriate cells of $\widehat{L}_{\text{an}}(P)$ and Y_p . The relative dimension of \mathcal{U}_s (we put $\dim \mathcal{U}_1 = 0$) is equal to $2\overline{lt}_p(s)$, defined as follows. We have $\Delta(G) = \Delta(P) \cup \Delta^p$. Let

$$\widehat{\Delta}(P) = \{l\delta + \alpha, \alpha \in \Delta(P), l \in \mathbf{Z}\}, \quad \widehat{\Delta}^p = \{l\delta + \alpha, \alpha \in \Delta^p, l \in \mathbf{Z}\}.$$

Then

$$\overline{lt}_p(s) = \# \{\alpha \in \widehat{\Delta}(P) \cap \widehat{\Delta}(G)_+ : s\alpha \in \widehat{\Delta}(G)_-\} - \# \{\alpha \in \widehat{\Delta}^p \cap \widehat{\Delta}(G)_+ : s\alpha \in \widehat{\Delta}(G)_-\}.$$

In particular,

$$\begin{aligned} \overline{lt}_b(s) &= \# \{\alpha \in \widehat{\Delta}(G)_+ : \alpha = l\delta + \beta, \beta \in \Delta_+, s\alpha \in \widehat{\Delta}(G)_-\} \\ &\quad - \# \{\alpha \in \widehat{\Delta}(G)_+ : \alpha = l\delta + \beta, \beta \in \Delta_-, s\alpha \in \widehat{\Delta}(G)_-\}, \end{aligned}$$

and if $s = s' \cdot t_\gamma$, $s' \in S(G)$, $b \in \Delta_G$, then $\overline{lt}_\gamma(s) = l(s') + (2Q, \gamma)$. We define also $lt_p(s) = \overline{lt}_p(s^{-1})$.

4.5. Connection with Wakimoto Modules

We suppose that in a semi-infinite case there are analogues of functors U, V, Z of Sect. 2, 3, which establish the connection of Wakimoto modules $W_{\chi, p}$ with sheaves on X_p .

The Wakimoto module $W_{s^* \chi, s p s^{-1}}$ corresponds to the space of “local semi-infinite cohomology” with support on the Schubert cell of the invertible sheaf $H_{\overline{\mathcal{U}}_s}^{\infty/2 - \overline{lt}_p(s)}(X_p, \xi_\gamma)$, where $\overline{lt}_p(s)$ is the (complex) relative codimension of the cell $\overline{\mathcal{U}}_s$. In the next section we will give algebraic construction of Wakimoto modules which is similar to finite-dimensional construction of 2.4 and clarifies the notion of local semi-infinite cohomology.

The constructible sheaf on X_p , corresponding to the Lg^A -module M , has the stalk at the point κ , isomorphic to the complex of semi-infinite homology (more exactly, its $w^* \chi$ eigenvalue component with respect to the action of \widehat{h})

$C_{\infty/2+*}^{w*\chi}(a_{p,\kappa}^*, M)$, where $a_{p,\kappa}^* = \kappa a_{p,\kappa}^* \kappa^{-1} = a_p^*$ is the Lie algebra of the stabilizer of $\kappa \in X_p$ and we take its decomposition

$$a_{p,\kappa}^* = a_{p,\kappa}^{*+} \oplus a_{p,\kappa}^{*-}, \quad a_{p,\kappa}^{*\pm} = \kappa(a_{p,\kappa}^* \cap \hat{n}_{\pm})\kappa^{-1}.$$

So, the Wakimoto module $W_{(sw)*\chi, sps^{-1}}$ corresponds to the constant sheaf on the Schubert cell \mathcal{U}_s . The irreducible representation with dominant integral highest weight L_χ corresponds to the constant sheaf on X_p . Irreducible representations $L_{s*\chi}$ correspond to semi-infinite analogues of Goresky-MacPherson sheaves [4].

4.6. Cousin-Grothendieck Resolution

There is a semi-infinite analogue of Cousin-Grothendieck resolution $R_p^{\infty/2+*}(\chi)$ of invertible sheaf $\tilde{\xi}_\chi$ with respect to Schubert stratification of X_p . Its terms are local cohomologies of $\tilde{\xi}_\chi$ with support on the Schubert cell, that is Wakimoto modules:

$$R_p^{\infty/2+i}(\chi) = \bigoplus_{\substack{\text{codim } \mathcal{U}_s = i \\ s \in S_{\text{aff}}}} H_{\mathcal{U}_s}^{\infty/2+i}(X_p, \tilde{\xi}_\chi) \simeq \bigoplus_{\substack{s \in S_{\text{aff}} \\ \bar{l}_p(s) = -i}} W_{s*\chi, sps^{-1}}.$$

The cohomologies of this complex coincide with cohomologies of $\tilde{\xi}_\chi$:

$$H^{\infty/2+i}(X_p, \tilde{\xi}_\chi) = 0, \quad i \neq 0, \quad H^{\infty/2}(X_p, \tilde{\xi}_\chi) \simeq L_\chi.$$

[analogously, $H^{\infty/2+i}(X_p, \tilde{\xi}_{s*\chi}) = 0$, if $i \neq \bar{l}_p(s)$, $H^{\infty/2+\bar{l}_p(s)}(X_p, \tilde{\xi}_{s*\chi}) \simeq L_\chi$]. This is the analogue of the Borel-Weil-Bott theorem.

So $R_p^{\infty/2+*}(\chi)$ is the resolution of an irreducible representation. It is the limit resolution of twisted resolutions on the usual flag manifold X . We call it two-sided BGG resolution. This is an analogue of generalized strong BGG resolution.

In Sect. 6 we will prove a similar two-sided BGG resolution in a different way. It is an analogue of generalized weak BGG resolution.

4.7. Grassmanian Model

There is a Grassmanian model of $L(\widetilde{P})$ similar to the Grassmanian model of the usual flag manifold [40]. We give this model for $L_{\text{pol}}SL_2$ [40], the other models for other algebras are direct generalizations of this.

Let $\mathcal{H} = L^2(S^1, \mathbb{C}^2)$ be Hilbert space. Let e_1 and e_2 be basic vectors in \mathbb{C}^2 . We choose the basis u_i , $i \in \mathbb{Z}$ in \mathcal{H} , such that $u_{2i} = e_1 z^i$, $u_{2i-1} = e_2 z^i$. The shift operator z transforms u_i to u_{2i+1} .

There is decomposition of \mathcal{H} into a direct sum of mutually orthogonal subspaces:

$$\mathcal{H} = H_+ \oplus H_-^{(1)} \oplus H_-^{(2)},$$

where H_+ is generated (over \mathbb{C}) by u_i , $i > 0$, $H_-^{(1)}$ – by u_{2i} , $i \leq 0$, $H_-^{(2)}$ – by u_{2i-1} , $i \leq 0$.

Let us consider the manifold Gr of all subspaces H of \mathcal{H} , commensurable with $H_-^{(1)}$ or $H_-^{(2)}$ (see [40]), characterized by the property:

$$zH \subset H \text{ and } z^{n-1}H/z^nH \text{ is one-dimensional.}$$

The manifold Gr is isomorphic to $\widetilde{L(B)}_{\text{pol}}$ of Lsl_2 . The group $L_{\text{pol}}SL_2$ acts on Gr naturally, the stabilizer of $H^{(1)}$ being the group $L_{\text{pol}}B_0$ (the connected component of the unit of $L_{\text{pol}}B$).

We say that an element $u \in \mathcal{H}$ is of finite order q if $u = \sum_{i \geq q} c_i u_i, c_i \in \mathbb{C}$. For $H \subset \mathcal{H}$ we put $Q_H = \{q \in \mathbb{Z} : H \text{ contains an element of order } q\}$. For the set of integers Q we put $\sum_Q = \{H \in Gr : Q_H = Q\}$.

The affine Weyl group $S_{\text{aff}}(SL_2)$ is freely generated by s_0 and s_1 . Let us denote

$$s_{(i)} = \underbrace{s_{0 \text{ or } 1} \dots s_1 s_0 s_1}_{-i \text{ times}}, \quad i \leq 0,$$

$$s_{(i)} = \underbrace{s_{0 \text{ or } 1} \dots s_0 s_1 s_0}_{i \text{ times}}, \quad i > 0.$$

Then $\bar{u}_{s(r)}$ – the cell of $\widetilde{L(B)}$ coincides with \sum_{Q_i} , where

$$Q_i = \{i, i-2, i-4, \dots\}.$$

It is the orbit of $H_{Q_i} = s_{(i)}$ under the action of \hat{N}_+ .

5. Wakimoto Modules: Construction

In this section we will construct Wakimoto modules algebraically as certain modules over the algebra of infinitesimal automorphisms of the bundle over the formal neighbourhood of the cell of the manifold $\widetilde{L(p)}$. This is the generalization of the finite-dimensional construction of twisted Verma modules 2.4. But in the infinite-dimensional case we meet with some homological problems.

5.1. The Case of Borel Subalgebra

We start with the most important case $p=b$. Denote $W_x = W_{x,b}$. Wakimoto modules W_x are boson representations of Lg^A , which are interesting in conformal field theory. It is possible to obtain explicit formulae for these representations. The formulae for Lsl_n^A are represented in Sect. 7.

Consider the formal neighbourhood \mathcal{N} of the cell \bar{Q}_1 of $\widetilde{L(B)}$. It is isomorphic to the linear space $N_- \otimes \mathbb{C}((t)) \simeq \mathbb{C}^{d(g)}((t)), d(g) = (\dim g - n)/2$.

First of all let us study Lie algebra of vector fields on \mathcal{N} .

Let $\bar{\Gamma}$ be Heisenberg algebra with generators $a_\alpha(m), a_\alpha^*(m), \alpha \in \Delta_-, m \in \mathbb{Z}$ and commutation relations

$$[a_\alpha(m), a_\beta(l)] = [a_\alpha^*(m), a_\beta^*(l)] = 0, \quad [a_\alpha(m), a_\beta^*(l)] = \delta_{\alpha,\beta} \delta_{m,-l}. \tag{1}$$

Let \bar{M} be an irreducible representation of $\bar{\Gamma}$ with vacuum vector, annihilated by $a_\alpha(m), m > 0, \alpha \in \Delta_-, a_\alpha^*(m), m \geq 0, \alpha \in \Delta_-$. Introduce grading on $\bar{\Gamma}$ and \bar{M} , putting $\text{deg } a_\alpha^*(m) = 1, \text{deg } a_\alpha(m) = -1$. Denote by A_0 the algebra of operators $\bar{M} \rightarrow \bar{M}$, commuting with all $a_\alpha^*(m)$ and by A_1 its normalizer in $\text{End } \bar{M}$. The space $A_1/A_0 = W = \bigoplus_{i \geq -1} W_i$ is a graded Lie algebra.

It is easy to see that W_1 consists of the operators $\sum_{\substack{\alpha \in A \\ m \geq l}} \varrho_\alpha(m) a_\alpha(m)$, where $\varrho_\alpha(m) \in \mathbf{C}, l \in \mathbf{Z}$. So W_{-1} is identified with \mathcal{N} . Denote $\mathcal{N}_+ = N_- \otimes t\mathbf{C}[[t]]$. We have

Proposition 2 [18]. 1. *The Lie algebra W_0 is isomorphic to the algebra of operators $a: \mathcal{N} \rightarrow \mathcal{N}$, such that $\dim((a(\mathcal{N}_+) + \mathcal{N}_+)/\mathcal{N}_+) < \infty$; $W_0 \oplus W_{-1}$ is normalizer of W_{-1} in W ;*

2. *The Lie algebra W is Cartan prolongation of the pair (W_0, W_{-1}) . W is identified with the Lie algebra of vector fields on \mathcal{N} .*

It is interesting to compute the cohomology of W with coefficients in A_0 , which is identified with the space of the functions on \mathcal{N} .

We can change W by the Lie algebra W_∞ of “finite” vector fields, which is the Cartan prolongation of the pair (gl_∞, V_∞) , where gl_∞ is the injective limit of gl_n and V_∞ is the injective limit of vector representation V_n of gl_n . Denote by $F(V_\infty)$ the space of functions on V_∞ and consider $H^*(W_\infty, F(V_\infty))$. $W_\infty = W_{\infty, -1} \oplus \tilde{W}_\infty$, where $\tilde{W}_\infty \simeq \bigoplus_{i \geq 0} W_{\infty, i}$. As is well-known,

$$H^*(W_\infty, F(V_\infty)) \simeq H^*(\tilde{W}_\infty, \mathbf{C}) \simeq H^*(gl_\infty, \mathbf{C}) \simeq A^*(e_1, e_2, \dots),$$

$$\deg e_i = 2i - 1, \quad i = 1, 2, \dots$$

Here the first isomorphism is given by the Schapiro lemma and the second is proved in [25].

We obtain the analogous result for W .

Theorem 1. $H^*(W, A_0) \simeq H^*(W_0, \mathbf{C})$.

Proof. The isomorphism $H^*(W, A_0) \simeq H^*(\bigoplus_{i \geq 0} W_i, \mathbf{C})$ is given by the Schapiro lemma and the proof of the isomorphism

$$H^*(\bigoplus_{i \geq 0} W_i, \mathbf{C}) \simeq H^*(W_0, \mathbf{C})$$

is the same as in [25]. \square

The cohomology ring $H^*(W_0, \mathbf{C})$ was computed in [22].

$$H^*(W_0, \mathbf{C}) \simeq S^*(c_1, c_2, \dots), \quad \deg c_i = 2i, \quad i = 1, 2, \dots$$

Note that $H^2(W_0, \mathbf{C}) \simeq \mathbf{C}$ is generated by the well-known Tate [42] or “Japanese” [12] or “wedge” [31a] cocycle c_1 and so $H^2(W, A_0) \simeq \mathbf{C}$. We can give the description of the cocycle \tilde{c}_1 generating $H^2(W, A_0)$. Let W_1 and W_2 be two vector fields on \mathcal{N} from W . Any point $\kappa \in \mathcal{N}$ determines two elements $\bar{w}_1(\kappa)$ and $\bar{w}_2(\kappa)$ from W_0 , which are linearizations of W_1 and w_2 at κ . We put

$$[\tilde{c}_1(w_1, w_2)](\kappa) = c_1(\bar{w}_1(\kappa), \bar{w}_2(\kappa)).$$

Now we pass to Lg^A . It is clear that Lg acts on \mathcal{N} and it gives the embedding $Lg \hookrightarrow W$.

Proposition 3 [18]. *Composition*

$$H^2(W, A_0) \rightarrow H^2(Lg, A_0) \rightarrow H^2(Lg, A_0/\mathbf{C}),$$

applied to \tilde{c}_1 is 0.

Proof. Short exact sequence $0 \rightarrow \mathbb{C} \rightarrow A_0 \rightarrow A_0/\mathbb{C} \rightarrow 0$ gives

$$H^2(Lg, \mathbb{C}) \xrightarrow{\varepsilon_1} H^2(Lg, A_0) \xrightarrow{\varepsilon_2} H^2(Lg, A_0/\mathbb{C}).$$

According to Schapiro lemma,

$$H^2(Lg, A_0) \simeq H^2(Lb_+, \mathbb{C}) \simeq H^2(Lh, \mathbb{C}).$$

Denote the image of the projection of \tilde{c}_1 onto $H^2(Lg, A_0)$ by \bar{c}_1 , and the corresponding element of $H^2(Lh, \mathbb{C})$ by \bar{c} . We should show that \bar{c} lies in the image of ε_1 . Recall that $H^2(Lg, \mathbb{C}) \simeq \mathbb{C}$ is generated by the central charge. It means that we should show that \bar{c} (central extension of Lh) is the restriction of the central extension of Lg . It follows from the computations below.

Let us compute \bar{c} . In order to do it we should express operators $H_\alpha(m) \in Lg^A$ (H_α is the coroot, dual to the root $\alpha \in A$) via operators $a_\beta(m), a_\beta^*(m)$.

We need some preparations. For any countable set $A(m), m \in \mathbb{Z}$, of operators we put $A(z) = \sum_{m \in \mathbb{Z}} A(m)z^m, \dot{A}(z) = z \frac{d}{dz} A(z)$. Introduce normal ordering $::$, as usual [19].

Operators $H_\alpha(m)$ act on \bar{M} as follows:

$$H_\alpha(z) = \sum_{\beta \in A_-} (\bar{\alpha}, \beta) : a_\beta(z) a_\beta^*(z) ::$$

Using the Wick theorem we obtain:

$$\begin{aligned} [H_\alpha(m), H_\beta(l)] &= -\delta_{m, -l} \cdot m \sum_{\gamma \in A_+} (\bar{\alpha}, \gamma) (\bar{\beta}, \gamma) \\ &= \delta_{m, -l} \cdot m (\bar{\alpha}, \bar{\beta}) (-c_g), \end{aligned}$$

where c_g is dual Coxeter number of g [29]. We see that \bar{c} is the restriction of the central extension of Lg and Proposition 3 follows. \square

We obtain the following result.

Theorem 2 [18]. *The embedding $Lg \hookrightarrow W$ is lifted to the embedding $Lg^A \hookrightarrow A_1$. The set of these embeddings is the principal homogeneous bundle over $H^1(Lg, A_0)$. There is natural homomorphism $h^* \rightarrow H^1(Lg, A_0)$. There is an n -parameter family of Lg^A -modules $\bar{W}_\chi, \chi \in h^*$ with central charge $-c_g$ in \bar{M} .*

We call \bar{W}_χ a restricted Wakimoto module. It is characterized by the homological property:

$$H^{\infty/2}(r_p \otimes \mathbb{C}((t)), \bar{W}_\chi) \simeq \mathbb{C}_\chi; \quad H^{\infty/2+i}(r_p \otimes \mathbb{C}((t)), \bar{W}_\chi) = 0, \quad i \neq 0.$$

\bar{W}_χ corresponds to the cell of $\overline{L(\bar{B})}$. We studied \bar{W}_χ in [19] (see also Appendix B).

In order to construct Wakimoto modules W_χ we should consider the fibering over \mathcal{N} with the fiber π_k -Fock representation of Lh^A with vacuum vector annihilated by $h \otimes \mathbb{C}[[t]]$ and central charge k' .

So we should consider the extension Γ of $\bar{\Gamma}$ by generators $b_i(m), i = 1, \dots, n, m \in \mathbb{Z}$, commuting with Γ and with commutation relations $[b_i(m), b_j(l)] = \delta_{m, -l} \langle H_i, H_j \rangle$. The subalgebra of Γ is identified with the algebra of infinitesimal automorphisms of our fibering. So Lg^A imbeds into Γ and it equips $\bar{W}_\chi \otimes \pi_k$ with the structure of an Lg^A -module with central charge $k = k' - c_g$ and highest weight $\chi = (\bar{\chi}, k)$. So we obtain an $(n + 1)$ -parameter family of Lg^A -modules, which is the family of Wakimoto modules W_χ [18].

5.2. General Case

Now we will construct $W_{\chi, p}$ with an arbitrary parabolic subalgebra p of \mathfrak{g} .

We introduce some notations. The subgroup V_p of G is the product of semi-simple subgroup: $G_p = G_p^{(1)} \cdot \dots \cdot G_p^{(q)}$ (where $G_p^{(i)}$ are simple subgroups of G_p) and of the abelian subgroup H_p . We assume that corresponding Lie algebras $\mathfrak{g}_p^{(i)}$, $i = 1, \dots, q$ and \mathfrak{h}_p are mutually orthogonal with respect to the Killing form.

Let $M_{(\chi, \mathbf{k}_p)}^p$ be the contragradient Verma module over $L\mathfrak{g}_p^A$:

$$M_{(\chi, \mathbf{k}_p)}^p = M_{(\chi_1, \mathbf{k}_p^{(1)})}^* \otimes \dots \otimes M_{(\chi_q, \mathbf{k}_p^{(q)})}^*,$$

where $M_{(\chi_i, \mathbf{k}_p^{(i)})}^*$ is a contragradient Verma module over $L\mathfrak{g}_p^{(i)A}$. Let $\pi_{(\chi', \mathbf{k}_p')}$ be a Fock representation of $L\mathfrak{h}_p^A$ with central charge \mathbf{k}_p' and highest weight χ' .

Consider the fibering over the formal neighbourhood \mathcal{N}_p of the cell $\overline{\mathcal{U}}_1 \subset \overline{L(P)}$ with fiber $M_{(\chi, \mathbf{k}_p)}^p \otimes \pi_{(\chi', \mathbf{k}_p')} = M_{\chi, \chi', \mathbf{k}_p, \mathbf{k}_p'}^p$.

We define the algebra W^p of vector fields on the space \mathcal{N}_p [which is isomorphic to $R_p \otimes \mathbb{C}((t))$] in the same way as in 5.1. Let Γ_p be a Heisenberg algebra with generators $a_\alpha(m)$, $a_\alpha^*(m)$, $\alpha \in \Delta^p$, $m \in \mathbb{Z}$ and commutation relations (1). Let \overline{M}_p be its irreducible representation with a vacuum vector, annihilated by $a_\alpha(m)$, $\alpha \in \Delta^p$, $m > 0$ and $a_\alpha^*(m)$, $\alpha \in \Delta^p$, $m \geq 0$. Denote by A_0^p the algebra of operators $\overline{M}_p \rightarrow \overline{M}_p$, commuting with $a_\alpha^*(m)$, $\alpha \in \Delta^p$, $m \in \mathbb{Z}$, and by A_1^p its normalizer $W^p = A_1^p/A_0^p$ is identified with a Lie algebra of vector fields in $\text{End } \overline{M}_p$ on \mathcal{N}_p .

The algebra Aut_p of infinitesimal automorphisms of the fibering defined above is a semi-direct sum of W^p and of the algebra

$$\text{End } M_{\chi, \chi', \mathbf{k}_p, \mathbf{k}_p'}^p \otimes A_0^p = \text{End}_p.$$

The extension of W^p by A_0^p , defined in 5.1 gives the extension of W^p by End_p and of Aut_p by End_p . Denote by \tilde{c} the element of $H^2(\text{Aut}_p, \text{End}_p)$, which corresponds to the representation of Aut_p in $M_{\chi, \chi', \mathbf{k}_p, \mathbf{k}_p'}^p \otimes \overline{M}_p$.

We want to obtain the condition, when $L\mathfrak{g}^A$ may be embedded into Aut_p . $L\mathfrak{g}$ acts on \mathcal{N}_p and hence imbeds into W^p . So $L\mathfrak{g}^A$ imbeds into Aut_p if and only if the composition

$$H^2(\text{Aut}_p, \text{End}_p) \rightarrow H^2(L\mathfrak{g}, \text{End}_p) \rightarrow H^2(L\mathfrak{g}, \text{End}_p/\mathbb{C}),$$

applied to \tilde{c} gives 0.

Proposition 4. *The composition*

$$H^2(\text{Aut}_p, \text{End}_p) \rightarrow H^2(L\mathfrak{g}, \text{End}_p) \rightarrow H^2(L\mathfrak{g}, \text{End}_p/\mathbb{C}),$$

applied to \tilde{c} , gives 0, if and only if the following conditions are satisfied:

$$\mathbf{k}_p^{(i)} = \mathbf{k}_p' - c_{\mathfrak{g}_p^{(i)}}.$$

Proof. The short exact sequence $0 \rightarrow \mathbb{C} \rightarrow \text{End}_p \rightarrow \text{End}_p/\mathbb{C} \rightarrow 0$ gives:

$$H^2(L\mathfrak{g}, \mathbb{C}) \rightarrow H^2(L\mathfrak{g}, \text{End}_p) \rightarrow H^2(L\mathfrak{g}, \text{End}_p/\mathbb{C}).$$

As in Proposition 3, we see that, according to the Schapiro lemma,

$$H^2(L\mathfrak{g}, \text{End}_p) \simeq H^2(L_p, M_{\chi, \chi', \mathbf{k}_p, \mathbf{k}_p'}^p) = \bigoplus_{i=1}^q H^2(L\mathfrak{g}_p^{(i)}, M_{(\chi_i, \mathbf{k}_p^{(i)})}^*) \oplus H^2(L\mathfrak{h}_p, \pi_{\chi', \mathbf{k}_p'}),$$

and we should show that the restriction of \tilde{c} to the latter cohomology space is the restriction of the central charge of Lg . In order to do it we should compute the extension on $h \subset p$.

For any $H_\beta \in h$ we have $H_\beta = H_\beta^p + \sum_{i=1}^q H_{\beta, i}$, where $H_\beta^p \in h_p$ and $H_{\beta, i} \in g_p^{(i)}$.

We have the action of $H_\beta(m)$ in $M_{\chi, \chi', k_p, k_p'}^* \otimes \bar{M}_p$:

$$H_\beta^{(z)} = \sum_{\alpha \in \Delta^p} : a_\alpha(z) a_\alpha^*(z) : (\alpha, \bar{\beta}) + \bar{H}_\beta^p(z) + \bar{H}_{\beta, i}(z),$$

where $H_\beta^p(m)$ denotes the action of $H_\beta^p(m)$ in $\pi_{\chi', k_p'}$, $\bar{H}_{\beta, i}(m)$ denotes the action of $H_{\beta, i}(m)$ in $M_{\chi_i, k_p^{(i)}}^*$.

We have:

$$[H_\beta(m), H_\gamma(l)] = m\delta_{m, -l} \left(- \sum_{\alpha \in \Delta^p} (\alpha, \bar{\beta}) (\alpha, \bar{\gamma}) + \sum_{i=1}^q k_p^{(i)} \langle H_{\beta, i}, H_{\gamma, i} \rangle + k_p' \langle H_\beta^p, H_\gamma^p \rangle \right). \tag{2}$$

Evidently, $\langle H_\beta, H_\gamma \rangle = \sum_{i=1}^q \langle H_{\beta, i}, H_{\gamma, i} \rangle + \langle H_\beta^p, H_\gamma^p \rangle$. We must obtain

$$[H_\beta(m), H_\gamma(l)] = m\delta_{m, -l} (\bar{\beta}, \bar{\gamma}) \cdot k. \tag{3}$$

So we see that the following conditions must be satisfied for it:

$$k_p^{(i)} = k_p' - c_{g_p^{(i)}}, \tag{4}$$

(where $c_{g_p^{(i)}}$ denotes dual Coxeter number of $g_p^{(i)}$).

Then (2) is (3) with $k = k_p' - c_g$.

The proposition is proved. \square

So we obtain that if (4) is satisfied, then $M_{\chi, \chi', k_p, k_p'}^* \otimes \bar{M}_p$ is equipped with the structure of an Lg^A -module, which is isomorphic to $W_{\chi, p}$, where $\chi = (\chi, \chi', k)$, with central charge $k = k_p' - c_g$.

Our construction shows that $W_{\chi, p}$ is the Lg^A -module, “semi-infinitely induced” from the $(Lg_p \oplus Lh_p)^A$ -module $M_{\chi, \chi', k_p, k_p'}^*$ and the existence of a central extension imposes the constraints (4) on the central charges. Indeed, it is possible to induce from any representation of $(Lg_p \oplus Lh_p)^A$, if the conditions (4) are satisfied.

In particular, if we induce from a Verma module, then we obtain the module $W_{\chi, p}^*$ corresponding to the cell of $X_{p^*}^+$.

Our construction also says that it is possible to put the sheaf of $(Lg_p \oplus Lh_p)^A$ -modules on $\widehat{L(p)}$ into correspondence with highest weight Lg^A -module, by taking the complex of semi-infinite homology $C_{\infty/2+*}(r_p \otimes \mathbf{C}((t)), M)$ as the stalk of this sheaf. The corresponding sheaves may be lifted (in some sense) to the constructible sheaves on X_p (see Sect. 4).

6. Two-Sided Bernstein-Gelfand-Gelfand Resolutions

In this section we construct and prove two-sided Bernstein-Gelfand-Gelfand (BGG) resolutions of irreducible module over Lg^A , consisting of Wakimoto modules. The approach we develop is an alternative to that of Sect. 4, where similar resolutions appeared to be Cousin-Grothendieck on the semi-infinite flag manifold.

Our resolutions are the subcomplexes of the semi-infinite de Rham complex in the neighbourhood of the big cell on the semi-infinite flag manifold (for a finite-dimensional counterpart see 2.4).

Let Γ_p^+ be the supersymmetric extension of Γ_p by odd generators $\varphi_\alpha(m)$, $\varphi_\alpha^*(m)$, $\alpha \in \Delta^p$, $m \in \mathbb{Z}$, commuting with Γ with the following anticommutation relations:

$$\begin{aligned} [\varphi_\alpha(m), \varphi_\beta(l)]_+ &= [\varphi_\alpha^*(m), \varphi_\beta^*(l)]_+ = 0, \\ [\varphi_\alpha(m), \varphi_\beta^*(l)]_+ &= \delta_{\alpha, \beta} \delta_{m, -l}. \end{aligned}$$

Let M_p^+ be the irreducible representation of Γ_p^+ with vacuum vector annihilated by $a_\alpha(m)$, $\varphi_\alpha(m)$, $\alpha \in \Delta^p$, $m > 0$ and by $a_\alpha^*(m)$, $\varphi_\alpha^*(m)$, $\alpha \in \Delta^p$, $m \geq 0$. Introduce grading on Γ_p^+ and M_p^+ putting $\deg a_\alpha(m) = \deg \varphi_\alpha(m) = -1$, $\deg a_\alpha^*(m) = \deg \varphi_\alpha^*(m) = 1$. Denote by A_0^p the algebra of operators $M_p^+ \rightarrow M_p^+$, commuting with all $a_\alpha^*(m)$, $\varphi_\alpha^*(m)$ and let A_1^p be the normalizer of A_0^p in the Lie superalgebra. The superalgebra $A_1^p/A_0^p = W_p^+ = \bigoplus_{i \geq -1} W_{p,i}^+$ is a graded Lie superalgebra. We see that $W_{p,-1}^+$ consists of the operators

$$\sum_{\alpha \in \Delta^p, m \geq l} \varrho_\alpha(m) a_\alpha(m) + \sum_{\alpha \in \Delta^p, m \geq l'} \varrho'_\alpha(m) \varphi_\alpha(m), \quad \text{where } l, l' \in \mathbb{Z},$$

and so $W_{p,-1}^+$ is identified with the tangent bundle $T^+ \mathcal{N}_p$ over \mathcal{N}_p [the formal neighbourhood of the cell $\overline{\mathcal{W}}_1 \subset \overline{L(p)}$] with the changed parity of fibers. W_p^+ is identified with Lie superalgebra of vector fields on $T^+ \mathcal{N}_p$. In the same way as in Sect. 5, we obtain: $H^2(W_p^+, A_0^p) \simeq \mathbb{C}$. The generator \tilde{c}_1^+ of $H^2(W_p^+, A_0^p)$ is induced by Tate-“Japanese” cocycle c_1^+ from $H^2(W_{p,0}^+, \mathbb{C}) \simeq \mathbb{C}$.

Now change the grading on Γ_p^+ and M_p^+ , putting $\deg_0 a_\alpha(m) = \deg_0 \varphi_\alpha^*(m) = 0$, $\deg \varphi_\alpha(m) = -1$, $\deg \varphi_\alpha^*(m) = 1$. There is a canonical element \tilde{d} (differential) in Γ_p^+ such that $\deg_0 \tilde{d} = 1$, $[\tilde{d}, \tilde{d}]_+ = 0$, which endows M_p^+ with the structure of complex $R_p^* = \bigoplus_{i \in \mathbb{Z}} R_p^i$. It is evident that their cohomologies are contained in the 0th dimension and equal to \mathbb{C} . This complex is the semi-infinite analogue of de Rham complex on \mathcal{N}_p .

On the other hand, let us consider the usual de Rham complex Ω_p^* on the big cell X_1 of the usual flag manifold $X(G_p)$ of the group LG_p^A , $\Omega_p^* = \bigoplus_{i \geq 0} \Omega_p^i$. Denote by F_p the space of the functions on X_1 . The tensor product complex $\bar{R}_p^* = R_p^* \otimes \Omega_p^* = \bigoplus_{i \in \mathbb{Z}} \bar{R}_p^i$ may be considered as the semi-infinite de Rham complex on the product of X_1 and \mathcal{N}_p . Denote by d its differential. Note that the product of \mathcal{N}_p and X_1 is isomorphic to the cell of the manifold \bar{X}_p (see 4.2). Denote by \bar{W}_p^+ the Lie superalgebra of vector fields on $T^+(\mathcal{N}_p \times X_1)$ (tangent bundle over $\mathcal{N}_p \times X_1$ with changed parity of fibers). The Lie algebra Lg acts on $T^+(\mathcal{N}_p \times X_1)$ and hence Lg imbeds into \bar{W}_p^+ , the image commuting with d . We have the following

Proposition 5. *The composition*

$$H^2(W_p^+; A_0^p) \rightarrow H^2(\bar{W}_p^+, A_0^p \otimes F_p) \rightarrow H^2(Lg \oplus \mathbb{C}d, A_0^p \otimes F_p)$$

(the first mapping is due to the Schapiro lemma) transforms \tilde{c}_1^+ to 0.

Proof is analogous to the proof of Propositions 3 and 4 (we should consider the superconformal current algebra Lg^+ [32] which imbeds into \tilde{W}_p^+ , and furthermore is the same as in the proof of Propositions 3 and 4).

So \bar{R}_p^* is equipped with the structure of complex of Lg^A -modules. Its cohomologies are trivial in all dimensions except 0th, where they are equal to \mathbb{C} . Simple calculation by virtue of those of the proof of Propositions 3 and 4 show that the central charge of Lg^A is equal to 0 (due to supersymmetric “cancellation of anomalies”). \square

Now let L_χ be an irreducible representation of Lg^A with dominant integral highest weight χ . Consider the complex $\bar{R}_p^*(\chi) = \bar{R}_p^* \otimes L_\chi$. It is an Lg^A -module with highest weight χ . Hence, the Casimir element Cas of Lg^A [29] acts on $\bar{R}_p^*(\chi)$, commuting with differential d . So $\bar{R}_p^*(\chi)$ decomposes into the direct sum of subcomplexes $\bar{R}_p^*(\chi)_\theta = \bigoplus \bar{R}_p^*(\chi)_\theta$, where

$$\bar{R}_p^*(\chi)_\theta = \{c \in \bar{R}_p^*(\chi) \mid \exists m : (\text{Cas} - \theta)^m c = 0\}.$$

Denote $R_p^*(\chi) = \bar{R}_p^*(\chi)_{(\chi + 2\rho, \chi)}$. Cohomologies of this complex are contained in the 0th dimension and are isomorphic to L_χ .

$R_p^*(\chi)$ is a two-sided BGG resolution corresponding to parabolic subalgebra p of g .

Theorem 3. $R_p^i(\chi) = \bigoplus_{\substack{s \in S_{\text{aff}} \\ lt_p(s) = -i}} W_{s^* \chi, p}$

Proof is equivalent to the standard [26]. It is easy to see that $R^*(\chi)$ is free over a_p^- and the dual to $R^*(\chi)$ is free over a_p^+ . So there is filtration of $R_p^*(\chi)$ by Lg^A -modules, whose adjoint quotients are isomorphic to Wakimoto modules $W_{\chi', p}$. Highest weights χ' of these modules lie on the set $\tilde{\chi} - \sum \gamma_i, \tilde{\chi} \in L_{\chi'}$, where γ_i are different roots of Lg^A . Applying the arguments of [26] we see that the eigenvalue of Cas is equal to $(\chi + 2\rho, \chi)$ only if

$$\chi' = s(\chi) - \sum_{i=1}^{l(s)} \gamma_i = s(\chi + \rho) - \rho = s^* \chi,$$

where $\{\gamma_1, \dots, \gamma_{l(s)}\} = \{\gamma \in \hat{\Delta}_+ : s^{-1}\gamma \in \hat{\Delta}_-\}$. In $R_p^*(\chi)$ the corresponding module $W_{s^* \chi, p}$ lies in the dimension $-lt_p(s)$. After all, the fact that filtration indeed splits follows from homological considerations in spirit of [41].

Our resolutions enable us to compute the semi-infinite cohomology of twisted parabolic subalgebras a_p and $r_p \otimes \mathbb{C}((t))$ with coefficients in the irreducible representation (semi-infinite Borel-Weil-Bott theorem).

Theorem 4. 1. $H^{\infty/2+i}(a_p, L_\chi) = \bigoplus_{\substack{s \in S_{\text{aff}} \\ lt_p(s) = -i}} \mathbb{C}_{s^* \chi}$ as \hat{h} -module,

2. $H^{\infty/2+i}(n_+ \otimes \mathbb{C}((t)), L_\chi) = \bigoplus_{\substack{s \in S_{\text{aff}} \\ lt_{b_+}(s) = -i}} \pi_{(s^* \tilde{\chi}, k + c_g)}$ (as an $L\hat{h}$ -module).

Proof. The spectral sequence corresponding to our resolution degenerates in the first term and gives the result. \square

We use this result for computing the (co)homology of Lie algebra $r_p \otimes \mathbb{C}[[t]]$ of currents on the line to a nilpotent subalgebra.

We will formulate the result only for the algebra $n_+ \otimes \mathbb{C}[[t]]$.

Denote $\bar{S}_{\text{aff}} = A_+ \times S$, where $A_+ = \bigoplus_{i=1}^n \mathbb{Z}_+ \alpha_i$. For $s \in \bar{S}_{\text{aff}}$ $lt_{b_+}(s) = l(s)$.

Theorem 5. $H_i(n_+ \otimes \mathbb{C}[t]) \simeq \bigoplus_{\substack{s \in S_{\text{aff}} \\ l(s)=i}} \mathbb{C}[h_j(m)]_{m=1, \dots, l_j(s)}^{j=1, \dots, n}$, as $h \otimes \mathbb{C}[t]$ module, where

$l_j(s), j=1, \dots, n$ are defined as follows:

$$q - s(q) = \sum_{j=1}^n l_j(s) \alpha_j, \quad l_j(s) \in \mathbb{Z}_+, \quad h_j \text{ are generators of } h.$$

Proof is based on the study of the action of the homologies of $n_+ \otimes \mathbb{C}[t]$ on its semi-infinite homology and follows from Theorem 4. \square

Detailed proof and related results will be published elsewhere.

Note that the standard methods of computing the (co)homologies of current algebras [17] fail in this case. Thus, Theorem 5 is one of the interesting applications of our theory.

7. Examples

7.1. Explicit Formulae for W_χ over Lsl_{n+1}^A

We give the explicit formulae for the action of the generators of Lsl_{n+1}^A which were obtained in [18]. For the simplest case Lsl_2^A they were obtained by Wakimoto in [44] – that is why we call W_χ Wakimoto modules. Note also that formulas for Lsl_2^A and Lsl_3^A were independently obtained by A. B. Zamolodchikov (unpublished).

Denote by $E_i, H_i, F_i, i=1, \dots, n$, standard generators of sl_{n+1} . Denote $a_{ij}(m) = a_{-\alpha_i - \dots - \alpha_j}(m), 1 \leq i \leq j \leq n, a_{ij}^*(m) = a_{-\alpha_i - \dots - \alpha_j}, 1 \leq i \leq j \leq n$. Commutation relations of $a_{ij}(m), a_{ij}^*(m)$ are given in Sect. 5.

Then

$$\begin{aligned} E_i(z) &= a_{ii}^* \left(\sum_{j=1}^{i-1} a_{j,i-1} a_{j,i-1}^* - \sum_{j=1}^i a_{ji} a_{ji}^* \right) : -v a_{ii}^* b_i - \chi_i a_{ii}^* \\ &\quad + \sum_{j=i+1}^n a_{i+1,j} a_{ij}^* - \sum_{j=1}^{i-1} a_{j,i-1} a_{ji}^* + (i+1 - v^2) a_{ii}^*, \\ H_i(z) &= 2 : a_{ii} a_{ii}^* : + \sum_{j=1}^{i-1} (: a_{ji} a_{ji}^* : - : a_{j,i-1} a_{j,i-1}^* :) \\ &\quad + \sum_{j=i+1}^n (: a_{ij} a_{ij}^* - : a_{i+1,j} a_{i+1,j}^* :) + v b_i + \chi_i, \\ F_i(z) &= a_{ii} + \sum_{j=i+1}^n a_{ij} a_{i+1,j}^*, \quad k = v^2 - (n+1), \\ &\quad (a_{ij}, a_{ij}^* \text{ denote } a_{ij}(z), a_{ij}^*(z)) \end{aligned}$$

define the action of Lsl_{n+1}^A in W_χ with $\chi = (\chi_1, \dots, \chi_n, v^2 - (n+1))$, the central charge is equal to $v^2 - (n+1)$. We denote this module by $W_{\bar{\chi}, v}$, where $\bar{\chi} = (\chi_1, \dots, \chi_n)$.

7.2. *Co-twisted Verma Modules over Lsl_2^A* with dominant weights are shown on the picture (Fig. 2). The points on the figures denote singular vectors in ${}^w M_\chi$ or in its quotient, which correspond to singular vectors in the Verma module M_χ . Interrelations between these vectors are expressed by the arrows.




Fig. 2. Twisted Verma modules and Wakimoto module over Lsl_2^A

The chain of arrows leads from one vector to another if and only if for any choice of vectors projecting under some factorizations into singular vectors, the latter vector is generated by the former.

In Fig. 2 the structure of the corresponding Wakimoto module is shown (the description of the structure of Wakimoto modules over Lsl_2^A was given in [19]). We see that this structure is “approximated” by the structures of M_χ^w .

7.3. Structure of Two-Sided BGG Resolutions over Lsl_2^A and Lsl_3^A

The affine Weyl group $S_{\text{aff}}(SL_2)$ is generated by two simple reflections s_0 and s_1 . Let

$$s^{(i)} = \underbrace{s_1 s_0 s_1 \dots s_0}_{-i \text{ times}}, \quad i \leq 0,$$

$$s^{(i)} = \underbrace{s_0 s_1 s_0 \dots s_0}_{i \text{ times}}, \quad i > 0.$$

Then $R_b^i(\chi) = W_{s^{(i)} * \chi}$. We want to give explicit formulae for the differential of $R_b^*(\chi)$.

In [21, 43, 19] composition vertex operators

$$B_{l_1, \dots, l_m}(\beta_1, \dots, \beta_m; \gamma_1, \dots, \gamma_m), \quad l_i \in \mathbb{Z}, \quad \beta_i \in \mathbb{C}, \quad \gamma_i \in \mathbb{C}, \quad \sum_{i=1}^m \gamma_i = 0,$$

were defined. $B_{l_1, \dots, l_m}(\beta_1, \dots, \beta_m; \gamma_1, \dots, \gamma_m)$ acts from $W_{\chi - \sum_{i=1}^m \beta_i v, v}$ to $W_{\chi, v}$.

Let $D_{l_1, \dots, l_m}(\gamma_1, \dots, \gamma_m)$ be the operator: $W_{\chi - 2m, v} \rightarrow W_{\chi, v}$ defined as follows:

$$D_{l_1, \dots, l_m}(\gamma_1, \dots, \gamma_m) = \sum_{i_1, \dots, i_m \in \mathbb{Z}} a(i_1) \dots a(i_m) \times B_{l_1, \dots, l_m} \left(\frac{1}{v}, \dots, \frac{1}{v}; \gamma_1, \dots, \gamma_m \right), \quad \sum_{i=1}^m \gamma_i = 0.$$

In [19] the following theorem is proved.




Fig. 3a, b. Two-sided BGG resolutions over Lsl_2^A and Lsl_3^A . a Lsl_2^A , b Lsl_3^A

Theorem 6. Let $\gamma_i = \frac{1}{v^2}(\bar{\chi} - 2(i-1)) - l_i$, $i = 1, \dots, m$. If $\sum_{i=1}^n \gamma_i = 0$ (Kac-Kazhdan equation [30]), then $D_{l_1, \dots, l_m}(\gamma_1, \dots, \gamma_m)$ is an intertwining operator between $W_{\bar{\chi} - 2m, v}$ and $W_{\bar{\chi}, v}$.

Now let $\chi = (m - 1, l + m - 2)$, $l, m \in \mathbb{Z}$, $l, m > 0$, by the integral dominant highest weight. Then the differential of $R_b^*(\chi)$ $d_i: R_b^{i-1}(\chi) \rightarrow R_b^i(\chi)$ is given by [20],

$$d_i = D_{i, \dots, i}(\gamma_1, \dots, \gamma_m), \quad \gamma_s = \frac{m + 1 - 2s}{m + l}, \quad \text{if } i \text{ is even,}$$

$$d_i = D_{i, \dots, i}(\gamma_1, \dots, \gamma_l), \quad \gamma_s = \frac{l + 1 - 2s}{m + l}, \quad \text{if } i \text{ is odd.}$$

The differential for the two-sided BGG resolution over other algebras may be also expressed via vertex operators. Two-sided BGG resolution over Lsl_2^A does exist for any highest weight $\chi = (\bar{\chi}, k)$ with $\bar{\chi}$ and k rational and $k > -2$ [19, 20].

The structure of the two-sided BGG resolution over Lsl_2^A and Lsl_3^A is shown in Fig. 3.

The points in the figure denote Wakimoto modules. The marked point denotes W_χ . The arrows show the action of the differential. The weight of the module situated in the given point is equal to $s * \chi$, where s is the product of simple reflections along the way from the marked point to the given point. Note that the pictures are composed of BGG resolutions over corresponding finite-dimension Lie algebras sl_2 and sl_3 .

Two-sided BGG resolutions $R_b^*(\chi)$ over Lg^A may be used in WZW models for computations of the correlation functions on the torus and higher genus surfaces in integral representation, in the same way as in Felder’s work [23].

Note that Felder’s resolution over Virasoro algebra [23] is closely connected with our two-sided resolution over Lsl_2^A via the functorial correspondence between Lsl_2^A -modules and modules over Virasoro algebra [20, 47].

Appendix A. Semi-Infinite (Co)homology [16]

Let \mathcal{L} be \mathbb{Z} -graded Lie algebra $\mathcal{L} = \bigoplus_{i \in \mathbb{Z}} \mathcal{L}_i$, $\dim \mathcal{L}_i < \infty$, $n = \mathcal{L}_1 \oplus \mathcal{L}_2 \oplus \dots$, $b = \mathcal{L}_0 \oplus \mathcal{L}_{-1} \oplus \dots$ – its subalgebras. Let M be such an \mathcal{L} -module that the n -submodule generated by any vector of M is finite-dimensional.

Put $\mathcal{L}^* = \bigoplus_{i \in \mathbb{Z}} \mathcal{L}_i^*$, $\mathcal{L}^* = n^\perp \oplus b^\perp$, where \perp denotes the orthogonal complement.

Choose a basis $\varphi_i, i \in \mathbb{Z}$ in \mathcal{L} and dual basis $\varphi_i^*, i \in \mathbb{Z}$ in \mathcal{L}^* . Let $Cl(\mathcal{L})$ be a Clifford algebra with generators $\varphi_i, \varphi_i^*, i \in \mathbb{Z}$ and anticommutation relations: $[\varphi_i, \varphi_j]_+ = [\varphi_i^*, \varphi_j^*]_+ = 0, [\varphi_i, \varphi_j^*]_+ = \delta_{ij}$. Introduce grading on $Cl(\mathcal{L})$, putting $\deg \varphi_i = -1, \deg \varphi_i^* = 1$. Let I be an irreducible representation of $Cl(\mathcal{L})$ with the vacuum vector annihilated by $\varphi_i \in n$ and $\varphi_i^* \in n^\perp$. The module I inherits grading.

Put $\varphi = \sum : \varphi_i \varphi_j^* \varphi_k^* : c_{jk}^i$, where c_{jk}^i are structural constants of \mathcal{L} and $: :$ denotes normal ordering. Define the operator $T: M \otimes I \rightarrow M \otimes I$ as follows:

$$T(m \otimes p) = m \otimes \varphi(p) + \sum \varphi_i(m) \otimes \varphi_i^*(p), \quad m \in M, p \in I.$$

It is easy to see: $T^2 = \text{Id} \otimes \sum b_{ij} \varphi_i^* \varphi_j^*, b_{ij} \in \mathbb{C}$. The expression $\sum b_{ij} \varphi_i^* \varphi_j^*$ determines the 2-form ω on \mathcal{L} , and simple calculations show that ω is cocycle from $H^2(\mathcal{L})$. If $H^2(\mathcal{L}) = 0$, then $w = \delta v$, where $v = \sum r_i \varphi_i^*$ and δ is differential in the cohomological complex of \mathcal{L} . Define the operator $d: M \otimes I \rightarrow M \otimes I$, putting $d = T - v$. Evidently, $d^2 = 0$. We obtain the complex $\{M \otimes I, d\}$, whose i^{th} cohomologies are semi-infinite cohomologies of \mathcal{L} with coefficients in M with respect to the decomposition $\mathcal{L} = n \oplus b$. We denote them $H^{\infty/2+i}(\mathcal{L}, M)$.

Semi-infinite homologies are connected with semi-infinite homologies by the rule: $H_{\infty/2+i}(\mathcal{L}, M) = H^{\infty/2-i}(\mathcal{L}, M)$.

Our considerations may be applied to the finite dimensional Lie algebra \mathcal{L} . The corresponding cohomology groups will be denoted $H_s^i(\mathcal{L}, M)$ and called a shifted cohomology with respect to the decomposition $\mathcal{L} = n \oplus b$.

Appendix B. Modules on the Singular Hyperplane

The singular hyperplane is the hyperplane $k = -c_g$ in $\hat{\mathfrak{h}}^*$. The structure of Lg^A -modules, whose highest weights lie on the singular hyperplane, differ from the structure of other Lg^A -modules. This is caused by the fact that Segal-Sugawara operators [28, 37] of Lg^A commute mutually and with Lg^A if $k = -c_g$. So they yield a great number of singular vectors in the Verma molecule. Let us consider a restricted Verma module $\bar{M}_{(\bar{\chi}, -c_g)} = \bar{M}_{\bar{\chi}}$, which is the quotient of the Verma module $M_{(\bar{\chi}, -c_g)}$ by a submodule, generated by all singular Segal-Sugawara vectors. In [28, 37, 45] it is proved that in general a point of the singular hyperplane (that is if $\bar{\chi}$ does not belong to other Kazhdan-Lusztig hyperplanes [30]) $\bar{M}_{\bar{\chi}}$ is irreducible and hence isomorphic to the restricted Wakimoto module $\bar{W}_{\bar{\chi}}$ defined in 5.1.

In [19] we used explicit formulae for $\bar{W}_{\bar{\chi}}$ for the study of the structure of a restricted Verma module. Here we sketch our main results.

First of all we consider $\bar{W}_{\bar{\chi}}$ and $\bar{M}_{\bar{\chi}}$ over $Lsl_2^A(c_g = 2)$.

Theorem B.1 [19]. *If $\bar{\chi} = 0, 1, 2, \dots$ then $\bar{M}_{\bar{\chi}}$ is isomorphic to $\bar{W}_{\bar{\chi}}$ and contains the unique singular vector of degree $-(\bar{\chi} + 1)\alpha_1$, the quotient by the submodule generated by this singular vector being irreducible.*

If $\bar{\chi} = -2, -3, \dots$, then $\bar{M}_{\bar{\chi}}$ is isomorphic to $\bar{W}_{\bar{\chi}}^$ and contains the unique singular vector of degree $(+\bar{\chi} + 1)\alpha_0$, the quotient by the submodule generated by this singular vector being irreducible.*

If $\bar{\chi} \neq 0, 1, 2$ and $\bar{\chi} \neq -2, -3, \dots$ then $\bar{M}_{\bar{\chi}}$ is irreducible and isomorphic to $\bar{W}_{\bar{\chi}}$.

Note that Malikov has proved this result by other means [38].

For a general affine Kac-Moody algebra Lg^A we have proved the theorem about the structure of $\bar{M}_{\bar{\chi}}$, if $\bar{\chi}$ is projective. We call $\bar{\chi}$ projective if the Verma module $M_{\bar{\chi}}$ over g is projective in the category \mathcal{O} [5]. It means that $M_{\bar{\chi}}$ is not contained in another Verma module over g as a proper submodule. In particular, if $M_{\bar{\chi}}$ is irreducible and is not contained in another Verma module, then $\bar{\chi}$ is projective, and the dominant integral weight is projective.

Theorem B.2 [19]. *If weight $\bar{\chi}$ is projective, then the singular vectors of $\bar{M}_{\bar{\chi}}$ over Lg^A coincide with singular vectors of $M_{\bar{\chi}}$ over g , and also $\bar{M}_{\bar{\chi}}$ is isomorphic to $\bar{W}_{\bar{\chi}}$. In particular, if $M_{\bar{\chi}}$ is irreducible and is not contained in another Verma module then $\bar{M}_{\bar{\chi}}$ and $\bar{W}_{\bar{\chi}}$ are irreducible (and mutually isomorphic).*

We also have the conjecture about non-projective highest weight modules.

Let $\bar{\chi}_1, \dots, \bar{\chi}_l$ be highest weights of Verma modules over g such that $M_{\bar{\chi}_i} \subset M_{\bar{\chi}}$ or $M_{\bar{\chi}} \subset M_{\bar{\chi}_i}$ or there is $\bar{\chi}_j$, such that $M_{\bar{\chi}_i} \subset M_{\bar{\chi}_j}$, $M_{\bar{\chi}} \subset M_{\bar{\chi}_j}$. Let $\bar{\chi}_{\max}$ be the maximal weight among $\bar{\chi}_1, \dots, \bar{\chi}_l, \bar{\chi}$. Then $\bar{\chi}_{\max}$ is the projective weight.

Conjecture B.3 [19]. *The number of singular vectors of $\bar{M}_{\bar{\chi}}$ coincide with the number of singular vectors of $\bar{M}_{\bar{\chi}_{\max}}$. In particular, if $\bar{\chi} = s * \bar{\chi}_0$, where $\bar{\chi}_0$ is the dominant integral weight, then the number of singular vectors in $\bar{M}_{\bar{\chi}}$ is equal to the order of the Weyl group $S(G)$.*

We also conjecture the acyclic resolution consisting of restricted Verma modules. This resolution seems to be the Cousin-Grothendieck resolution of the invertible sheaf on the manifold $\bar{L}(\bar{B})$ with respect to Schubert filtration.

Conjecture B.4 [19]. *There is an acyclic resolution $A^i(\bar{\chi})$ ($\bar{\chi}$ is the dominant integral weight) of Lg^A -modules with central charge $-c_g$, so that $A^i = \bigoplus_{\substack{lt(s) = -i \\ s \in S_{\text{aff}}}} \bar{M}_{s(\bar{\chi} + \varrho) - \varrho}$.*

Acknowledgements. We would like to thank A. A. Beilinson and D. B. Fuchs for helpful discussions.

References

1. Beilinson, A.: Localization of representations of reductive Lie algebras. In: Proceedings of ICM, Warsaw 1984, Vol. 2, pp. 699–710
2. Beilinson, A., Bernstein, J.: Localization de g -modules. C.R. Acad. Sci. Paris **192**, 15–18 (1981)
3. Beilinson, A., Bernstein, J.: Generalization of a theorem of Casselman. Report on Utah Conference on representation theory, April 1982. In: Representation theory of reductive groups. Trombi, P. (ed.) DM 40. Boston: Birkhäuser 1983
4. Bernstein, J.: Lectures on D -modules (unpublished)
5. Bernstein, J., Gelfand, I., Gelfand, S.: On the category of g -modules. Funkt. Anal. Prilozhen. **10**, 1–8 (1976) (in Russian)
6. Bernstein, J., Gelfand, I., Gelfand, S.: Schubert cells and cohomology of spaces G/P . Usp. Mat. Nauk (= Russ. Math. Surv.) **28**, 3–26 (1973) (in Russian)
7. Bernstein, J., Gelfand, I., Gelfand, S.: Differential operators on the base affine space and a study of g -modules. In: Lie groups and their representations, pp. 21–64. New York: Wiley 1975
8. Borel, A.: Kählerian coset spaces of semi-simple Lie groups. Proc. Natl. Acad. USA **40**, 1147–1151 (1954)

9. Bott, R.: Homogeneous vector bundles. *Ann. Math.* **66**, 203–248 (1957)
10. Brylinsky, J.L., Kashiwara, M.: Kazhdan-Lusztig conjecture and holonomic systems. *Invent. Math.* **64**, 487–410 (1981)
11. Conley, C.C., Zehnder, E.: Morse type index theorem for flows and periodic solutions for hamiltonian equation. *Commun. Pure and Appl. Math.* **34**, 207–253 (1984)
12. Date, E., Jimbo, M., Kashiwara, M., Miwa, T.: Transformation groups for soliton equations. In: *Proceedings of RIMS symposium*. Jimbo, M., Miwa, T. (eds), pp. 39–120. Singapore: World Scientific 1983
13. Deodhar, V., Gabber, O., Kac, V.: Structure of some categories of representations of infinite-dimensional Lie algebras. *Adv. Math.* **45**, 92–116 (1982)
14. Dotsenko, V.I.S., Fateev, V.A.: Conformal algebra and multipoint correlation functions in 2D statistical models. *Nucl. Phys. B* **240**, F 312, 312–348 (1984)
15. Dotsenko, V.I.S., Fateev, V.A.: Private communications
16. Feigin, B.L.: Semi-infinite homology of Kac-Moody and Virasoro Lie algebras. *Usp. Mat. Nauk* (= *Russ. Math. Surv.*) **39**, 195–196 (1984) (in Russian)
17. Feigin, B.L.: On the cohomology of the Lie algebra of vector fields and of the current algebra. *Sel. Math. Sov.* **7**, 49–62 (1988)
18. Feigin, B.L., Frenkel, E.V.: The family of representations of affine Lie algebras. *Usp. Math. Nauk* (= *Russ. Math. Surv.*) **43**, 227–228 (1988) (in Russian)
19. Feigin, B.L., Frenkel, E.V.: Representations of affine Kac-Moody algebras and bosonization. To be published in: V. G. Knizhnik Memorial volume. Singapore: World Scientific 1989
20. Feigin, B.L., Frenkel, E.V.: Representations of affine Kac-Moody algebras, bosonization and resolutions. To be published in *Lett. Math. Phys.* **19**, N3 (1990)
21. Feigin, B.L., Fuchs, D.B.: Representations of the Virasoro algebra. To be published in: *Representations of infinite-dimensional Lie groups and Lie algebras*. New York: Gordon and Breach 1989
22. Feigin, B.L., Tsygan, B.L.: Cohomology of generalized Jakobean matrices. *Funct. Anal. Prilozhen.* **17**, 86–87 (1983)
23. Felder, G.: BRST approach to minimal models. *Nucl. Phys. B* **317**, 215–236 (1989)
24. Floer, A.: Proof of Arnold conjecture for surfaces and generalization to certain Kähler manifolds. *Duke Math. J.* **53**, 1–32 (1986)
25. Fuchs, D.B.: *Cohomology of infinite-dimensional Lie algebras*. New York: Plenum Press 1986
26. Garland, H., Lepowsky, J.: Lie algebra homology and the Macdonald-Kac formulas. *Invent. Math.* **34**, 37–76 (1976)
27. Gerasimov, A., Marshakov, A., Morosov, A., Olshansky, M., Shatashvili, S.: Moscow preprint, 1989
28. Hayashi, T.: Sugawara construction and Kac-Kazhdan conjecture. *Invent. Math.* **94**, 13–52 (1988)
29. Kac, V.G.: *Infinite-dimensional Lie algebras*. Boston, MA: Birkhäuser 1983
30. Kac, V.G., Kazhdan, D.: Structure of representations with highest weight of infinite-dimensional Lie algebras. *Adv. Math.* **34**, 97–108 (1979)
31. Kac, V., Peterson, D.: Unpublished, Kac, V.G.: Constructing groups associated to infinite-dimensional Lie algebras. In *MSRI Publ.*, 4. Berlin, Heidelberg, New York: Springer 1985
- 31a. Kac, V., Peterson, D.: Spin and wedge representations of infinite-dimensional Lie algebras and groups. *Proc. Natl. Acad. Sci. USA* **78**, 3308–3312 (1981)
32. Kac, V., Todorov, I.: Superconformal current algebras and their unitary representations. *Commun. Math. Phys.* **102**, 337–347 (1985)
33. Kazhdan, D., Lusztig, G.: Representation of Coxeter groups and Hecke algebras. *Invent. Math.* **53**, 165–184 (1979)
34. Kazhdan, D., Lusztig, G.: Schubert varieties and Poincaré duality. *Proc. Symp. Pure Math.* **36**, 185–203 (1980)
35. Kempf, G.: Cousin-Grothendieck complex of induced representations. *Adv. Math.* **29**, 310–396 (1978)

36. Kostant, B., Kumar, S.: The nil Hecke ring and cohomology of G/P for a Kac-Moody group. *Adv. Math.* **62**, 187–237 (1986)
37. Malikov, F.: Singular vectors in Verma modules over affine algebras. *Funct. Anal. Prilozhen.* **23**, 76–77 (1989)
38. Malikov, F.: Verma modules over affine rank 2 algebras. To appear in *Algebra and Analysis* (1989) (in Russian)
39. Peterson, D., Kac, V.: Infinite flag varieties and conjugacy theorem. *Proc. Natl. Acad. USA* **80**, 1778–1782 (1983)
40. Pressley, A., Segal, G.: *Loop groups*. Oxford: Clarendon Press 1987
41. Rocha-Caridi, A., Wallach, N.: Projective modules over graded Lie algebras. *Math. Z.* **180**, 151–177 (1982)
42. Tate, J.: Residues of differentials on curves. *Ann. Sci. E.N.S.* **4** Ser., t. 1, 149–159 (1968)
43. Tsuchiya, A., Kanie, Y.: Fock space representations of the Virasoro algebra – Intertwining operators. *Publ. RIMS, Kyoto Univ.* **22**, 259–327 (1986)
44. Wakimoto, M.: Fock representations of affine Lie algebra $A_1^{(1)}$. *Commun. Math. Phys.* **104**, 604–609 (1986)
45. Wallach, N.: A Class of non-standard modules for affine Lie algebras. *Math. Z.* **196**, 303–313 (1987)
46. Zamolodchikov, A.: Unpublished
47. Bershadsky, M., Ooguri, H.: Hidden $sl(n)$ symmetry in conformal field theory, IASSNS HEP-89/09
48. Bernard, D., Felder, G.: Fock representations and BRST cohomology of $sl(2)$ current algebra, ETH-TH/89-26
49. Bouwknegt, P., McCarthy, J., Pilch, K.: MIT preprint CTP #1797

Communicated by L. Alvarez-Gaumé

Received June 6, 1989

Note added in proof. After this paper had been accepted for publication, we received preprint [48], where a two-sided BGG resolution (BRST complex) was constructed for Lsl_2^A . The formulas obtained in [48] coincide with ours from 7.3. In [49] certain explicit formulas for the differential of two-sided BGG resolution (BRST operator) were given using vertex operators introduced in [19]. We would like to thank P. Bouwknegt and J. McCarthy for interesting discussions.

