

## SOME INEQUALITIES FOR $t$ -DESIGNS

RYUZABURO NODA

(Received March 12, 1975)

### 1. Introduction

D.K. Ray-Chaudhuri and R.M. Wilson [3] proved  $b \geq \binom{v}{s}$  for  $2s$ -designs with  $v \geq k+s$ , generalizing Fischer's inequality  $b \geq v$  for 2-designs, and Petrenjuk's inequality  $b \geq \binom{v}{2}$  for 4-designs. In this note we introduce a notion of rank  $s$  tactical decompositions of  $2s$ -designs, and generalize some of well known results for 2-designs.

**DEFINITION.** A rank  $s$  tactical decomposition of a  $2s$ -design  $(X, \mathcal{B})$  is a partition of the set  $X^{(s)}$  of all  $s$ -element subsets of  $X$  into  $s$ -point classes  $X_1, X_2, \dots, X_m$ , together with a partition of  $\mathcal{B}$  into block classes  $\mathcal{B}_1, \mathcal{B}_2, \dots, \mathcal{B}_{m'}$ , such that the number of elements of  $X_i$  contained in a block  $B$  of  $\mathcal{B}_j$  depends only on  $i$  and  $j$ , (and does not depend on the choice of  $B$  in  $\mathcal{B}_j$ ) and the number of blocks in  $\mathcal{B}_h$  containing an element  $\{p_1, p_2 \dots p_s\}$  of  $X_g$  depends only on  $h$  and  $g$ .

Our first result is:

**Theorem 1.** Let a  $2s$ - $(v, k, \lambda)$  design  $(X, \mathcal{B})$  with  $v \geq k+s$  admit a rank  $s$  tactical decomposition with  $m$   $s$ -point classes and  $m'$  block classes. Then  $m \leq m'$ .

The case  $s=1$  in the above was proved by W.M. Kantor (Theorem 4.1 [2]). Our proof, which will be given in section 2, seems to be more elementary.

If  $G$  is a group of automorphisms of a  $2s$ -design  $(X, \mathcal{B})$ , then the orbits of  $G$  on  $X^{(s)}$ , together with the orbits of  $G$  on  $\mathcal{B}$ , form a rank  $s$  tactical decomposition of  $(X, \mathcal{B})$ . Therefore, by Theorem 1, we have

**Corollary 2.** A group of automorphisms of a  $2s$ - $(v, k, \lambda)$  design  $(X, \mathcal{B})$  with  $v \geq k+s$  has at least as many orbits on  $\mathcal{B}$  as on  $X^{(s)}$ . In particular a block transitive automorphism group of  $(X, \mathcal{B})$  is  $s$ -homogeneous on points.

The following is a slight extension of a theorem of D.K. Ray-Chaudhuri and R.M. Wilson (Theorem 2 [3]).

**Theorem 3.** Let a  $2s$ - $(v, \mathcal{B}, \lambda)$  design  $(X, \mathcal{B})$  with  $v \geq k+s$  admit a rank

$s$  tactical decomposition with  $m$   $s$ -point classes and  $m'$  block classes. Then  $b$   
 $(=|\mathcal{B}|) \geq \binom{v}{s} + m' - m.$

Extending the notion of parallelisms of 2-designs we also introduce the following.

**DEFINITION.** A rank  $s$  parallelism of a  $2s$ -design  $(X, \mathcal{B})$  is an equivalence relation on  $\mathcal{B}$  with the property that each element of  $X^{(s)}$  lies in a unique member of each equivalent class. Equivalently a rank  $s$  parallelisms of a  $2s$ -design  $(X, \mathcal{B})$  is a partition of  $\mathcal{B}$  into "rank  $s$  parallel classes", each of which is a partition of  $X^{(s)}$ .

It is easy to see that a rank  $s$  parallelism of a  $2s$ -design is a rank  $s$  tactical decomposition with one  $s$ -point class and  $\lambda_s$  block classes, each of which consists of  $\binom{v}{s} / \binom{k}{s}$  blocks. Here, as usual,  $\lambda_s$  denotes the number of blocks containing given  $s$  points. Thus, by Theorem 3 (or by Theorem 2 of [3]), we have

**Corollary 4.** Let a  $2s$ - $(v, k, \lambda)$  design with  $v \geq k + s$  have a rank  $s$  parallelism. Then  $b \geq \binom{v}{s} + \lambda_s - 1.$

Corollary 4 is a generalization of Bose's inequality  $b \geq v + r - 1$  for 2-designs with a parallelism [1]. The author does not know whether there exist  $2s$ -designs,  $s \geq 2$ , with a rank  $s$  parallelism. But the following is true.

**Theorem 5.** If  $s \geq 2$ , there exist no  $2s$ - $(v, k, \lambda)$  designs,  $v \geq k + s$ , with a rank  $s$  parallelism having the smallest number  $b = \binom{v}{s} + \lambda_s - 1$  of blocks.

In the case  $s = 1$ , as is well known, there exist infinitely many 2-designs with a parallelism having  $v + \lambda_1 - 1$  blocks.

**2. Proof of Theorem 1**

Let  $N$  be the  $\binom{v}{s} \times b$ -matrix whose rows are numbered by elements of  $X^{(s)}$  and columns by elements of  $\mathcal{B}$ , and whose  $(\{i_1, i_2, \dots, i_s\}, B)$  entry is 1 or 0 according as  $\{i_1, i_2, \dots, i_s\} \subset B$  or not. Then  $N$  has rank  $\binom{v}{s}$  by (the proof of) Theorem 1 [3]. So our Theorem 1 is an immediate consequence of the following.

**Lemma.** Let  $M$  be a (real)  $n \times b$ -matrix with rank  $n$ . Assume that  $M$  can be decomposed into  $mm'$  rectangular submatrices  $M_{ij}$ ,  $1 \leq i \leq m$ ,  $1 \leq j \leq m'$ , such that  $M_{ij}$  is an  $n_i \times b_j$ -matrix with constant column sum  $k_{ij}$ . Then  $m \leq m'$ .

Proof. Let  $x_i$  ( $1 \leq i \leq n$ ) denote the  $i$ -th row vector of  $M$ . Set  $y_1 = x_1 + \dots + x_{n_1}$ ,  $y_2 = x_{n_1+1} + \dots + x_{n_1+n_2}$ ,  $\dots$ ,  $y_m = x_{n_1+n_2+\dots+n_{m-1}+1} + \dots + x_{n_1+n_2+\dots+n_m}$ .

Then  $y_i$  is the vector of the form:

$$y_i = (\overbrace{k_{i_1} \dots k_{i_1}}^{b_1}, \overbrace{k_{i_2} \dots k_{i_2}}^{b_2}, \dots, \overbrace{k_{i_{m'}} \dots k_{i_{m'}}}^{b_{m'}}), \quad 1 \leq i \leq m_i.$$

Then since the  $m$  vectors  $y_i$  are linearly independent, it follows that  $m \leq m'$ .

**3. Proof of Theorem 3**

We make use of an argument of D.K. Ray-Chaudhuri and R.M. Wilson [3]. Let  $V_s$  denote the free vector space over the rationals generated by  $X^{(s)}$ . Clearly  $V_s$  is  $\binom{v}{s}$  dimensional over rationals. Now for each  $A \in \mathcal{B}$ , define a vector  $\hat{A} \in V_s$  as the sum of all  $s$ -subsets of  $A$ , *i.e.*

$$\hat{A} = \sum(S: S \in X^{(s)}, S \subseteq A).$$

D.K. Ray-Chaudhuri and R.M. Wilson showed that the vectors  $\{\hat{A}: A \in \mathcal{B}\}$  span  $V_s$ . Put  $\hat{X}_j = \sum(S: S \in X_j)$ . Then, by our assumption

$$\sum\{\hat{A}: A \in \mathcal{B}_i\} = \sum_{j=1}^m \lambda_{ij} \sum(S: S \in X_j) = \sum_{j=1}^m \lambda_{ij} \hat{X}_j, \quad 1 \leq i \leq m'$$

So, if we choose one block  $A_i$  from each  $\mathcal{B}_i$ , then

$$\{\hat{A}: A \in \mathcal{B} - \{A_1, \dots, A_{m'}\}\} \cup \{\hat{X}_1, \hat{X}_2, \dots, \hat{X}_m\}$$

spans  $V_s$ . The stated inequality follows.

**4. Proof of Theorem 5**

Assume by way of contradiction that there exists a  $2s$ - $(v, k, \lambda)$  design  $(X, \mathcal{B})$ ,  $s \geq 2$ ,  $v \geq k + s$  with a rank  $s$  parallelism having the smallest number  $b = \binom{v}{s} + \lambda_s - 1$  of blocks. Then we have

$$\begin{aligned} \frac{\binom{v}{s}}{\binom{k}{s}} \lambda_s &= \binom{v}{s} + \lambda_s - 1, \\ \lambda_s \left\{ \frac{\binom{v}{s}}{\binom{k}{s}} - 1 \right\} &= \binom{v}{s} - 1 \end{aligned} \tag{4.1}$$

Case 1.  $s=2r$  is even.

Applying the theorem of D.K. Ray-Chaudhuri and R.M. Wilson (Theorem 1 [3]), to a contracted  $s$ - $(v-s, k-s, \lambda)$  design of  $(X, \mathcal{B})$ , we have

$$\lambda_s \geq \binom{v-s}{r} \quad (4.2)$$

Then (4.1) and (4.2) yield

$$\binom{v-s}{r} \left\{ \frac{\binom{v}{s}}{\binom{k}{s}} - 1 \right\} \leq \binom{v}{s} - 1, \quad (4.3)$$

Now let  $\mathcal{B}_1$  be a rank  $s$  parallel class of  $(X, \mathcal{B})$ . Then  $(X, \mathcal{B}_1)$  is a  $s$ - $(v, k, 1)$  design, and hence, again by the theorem of D.K. Ray-Chaudhuri and R.M. Wilson, we have

$$\frac{\binom{v}{s}}{\binom{k}{s}} \geq \binom{v}{r} \quad (4.4)$$

Then (4.3) and (4.4) imply

$$\begin{aligned} \binom{v-s}{r} \left\{ \binom{v}{r} - 1 \right\} &\leq \binom{v}{s} - 1 \\ \left\{ \binom{v-s}{r} - 1 \right\} \binom{v}{r} &\leq \binom{v-s}{r} \left\{ \binom{v}{r} - 1 \right\} \leq \binom{v}{s} \\ \binom{v-s}{r} - 1 &\leq \frac{\binom{v}{s}}{\binom{v}{r}} = \frac{(v-r)(v-r-1)\cdots(v-2r+1)}{2r(2r-1)\cdots(r+1)} \end{aligned} \quad (4.5)$$

On the other hand, since  $v \geq 6r$ , we have

$$\frac{v-s-i}{r-i} \geq \frac{v-r-i}{2r-i}, \quad 0 \leq i \leq r-2 \quad (4.6)$$

and

$$v-3r+1 \geq \frac{v-2r+1}{r+1} + 1. \quad (4.7)$$

Then (4.6) and (4.7) yield

$$\begin{aligned} \binom{v-s}{r} - 1 &= \frac{(v-s)}{r} \cdot \frac{(v-s-1)}{r-1} \cdots \frac{(v-3r+2)}{2} \cdot (v-3r+1) - 1 \\ &\geq \frac{(v-r)}{2r} \cdot \frac{(v-r-1)}{2r-1} \cdots \frac{(v-2r+2)}{r+2} \left\{ \frac{v-2r+1}{r+1} + 1 \right\} - 1 \end{aligned}$$

$$\begin{aligned}
 &= \frac{\binom{v}{s}}{\binom{v}{r}} + \frac{(v-r)\binom{v}{s-1}}{s\binom{v}{r+1}} - 1 \\
 &\cong \frac{\binom{v}{s}}{\binom{v}{r}}.
 \end{aligned}$$

This contradicts (4.5).

Case 2.  $s=2r+1$  ( $r \geq 1$ ) is odd.

Applying the theorem of D.K. Ray-Chaudhuri and R.M. Wilson to a contracted  $2(s-1)-(v-2, k-2, \lambda)$  design of  $(X, \mathcal{B})$ , we have

$$\lambda_2 = \frac{\binom{v-2}{s-2}}{\binom{k-2}{s-2}} \lambda_s \cong \binom{v-2}{s-1} \tag{4.8}$$

Then (4.1) and (4.8) yield

$$\begin{aligned}
 &\frac{\binom{k-2}{s-2}(v-s)}{s-1} \left\{ \frac{\binom{v}{s}}{\binom{k}{s}} - 1 \right\} \cong \binom{v}{s} - 1 \\
 &\frac{s(v-s)}{k(k-1)} \left\{ \binom{v}{s} - \binom{k}{s} \right\} \cong \binom{v}{s} - 1 \tag{4.9}
 \end{aligned}$$

On the other hand applying Fischer's inequality to a contracted  $2-(v-s+2, k-s+2, 1)$  design of a  $s-(v, k, 1)$  design  $(X, \mathcal{B}_1)$ , where  $\mathcal{B}_1$  is a rank  $s$  parallel class of  $(X, \mathcal{B})$ , we have

$$(k-s+2)(k-s+1) \leq v-s+1. \tag{4.10}$$

We shall now show that

$$\binom{v}{s} - \binom{k}{s} \cong \frac{4}{5} \left\{ \binom{v}{s} - 1 \right\} \tag{4.11}$$

Deny (4.11). Then

$$\begin{aligned}
 &\frac{1}{5} \binom{v}{s} \cong \binom{k}{s} \\
 &v(v-1) \cdots (v-s+1) \cong k(k-1) \cdots (k-s+1) 5 \tag{4.12}
 \end{aligned}$$

Then (4.10) and (4.12) give

$$\begin{aligned} v(v-1) \cdots (v-s+2) &\cong k(k-1) \cdots (k-s+3) 5 \\ v(v-1) \cdots (v-s+3) &\cong k(k-1) \cdots (k-s+3) \\ v &\cong k, \text{ a contradiction.} \end{aligned}$$

Now by (4.9) and (4.11) we obtain

$$s(v-s) \leq \frac{5}{4} k(k-1)$$

Combining this with (4.10) gives

$$(k-s+2)(k-s+1) \leq \frac{5k(k-1)}{4s} + 1 \quad (4.13)$$

Then since  $k \geq 2s$  (4.13) implies

$$\left(\frac{1}{2}k+2\right)\left(\frac{1}{2}k+1\right) \geq \frac{5k(k-1)}{4s} + 1 \quad (4.14)$$

If  $s \leq 5$  then (4.14) gives

$$\begin{aligned} \left(\frac{1}{2}k+2\right)\left(\frac{1}{2}k+1\right) &\leq \frac{k(k-1)}{4} + 1 \\ \frac{3}{2}k+2 &\leq -\frac{1}{4}k+1, \text{ a contradiction.} \end{aligned}$$

So we must have  $s=3$ . But then (4.13) gives

$$\begin{aligned} (k-1)(k-2) &\leq \frac{5}{12}k(k-1)+1 \\ 7k^2-31k+12 &\leq 0 \\ k &\leq 4, \text{ a contradiction.} \end{aligned}$$

OSAKA UNIVERSITY

---

### References

- [1] R.C. Bose: *A note on the resolvability of balanced incomplete block designs*, *Sankhya* **6** (1942), 105-100.
- [2] W.M. Kantor: *Automorphism groups of designs*, *Math. Z.* **109** (1969), 246-252.
- [3] D.K. Ray-Chaudhuri and R.M. Wilson: *On  $t$ -designs*, *Osaka. J. Math.* **12** (1975), 737-744.