

BOUNDS FOR THE PROBABILITY OF A UNION, WITH APPLICATIONS¹

BY EUSTRATIOS G. KOUNIAS

The University of Connecticut

1. Inequalities. Consider a probability space $(\Omega, \mathfrak{F}, P)$ and the sequence of events $A_i \in \mathfrak{F}, i = 1, 2, \dots, n$. Let $I_i(\omega)$ be the indicator random variable of the event $A_i, i = 1, 2, \dots, n$, then $\max_{i=1, \dots, n} I_i(\omega)$ is the indicator random variable of the event $\bigcup_{i=1}^n A_i$.

In the present paper we assume that we only know $P_i = P(A_i)$ and $P_{i,j} = P(A_i \cap A_j)$ for all $i, j = 1, 2, \dots, n$, and we find bounds for the $P(\bigcup_{i=1}^n A_i)$.

It is not difficult to prove that the inequality

$$(1) \quad \sum_{i \in J} I_i(\omega) - \sum_{i < j, i, j \in J} I_i(\omega) I_j(\omega) \leq \max_{i \in J} I_i(\omega) \leq (1 - I_k(\omega)) \sum_{i \in J} I_i(\omega) + I_k(\omega)$$

holds for all points $\omega \in \Omega$, all subsets J of the set $\{1, 2, \dots, n\}$ and all $k \in J$.

Taking expectations in (1) we obtain

$$(2) \quad \sum_{i \in J} P_i - \sum_{i < j, i, j \in J} P_{i,j} \leq P(\bigcup_{i \in J} A_i) \leq \sum_{i \in J} P_i - \sum_{i \in J, i \neq k} P_{k,i}$$

Hence, the best lower bound for the $P(\bigcup_{i=1}^n A_i)$, among the lower bounds (2), is

$$(3) \quad \max_J (\sum_{i \in J} P_i - \sum_{i < j, i, j \in J} P_{i,j})$$

Now if we take disjoint subsets J_1, \dots, J_m such that $\bigcup_{i=1}^m J_i = \{1, \dots, n\}$ and denote by $B_r = \bigcup_{i \in J_r} A_i$, then $\bigcup_{r=1}^m B_r = \bigcup_{i=1}^n A_i$ and using (2) obtain $P(B_r) \leq \min(1, \sum_{i \in J_r} P_i - \max_{k \in J_r} \sum_{i \in J_r, i \neq k} P_{k,i}) = T_r$ and $P(\bigcup_{i=1}^n A_i) = P(\bigcup_{r=1}^m B_r) \leq \sum_{r=1}^m T_r$. In particular for $m = 1$ we obtain the upper bound

$$(4) \quad \min [\sum_{i=1}^n P_i - \max_{k=1, 2, \dots, n} \sum_{i=1, i \neq k}^n P_{k,i}, 1]$$

Actually, it is easy to construct examples where the bounds (3) and (4) are attained.

Below we derive another lower bound which is not as sharp but more elegant than (3). For this, let us use the following notation:

$$P' = (P_1, \dots, P_n), \quad Q = \{P_{i,j}\}, \quad I'(\omega) = (I_1(\omega), \dots, I_n(\omega)),$$

i.e., $P, I(\omega)$ are $n \times 1$ vectors and Q is $n \times n$ matrix. Let us also denote by Q^- the generalized inverse of Q , i.e., satisfying $QQ^-Q = Q$. Such an inverse always exists, (see [4], page 24). The lower bound (5), we derive below, for non-singular Q was obtained by Gallot [3].

LEMMA 1.1. *The vector P is in the range of Q .*

PROOF. If Q is non-singular, the lemma is obvious. If Q is singular, then for

Received 5 September 1967; revised 25 July 1968.

¹ This research was supported in part by the University of Connecticut Research Foundation, Grant No. 5.171-35-010.



every vector a in the null space of Q , i.e., $Qa = 0$, we have

$$E(a'I)^2 = a'Qa = 0.$$

Hence, $a'I = 0$ with probability one and, taking expectations, we obtain $a'P = 0$, i.e., P is orthogonal to the null space of Q and thus lies in the range of Q . Q.E.D.

THEOREM 1.1.

$$(5) \quad P(\mathbf{U}_{i=1}^n A_i) \geq P'Q^-P.$$

PROOF. For any vector $a' = (a_1, \dots, a_n)$, the following inequality

$$(6) \quad (a'I(\omega))^2 - 2(a'I(\omega)) + \max_{i=1, \dots, n} I_i(\omega) \geq 0$$

holds for all points $\omega \in \Omega$; hence, taking expectations, we obtain

$$(7) \quad P(\mathbf{U}_{i=1}^n A_i) \geq 2a'P - a'Qa.$$

The vector a which maximizes $2a'P - a'Qa$ satisfies the relation $Qa = P$. Therefore,

$$P(\mathbf{U}_{i=1}^n A_i) \geq a'Qa \quad \text{subject to} \quad Qa = P,$$

but

$$a'Qa = a'QQ^-Qa = P'Q^-P. \quad \text{Q.E.D.}$$

2. Related work. Recently, Dawson and Sankoff [2] have proved a result equivalent to

$$(8) \quad P(\mathbf{U}_{i=1}^n A_i) \geq 2(B + C)/(2 + \rho) - 2C/(1 + \rho)$$

where

$$B = \sum_{i=1}^n P_i, \quad C = \sum_{i=1}^n \sum_{i < j} P_{ij}, \quad \text{and} \quad \rho = [2C/B].$$

They show that (8) is stronger than

$$(9) \quad P(\mathbf{U}_{i=1}^n A_i) \geq B^2/(2C + B)$$

which was derived by Chung and Erdős [1]. Nevertheless, inequality (8) is not better than (5) as will be clear from the examples below.

In the following numerical examples, inequalities (3), (4), (5), (8), and (9) are compared.

(1) For $n = 3$ and

$$\begin{aligned} P_1 &= 0.5, & P_2 &= 0.4, & P_3 &= 0.3, \\ P_{12} &= 0, & P_{13} &= 0.25, & P_{23} &= 0.05, \end{aligned}$$

the lower bounds (3), (5), (8), and (9) are 0.9, 0.9, 0.9, and 0.8, respectively. The upper bound (4) is 0.9.

(2) For $n = 3$ and

$$P_1 = 0.50, \quad P_2 = 0.40, \quad P_3 = 0.60,$$

$$P_{12} = 0.20, \quad P_{13} = 0.30, \quad P_{23} = 0.10,$$

the lower bounds (3), (5), (8), and (9) are 0.9, 0.9, 0.858 and 0.833 respectively. The upper bound (4) is 1.

(3) For $n = 3$ and

$$P_1 = \frac{1}{3}, \quad P_2 = \frac{2}{3}, \quad P_3 = \frac{1}{3},$$

$$P_{12} = \frac{1}{6}, \quad P_{13} = \frac{1}{3}, \quad P_{23} = \frac{1}{6},$$

the lower bounds (3), (5), (8), and (9) are 0.83, 0.76, 0.66, and 0.66, respectively. The upper bound (4) equals 0.83.

Notice that inequality (8) is equivalent to (5) in the first example, better in the second, and worse in the last example in which Q is singular. For $n = 2$, (8) gives the exact result i.e., equality holds, whereas equality holds in (5) only if Q is singular or $P_1 > P_2 = P_{12}$.

3. Applications. Assume now that we have an infinite sequence of events $A_i \in \mathcal{F}$, $i = 1, 2, \dots$, and denote by $D'_{r,n} = (P_r, \dots, P_{r+n})$, $Q_{r,n} = \{P_{i,j}\}$, $i, j = r, r + 1, \dots, r + n$, and the quadratic form $D'_{r,n} Q_{r,n}^- D_{r,n} = c_{r,n}$.

LEMMA 3.1. *The $\lim_{r \rightarrow \infty} \lim_{n \rightarrow \infty} c_{r,n} = c$ exists and lies between zero and one.*

PROOF. By the definition of $c_{r,n}$ and relation (7), we obtain

$$1 \geq c_{r,n+1} = \max_a [2a'D_{r,n+1} - a'Q_{r,n+1}a]$$

$$\geq \max_b [2b'D_{r,n+1} - b'Q_{r,n+1}b] = c_{r,n} \geq 0$$

where $a' = (a_0, \dots, a_{n+1})$ and $b' = (b_0, \dots, b_n, 0)$.

Thus, for any r the sequence $c_{r,n}$, $n = 1, 2, \dots$, is increasing and bounded by 1; hence,

$$\lim_{n \rightarrow \infty} c_{r,n} = c_r.$$

Similarly, we prove that $0 \leq c_{r,n} \leq c_{r-1,n+1} \leq 1$ so that $1 \geq c_{r-1} \geq c_r \geq 0$ and the sequence c_r is decreasing and bounded below by 1; hence, the result. Q.E.D.

THEOREM 3.1.

$$\liminf_{r \rightarrow \infty} \liminf_{n \rightarrow \infty} \left(\sum_{i=r}^n P_i - \max_{k=r, \dots, n} \sum_{i=r, i \neq k}^n P_{k,i} \right)$$

$$\geq P(\bigcap_{r=1}^{\infty} \bigcup_{n=r}^{\infty} A_n) \geq \lim_{r \rightarrow \infty} \lim_{n \rightarrow \infty} c_{r,n}.$$

PROOF.

$$P(\bigcap_{r=1}^{\infty} \bigcup_{n=r}^{\infty} A_n) = \lim_{r \rightarrow \infty} P(\bigcup_{n=r}^{\infty} A_n).$$

But, by virtue of (4) and (5) we have

$$\liminf_{n \rightarrow \infty} \left(\sum_{i=r}^n P_i - \max_{k=r, \dots, n} \sum_{i=r, i \neq k}^n P_{k,i} \right)$$

$$\geq P(\bigcup_{n=r}^{\infty} A_n) \geq \lim_{n \rightarrow \infty} c_{r,n} = c_r$$

and hence the result. Q.E.D.

COROLLARY 3.1. *The lower bound of Theorem 3.1 is better than the lower bound of Erdős-Rényi ([5], p. 326), i.e.,*

$$(10) \quad \lim_{r \rightarrow \infty} \lim_{n \rightarrow \infty} c_{r,n} \geq \limsup_{n \rightarrow \infty} [(\sum_{i=1}^n P_i)^2 / \sum_{i=1}^n \sum_{j=1}^n P_{ij}].$$

PROOF. First, observe that

$$c_{r,n} = D'_{r,n} Q_{r,n}^- D_{r,n} \\ = \max_a (2a' D_{r,n} - a' Q_{r,n} a) \geq (\sum_{i=r}^{n+r} P_i)^2 / \sum_{i=r}^{n+r} \sum_{j=r}^{n+r} P_{ij}$$

because the last expression is found by setting the $n \times 1$ vector $a' = c(1, 1, \dots, 1)$ with

$$c = \sum_{i=r}^{n+r} P_i / \sum_{i=r}^{n+r} \sum_{j=r}^{n+r} P_{ij}.$$

This shows that (5) is stronger than (9) which in consequence proves our claim.

Q.E.D.

The following example, although not very interesting, shows that strict inequality might hold in (10) and hence the Borel-Cantelli type lemma obtainable from Theorem 3.1 is stronger than the one obtained by Erdős and Rényi. Let the events $A_i, i = 1, 2, \dots$, such that,

$$P_{2i-1} = p, \quad P_{2i} = q, \quad \text{with } 0 < q < p \leq 1, \quad \text{and} \\ P_{i,j} = p, \quad \text{if } i \text{ and } j \text{ are even,} \\ P_{i,j} = q, \quad \text{if } i \text{ and } j \text{ are not both even, for all } i, j, = 1, \dots,$$

then we easily see that $c_r = p$ for all $r = 1, \dots$, so that $\lim_{r \rightarrow \infty} c_r = p$, whereas $\lim_{n \rightarrow \infty} (\sum_{i=1}^n P_i)^2 / (\sum_{i=1}^n \sum_{j=1}^n P_{i,j}) = (p + q)^2 / (p + 3q)$ which is less than p .

As the last application, consider the sequence of random variables $X_n, n = 1, \dots$, and let A_n be the event $|X_n| > e$ for some $e > 0$, then

$$P(\sup_{n \geq r} |X_n| > e) = P(\bigcup_{n=r}^{\infty} A_n)$$

and

$$\liminf_{r \rightarrow \infty} \liminf_{n \rightarrow \infty} (\sum_{i=r}^n P_i - \max_{r \leq k \leq n} \sum_{i=r, i \neq k}^n P_{k,i}) \\ \geq \lim_{r \rightarrow \infty} P(\sup_{n \geq r} |X_n| > e) = P(\bigcap_{r=1}^{\infty} \bigcup_{n=r}^{\infty} A_n) \geq \lim_{r \rightarrow \infty} \lim_{n \rightarrow \infty} c_{r,n}.$$

For example, if $P(X_n = 0) = 1 - p/n, P(X_n = 1) = p/n$ and $P(X_n = 1, X_k = 1) = p \min(1/n - 1/n^2, 1/k - 1/k^2)$ for $n \neq k$ where $0 \leq p \leq 6/\pi^2$ then from (4)

$$P(\bigcup_{n=r}^i A_n) \leq \sum_{n=r}^i P_n - \sum_{n=r+1}^i P_{r,n} \\ = (\sum_{n=r}^i (1/n) - \sum_{n=r+1}^i (1/n - 1/n^2))p = 1/r + \sum_{n=r+1}^i (1/n^2).$$

Hence $P(\bigcup_{n=r}^{\infty} A_n) \leq 1/r + \sum_{n=r+1}^{\infty} (1/n^2)$, and $\lim_{r \rightarrow \infty} P(\bigcup_{n=r}^{\infty} A_n) = 0$, i.e., $X_n \rightarrow_{a.s.} 0$ as $n \rightarrow \infty$.

Acknowledgment. I wish to thank the referee for his suggestions which contributed to the presentation of this paper.

REFERENCES

- [1] CHUNG, K. L. and ERDÖS, P. (1952). On the application of the Borel-Cantelli lemma. *Trans. Amer. Math. Soc.* **72** 179-186.
- [2] DAWSON, D. A. and SANKOFF, D. (1967). An inequality for probabilities. *Proc. Amer. Math. Soc.* **18** 504-507.
- [3] GALLOT, S. (1966). A bound for the maximum of a number of random variables. *J. Appl. Prob.* **3** 556-558.
- [4] RAO, C. R. (1965). *Linear Statistical Inference and Its Applications*. Wiley, New York.
- [5] RÉNYI, A. (1962). *Wahrscheinlichkeitsrechnung*. Veb Deutscher Verlag der Wissenschaften, Berlin.