

BOUNDS, INEQUALITIES, AND MONOTONICITY PROPERTIES FOR SOME SPECIALIZED RENEWAL PROCESSES¹

BY MARK BROWN

City College, CUNY Florida State University, and Memorial Sloan-Kettering
Cancer Center

Renewal processes with increasing mean residual life and decreasing failure rate interarrival time distributions are investigated. Various two-sided bounds are obtained for $M(t)$, the expected number of renewals in $[0, t]$. It is shown that if the interarrival time distribution has increasing mean residual life with mean μ , then the expected forward recurrence time is increasing in $t > 0$, as is $M(t) - t/\mu$. If the interarrival time distribution has decreasing failure rate then $M(t)$ is concave, and the forward and backward recurrence time distributions are stochastically increasing in $t > 0$.

1. Introduction and summary. A random variable X with cdf F is defined to have an increasing mean residual life (IMRL) distribution on $[0, \infty)$ if $\mu_1 = EX < \infty$, $F(0^-) = 0$, $F(0) < 1$, and $E(X - t | X > t)$ is increasing for $t \geq 0$. The term increasing (decreasing) is used for monotone nondecreasing (nonincreasing). A random variable X is defined to have a decreasing failure rate (DFR) distribution on $[0, \infty)$ if $F(0^-) = 0$, $F(0) < 1$, and $\Pr(X > s + t | X^* > t)$ is increasing in $t \geq 0$ for each $s > 0$. The DFR property is equivalent to stochastically increasing residual life, which, when $EX < \infty$, implies increasing mean residual life. Thus F DFR and $EX < \infty$ implies F IMRL. It is easy to construct examples for which F is IMRL but not DFR.

In this paper renewal processes with IMRL and DFR interarrival times are studied. A summary of the main results now follows.

If F is DFR then the renewal function, M , is concave and the renewal density, m , which necessarily exists on $(0, \infty)$, is decreasing. The renewal age process, $\{A(t), t \geq 0\}$, and the forward recurrence time process, $\{Z(t), t \geq 0\}$, are stochastically increasing in t . For any Borel set $A \subset [0, \infty)$, $N(\{A + t\})$, the number of renewals in $A + t = \{x + t, x \in A\}$ is stochastically decreasing in $t \geq 0$.

If F is IMRL then the expected forward recurrence time is increasing as is $M(t) - t/\mu_1$. For any Borel set $A \subset [0, \infty)$, $EN(A) \geq \mu_1^{-1}l(A)$, where l is Lebesgue measure.

Bounds for the renewal function are obtained for F IMRL, with improvements for F DFR. If F is IMRL with renewal function M , $\mu_i = EX^i = \int_0^\infty x^i dF(x)$, and

Received November 8, 1976; revised September 6, 1978.

¹Research partially supported by ONR Contract N00014-76-C-0475, and ARO Contract PAAG27-77G-0031.

AMS 1970 subject classifications. Primary 60K05; secondary 60699.

Key words and phrases. Renewal theory, IMRL and DFR distributions, monotonicity properties for stochastic processes, almost sure constructions, future discounted reward process, forward and backward recurrence times, bounds and inequalities for stochastic processes.

$U(t) = t/\mu_1 + \mu_2/2\mu_1^2$, then $\mu_{k+2} < \infty$ for an integer $k \geq 0$ implies

$$(1) \quad U(t) \geq M(t) \geq U(t) - \min_{0 \leq i \leq k} c_i t^{-i}.$$

In (1) c_i is an explicitly computed expression involving μ_1, \dots, μ_{i+2} . Quantities $0 = v_{-1} < v_0 \leq v_1 \dots \leq v_k = \infty$ are found such that for $t \in [v_{j-1}, v_j]$, $c_j t^{-j} = \min_{0 \leq i \leq k} c_i t^{-i}$; thus the lower bound in (1) reduces to $U(t) - c_j t^{-j}$ for $t \in [v_{j-1}, v_j]$. It is further shown that if $\psi_F(a_0) = \int_0^\infty e^{a_0 t} dF(t) < \infty$ for an $a_0 > 0$, then for $0 < a \leq a_0$:

$$(2) \quad U(t) \geq M(t) \geq U(t) - c(F, a)(e^{at} - 1)^{-1}$$

where $c(F, a) = (\mu_1 a)^{-1} - (\mu_2/2\mu_1^2) - (\psi_F(a) - 1)^{-1}$ does not depend on t .

The results follow from a construction of two dependent renewal processes on the same probability space, one an ordinary renewal process and the other an appropriate delayed renewal process. The delayed renewal process has the property that for all k its k th renewal coincides with the $(N + k)$ th renewal of the ordinary renewal process, where N is a random variable. By having a model in which the two processes differ in an easily understandable manner they can readily be compared. The approach of comparing two processes by constructing a convenient bivariate version is widely used (see, for example, [9], [13], [19], [20], [21] and [22]) although the present construction appears new.

If $X - t|X > t$ is stochastically increasing (increasing in mean) it follows (Theorems 2 and 3) that $Z(t)$, the analogous quantity for the renewal process, is stochastically increasing (increasing in mean). More generally it would be of interest to understand the extent to which monotonicity and aging properties of a distribution are inherited in some fashion by its renewal process. Much work remains to be done on this problem.

The theory of DFR distributions is developed in Barlow [1], Barlow, Marshall and Proschan [6], Barlow and Marshall [2], [3], Barlow and Proschan [4], and Esary, Marshall and Proschan [10]. IMRL distributions are examined in Bryson and Siddiqui [8], Haines and Singpurwalla [12], and Barlow, Marshall and Proschan [6].

Mixtures of DFR distributions are DFR ([5], page 103); in particular, mixtures of exponential distributions are DFR. One would, therefore, expect that data collected by combining approximately exponentially distributed subpopulations would have an empirical distribution which would be DFR in appearance. This phenomena is discussed in Barlow and Proschan ([5], page 103). Keilson [14] has shown that a large class of first passage time distributions for Markov processes are DFR.

2. Representations. Let $X_0 \equiv 0$ and X_1, X_2, \dots be i.i.d. with distribution F . Define $S_0 = 0$, $S_n = \sum_{i=1}^n X_i$, $n = 1, 2, \dots$, $N(t) = \max\{i: S_i \leq t\} = \{\# S_i \leq t\} - 1$, and $M(t) = EN(t) + 1 = E\{\# S_i \leq t\}$. Further, define $Z(t) = \sum_1^{N(t)+1} X_i - t$, the forward recurrence time at t , and $A(t) = t - \sum_1^{N(t)} X_i$, the renewal age at t . The

process $\{N(t), t \geq 0\}$ is called an ordinary F renewal process. A class of delayed renewal processes is defined by treating the item at time zero as having a random age T ; when $T = t$ the residual life, which is the first renewal epoch, X'_0 , for the delayed process, has distribution $\bar{F}'_t(x) = \bar{F}(t+x)/\bar{F}(t)$, where $\bar{F} = 1 - F$. If H is the cdf of T then $\Pr(X'_0 > t) = \int_0^\infty \bar{F}'_t(x) dH(x)$. Then let $\{X'_i, i \geq 1\}$ be i.i.d. with distribution F , and independent of X'_0 . Define $S'_n = \sum_0^n X'_i, n \geq 0, S'_{-1} = 0, N'(t) = \max\{i: S'_i \leq t\}$ and $M'(t) = EN'(t) + 1$. The process $\{N'(t), t \geq 0\}$ is called a delayed F renewal process with initial age distribution H . Note that this definition does not allow an arbitrary distribution for X'_0 as the standard definition ([1] page 184). Define $A'(t) = t$ for $0 \leq t < S'_0, A'(t) = t - S'_{N'(t)}$ for $t \geq S'_0$. For a delayed renewal process, $Z', A', N', M', X'_i, S'_n$ will denote the analogues of Z, A, N, M, X_i, S_n for an ordinary renewal process. A delayed renewal process with initial age distribution $G(x) = \mu_1^{-1} \int_0^x \bar{F}(y) dy$ is called a stationary F renewal process. A stationary F renewal process satisfies $M'(t) = t/\mu_1$ and $Z'(t) \sim G$ for all $t \geq 0$ ([11] page 354). Note that the hazard function, h^* , of G satisfies $h^*(t) = 1/E(X - t|X > t)$, where $X \sim F$. Thus F IMRL $\Leftrightarrow G$ DFR.

For a distribution F satisfying $F(0^-) = 0$ and $F(0) < 1$, the definition of DFR given in Section 1 is equivalent to each of the conditions below:

- (i) $\bar{F} = 1 - F$ is log convex.
- (ii) F is absolutely continuous on $(0, \infty)$ with a density f possessing a version for which the hazard function $h(x) = f(x)/\bar{F}(x)$ is decreasing.

The fact that a DFR distribution on $[0, \infty)$, is absolutely continuous except perhaps for an atom at $\{0\}$ can be proved following an argument of Barlow and Proschan [5] page 77. The other implications are straightforward.

We will either assume F IMRL or DFR on $[0, \infty)$. Two dependent renewal processes will be constructed. Process 1 will be an ordinary F renewal process. Process 2 will be a stationary F renewal process in the IMRL case, and delayed F renewal process of the type described above in the DFR case. The special feature of process 2 is that $S'_i = S_{N+i}, i = 0, 1, \dots$, for a random integer N . Process 1 and 2 differ only in that process 2 has zero renewals in $[0, S_N)$ while process 1 has N renewals in this interval. The construction is based on a simple idea which is obscured by the details of the construction and proof. The hazard function of F is decomposed into two components. The first component causes failure for both processes, the second component only causes failure for process 2. The construction uses the following lemma.

LEMMA 1. Let X be distributed as F where F will either be assumed IMRL on $[0, \infty)$ or DFR on $[0, \infty)$. Set $\bar{K}(t) = \bar{G}(t)$ in the IMRL case and $\bar{K}(t) = \int_0^\infty (\bar{F}(t+y)/\bar{F}(y)) dH(y)$, where H is an arbitrary probability distribution on $[0, \infty)$, in the DFR case. Define $\bar{K}_v(t) = \bar{K}(t+v)/\bar{K}(v), \bar{J}_v(t) = \bar{F}(t)/\bar{K}_v(t)$. Then \bar{J}_v is the survival function of a possibly defective distribution on $[0, \infty)$.

PROOF. In the IMRL case $\bar{J}_v(t) = (\bar{F}(t)/\bar{G}(t))(\bar{G}(t)/\bar{G}(t+v))\bar{G}(v)$ and since G is DFR both $\bar{F}(t)/\bar{G}(t)$ and $\bar{G}(t)/\bar{G}(t+v)$ are decreasing. Thus \bar{J}_v is decreasing.

In the DFR case $\bar{J}_v(t) = \bar{K}(v)[\int_0^\infty (\bar{F}(t+v+y)/\bar{F}(t))(\bar{F}(y))^{-1} dH(y)]^{-1}$. Since F is DFR the denominator is increasing, and then, since the numerator is constant, \bar{J}_v is decreasing in t . Thus, in both cases \bar{J}_v is decreasing. In addition J_v is right continuous, equals 1 for $t < 0$, and is always between 0 and 1. It is thus the survival function of a possibly defective distribution on $[0, \infty)$. \square

We proceed with the construction. Again $\bar{K}(t) = \bar{G}(t)$ when F is assumed IMRL, and $\bar{K}(t) = \int_0^\infty (\bar{F}(t+y)/\bar{F}(y)) dH(y)$ with H an arbitrary distribution on $[0, \infty)$ when F is assumed DFR.

Construct Z_1 and W_1 independent with $Z_1 \sim K$, $W_1 \sim J$ where $\bar{J}(t) = \bar{F}(t)/\bar{K}(t)$. If J is defective let $\Pr(Z_1 = \infty) = \lim_{t \rightarrow \infty} \bar{J}(t)$. If $Z_1 \leq W_1$ set $X_1 = X'_0 = Z_1$ and $X_j = X'_{j-1} = Y_{j-1}$, $j = 2, 3, \dots$ where $\{Y_i, i \geq 1\}$ is an i.i.d. sequence with distribution F independent of (Z_1, W_1) . If $Z_1 > W_1$, set $X_1 = W_1$ and go to stage 2. At stage 2 construct Z_2 and W_2 conditionally independent of each other and of (Z_1, W_1) given W_1 , with $Z_2|W_1 = v$ having distribution $\bar{K}_v(t) = \bar{K}(t+v)/\bar{K}(v)$, and $W_2|W_1 = v$ distribution $\bar{J}_v(t) = \bar{F}(t)/\bar{K}_v(t)$. If $Z_2 \leq W_2$ then set $X_2 = Z_2$, $X'_0 = W_1 + Z_2$ and $X_j = X'_{j-2} = Y_{j-2}$, $j = 3, 4, \dots$ where $\{Y_i, i = 1, 2, \dots\}$ is i.i.d. with distribution F and independent of (Z_1, W_1, Z_2, W_2) . If $W_2 < Z_2$ set $X_2 = W_2$ and go to stage 3. We reach stage m if and only if $W_i < Z_i$, $i = 1, \dots, m-1$, in which case $X_i = W_i$, $i = 1, \dots, m-1$. At stage m we construct Z_m and W_m conditionally independent of each other and of $(Z_1, W_1) \dots (Z_{m-1}, W_{m-1})$ given $\sum_{i=1}^{m-1} W_i$, with $(Z_m|\sum_{i=1}^{m-1} W_i = v) \sim K_v$, $(W_m|\sum_{i=1}^{m-1} W_i = v) \sim J_v$. If $Z_m \leq W_m$ set $X_m = Z_m$, $X'_0 = \sum_{i=1}^{m-1} W_i + Z_m$, $X'_j = X'_{j-m} = Y_{j-m}$, $j = m+1, \dots$ where $\{Y_i, i \geq 1\}$ is i.i.d. with distribution F and independent of $(Z_1, W_1, Z_2, W_2, \dots, Z_{m-1}, W_{m-1})$. If $Z_m > W_m$ then go to stage $m+1$ and repeat.

THEOREM 1. *Under the above construction:*

- (i) $\{X_i, i \geq 1\}$ is an i.i.d. sequence with distribution F .
- (ii) $\{X'_i, i \geq 0\}$ are independent, $X'_0 \sim K$, $X'_j \sim F$ for $j \geq 1$.
- (iii) $S'_i = S_{N+i}$ for $i \geq 0$, where $N = \min\{i: Z_i \leq W_i\}$, and $\Pr(N < \infty) = 1$.

PROOF. (i) Define $N = \min\{i: Z_i \leq W_i\}$, $N = \infty$ if $W_i < Z_i$ for all i . Now $(X_i|N \geq i, (W_j, Z_j) = (w_j, z_j), j = 1, \dots, i-1) \sim \min(Z_v^*, W_v^*)$ where $v = \sum_{i=1}^{i-1} w_j$, $Z_v^* \sim K_v$, $W_v^* \sim J_v$ and Z_v^* and W_v^* are independent. Since $\bar{K}_v(t)\bar{J}_v(t) = \bar{F}(t)$, $(X_i|N \geq i, (W_j, Z_j) = (w_j, z_j), j = 1, \dots, i-1) \sim F$. Thus $X_i \sim F$ independent of X_1, \dots, X_{i-1} for all i ; thus $\{X_j, j \geq 1\}$ is i.i.d. with distribution F .

(ii) In our construction we generated W_1, \dots, W_N . It will not be convenient to continue constructing W_j 's for $j > N$. At stage j construct W_j to be conditionally independent of W_1, \dots, W_{j-1} given $\sum_{i=1}^{j-1} W_i$, with $(W_j|\sum_{i=1}^{j-1} W_i = v) \sim J_v$. Since K is DFR, $\inf_i \Pr(W_i > t) \geq \inf_v \Pr(W_v > t) = \bar{F}(t)/\lim_{v \rightarrow \infty} \bar{K}_v(t) > 0$. Therefore, $\Pr(\sum_{i=1}^\infty W_i = \infty) = 1$, so that given t , with probability 1 there exists j so that

$\sum_1^{j-1} W_i < t \leq \sum_1^j W_i$. Then:

$$\Pr(X'_0 > t | w_1, w_2, \dots) = \left(\prod_{i=1}^{j-1} \bar{K}_{\sum_1^{i-1} w_i}(w_i) \right) \left(\bar{K}_{\sum_1^{j-1} w_i}(t - \sum_1^{j-1} w_i) \right) = \bar{K}(t).$$

Thus $X'_0 \sim K$.

(iii) Since $\sum_1^\infty W_i = \infty$ with probability 1, $X'_0 < \infty$ if and only if $N < \infty$. By (ii), $X'_0 \sim K$ thus $\Pr(X'_0 < \infty) = 1$ and, therefore, $\Pr(N < \infty) = 1$. By construction $S'_i = S_{N+i}$ for $i \geq 0$. \square

3. Some properties of IMRL and DFR renewal processes. We will need the following well-known result (Feller [11] page 148).

LEMMA 2. Let F be a distribution on $[0, \infty)$. If $\mu_1 < \infty$ define $\bar{F}_1(t) = \int_t^\infty \bar{F}(x) dx$; if $\mu_2 < \infty$ define $\bar{F}_2(t) = \int_t^\infty \bar{F}_1(x) dx$. Then:

(i) $\mu_k < \infty \Rightarrow t^k \bar{F}(t) \rightarrow 0$ as $t \rightarrow \infty$; for $k \geq 0, \mu_{k+1} < \infty \Rightarrow \int_0^\infty t^k \bar{F}(t) dt = \mu_{k+1}/k + 1 < \infty$ and $t^k \bar{F}_1(t) \rightarrow 0$ as $t \rightarrow \infty$; for $k \geq 0, \mu_{k+2} < \infty \Rightarrow \int_0^\infty t^k \bar{F}_1(t) dt = \mu_{k+2}/(k+1)(k+2) < \infty$ and $t^k \bar{F}_2(t) \rightarrow 0$ as $t \rightarrow \infty$; for $k \geq 0, \mu_{k+3} < \infty \Rightarrow \int_0^\infty t^k \bar{F}_2(t) dt = \mu_{k+3}/(k+1)(k+2)(k+3) < \infty$.

(ii) For $a > 0, \psi_F(a) = \int_0^\infty e^{at} dF(t) < \infty$ implies $e^{at} \bar{F}(t), e^{at} \bar{F}_1(t),$ and $e^{at} \bar{F}_2(t)$ converge to 0 as $t \rightarrow \infty$. Moreover $\int_0^\infty e^{at} \bar{F}(t) dt = a^{-1}(\psi_F(a) - 1) < \infty,$ $\int_0^\infty e^{at} \bar{F}_1(t) dt = a^{-2}(\psi_F(a) - a\mu_1 - 1) < \infty,$ and $\int_0^\infty e^{at} \bar{F}_2(t) dt = a^{-3}(\psi_F(a) - a^2(\mu_2/2) - a\mu_1 - 1) < \infty.$

If $\mu_2 = EX^2 < \infty,$ define $L(t) = M(t) - t/\mu_1 - \mu_2/2\mu_1^2$. Define $q = \bar{F}(0) = \Pr(X > 0)$. For nonnegative extended real-valued random variables $EX \geq EY$ is defined to mean $\infty \geq EX \geq EY$. For extended real-valued random variables, $X \geq_{st} Y$ (X stochastically greater than Y) means $\Pr(X > a) \geq \Pr(Y > a)$ for all finite a . In Theorems 2 and 3 below, many of the random variables may be improper (assign positive probability to $\pm\infty$) and the above conventions for $EX \geq EY$ and $X \geq_{st} Y$ apply.

THEOREM 2. Let $\{N(t), t \geq 0\}$ be an ordinary renewal process with F IMRL, and $\{N'(t), t \geq 0\}$ a stationary F renewal process. Then:

(i) $M(t) - t/\mu_1$ and $EZ(t)$ (expected forward recurrence time at t) are increasing in $t \geq 0$. If $\mu_2 < \infty$ then $L(t) = M(t) - t/\mu_1 - \mu_2/2\mu_1^2 \uparrow 0$ as $t \rightarrow \infty$.

(ii) If g is a nonnegative measurable function then $\sum_0^\infty g(S_i)$ is stochastically larger than $\sum_0^\infty g(S'_i)$. In particular $N(A)$ is stochastically larger than $N'(A)$ for all Borel sets $A \subset [0, \infty)$, and $EN(A) \geq EN'(A) = \mu_1^{-1}l(A)$, where l is Lebesgue measure. Furthermore, $M(t+h) - M(t) \geq h/\mu_1$ for all $t \geq 0, h \geq 0,$ and converges to h/μ_1 as $t \rightarrow \infty$.

(iii) If $\mu_2 < \infty$ and g is a bounded measurable function which converges to 0 as $t \rightarrow \infty$ then $|\int_0^t g(t-x) dM(x) - \mu_1^{-1} \int_0^t g(x) dx| \rightarrow 0$ as $t \rightarrow \infty$.

(iv) If $\mu_2 < \infty$ then $0 \geq L(t) \geq -\mu_1^{-1} \int_t^\infty (\bar{G}(x) - q^{-1} \bar{F}(x)) dx \geq -\mu_1^{-1} \int_t^\infty \bar{G}(x) dx$ where $q = \bar{F}(0), L(t) = M(t) - t/\mu_1 - \mu_2/2\mu_1^2$.

(v) For $k \geq 0, \mu_{k+2} < \infty \Rightarrow \lim_{t \rightarrow \infty} t^k L(t) = 0$ and $\mu_{k+3} < \infty \Rightarrow 0 \geq \int_0^\infty t^k L(t) dt$

$> -\infty$; for $a > 0$, $\psi_F(a) = \int_0^\infty e^{at} dF(t) < \infty \Rightarrow \lim_{t \rightarrow \infty} e^{at} L(t) = 0$ and $0 > \int_0^\infty e^{at} L(t) dt > -\infty$.

PROOF. Recall that process 1 is an ordinary F renewal process, process 2 a stationary F renewal process, and $S'_i = S_{N+i}$, $i = 1, 2, \dots$. Note that by Wald's identity $EN = \mu_2/2\mu_1^2$ for $\mu_1 < \infty$ with both sides equal to ∞ if $\mu_2 = \infty$.

(i) $M(t) - t/\mu_1 = E(N(t) - N'(t))$. Since $N(t) - N'(t) \uparrow N$, $M(t) - t/\mu_1 \uparrow EN = \mu_2/2\mu_1^2$ by the monotone convergence theorem. Thus if $\mu_2 < \infty$ then $0 \geq L(t) = M(t) - t/\mu_1 - \mu_2/2\mu_1^2 \uparrow 0$. Since $Z(t) = \sum_1^{N(t)+1} X_i - t$, it follows by Wald's identity that $EZ(t) = \mu_1(M(t) - t/\mu_1)$, thus $EZ(t) \uparrow$.

(ii) By construction, $\sum_0^\infty g(S_i) \geq \sum_N^\infty g(S_i) = \sum_0^\infty g(S'_i)$, thus $\sum g(S_i)$ is stochastically larger than $\sum g(S'_i)$. Setting $g = I_A$, the indicator function of the Borel set $A \subset [0, \infty)$, gives $N(A)$ stochastically greater than $N'(A)$; taking expectations we get, $EN(A) \geq EN'(A) = \mu_1^{-1}l(A)$. Setting $A = [t, t+h)$ gives $M(t+h) - M(t) \geq h/\mu_1$. This last inequality and the elementary renewal theorem ($M(t)/t \rightarrow \mu_1^{-1}$) imply that $M(t+h) - M(t) \rightarrow h/\mu_1$ as $t \rightarrow \infty$. (The convergence of $M(t+h) - M(t)$ also follows from Blackwell's theorem, as an IMRL distribution must be nonlattice.)

(iii) Define $Y(t) = \sum_{i=0}^{\min(N-1, N(t))} g(t - S_i)$. Now $\int_0^t g(t-x) dM(x) - \mu_1^{-1} \int_0^t g(x) dx = E[\sum_{i=0}^{N(t)} g(t - S_i) - \sum_{i=0}^{N(t)} g(t - S'_i)] = EY(t)$. Now $Y(t) \rightarrow 0$ a.s. as $t \rightarrow \infty$ and $|Y(t)| \leq N(\sup|g|)$ for all t . Since $\mu_2 < \infty$ the result follows from the dominated convergence theorem. An alternative proof is based on the fact that $|\int_0^t g(t-x) dM(x) - \mu_1^{-1} \int_0^t g(x) dx| \leq \sup(|g|)$ multiplied by the total variation of $M(x) - x/\mu_1$ on $[0, \infty)$; the latter quantity is finite when $\mu_2 < \infty$ by part (i) of this theorem.

(iv) By the argument in part (i), $L(t) = \mu_1^{-1}E(Z(t) - Z'(t))$. By construction:

$$\begin{aligned} 0 \geq Z(t) - Z'(t) &= 0 && \text{if } t \geq X'_0 \\ &= Z(t) - (X'_0 - t) && \text{if } t < X'_0. \end{aligned}$$

Thus $\mu_1 L(t) = \bar{G}(t)E(Z(t)|X'_0 > t) - \int_t^\infty \bar{G}(x) dx$. Since $Z(t)|X'_0 > t$ is a mixture of distributions of the form $X - v|X > v$ where $X \sim F$, and F is IMRL, it follows that $E(Z(t)|X'_0 > t) \geq E(X|X > 0) = q^{-1}\mu_1$. Thus $0 \geq L(t) \geq -\mu_1^{-1} \int_t^\infty (\bar{G}(x) - q^{-1}\bar{F}(x)) dx \geq -\mu_1^{-1} \int_t^\infty \bar{G}(x) dx$.

(v) These results follow from (iv) and Lemma 2.

THEOREM 3. Suppose that F is DFR on $[0, \infty)$. Then:

(i) $A(t)$ and $Z(t)$ are stochastically increasing in $t \geq 0$.

(ii) The renewal density m on $(0, \infty)$, has a version which is decreasing and converges to μ_1^{-1} as $t \rightarrow \infty$ (if $\mu_1 = \infty$ then $\mu_1^{-1} = 0$); $M(t)$ is concave.

(iii) If g is a nonnegative measurable function then $V(t) = \sum_{i=1}^\infty g(S_{N(t)+i} - t)$ is stochastically decreasing in $t \geq 0$. In particular, for any Borel set $A \subset [0, \infty)$, $N(\{A + t\})$ is stochastically decreasing in $t \geq 0$; $N(t+h) - N(t)$ is stochastically decreasing in $t \geq 0$ for each $h \geq 0$, and $M(t+h) - M(t) \downarrow h/\mu_1$ as $t \rightarrow \infty$.

(iv) If g is a measurable function, $R^\infty \rightarrow \bar{R} = [-\infty, \infty]$, which is decreasing (increasing) then $W(t) = g(\{S_{N(t)+i} - t, i \geq 1\})$ is stochastically decreasing (increasing) in $t \geq 0$. (By g increasing we mean $x_i \leq y_i, i \geq 1$, implies $g(x) \leq g(y)$.)

PROOF. (i) Give process 2 initial age distribution $F_A(s)$, the age distribution of process 1 at time s . Then $Z'(t) \sim Z(s + t)$, and $A'(t) \sim \min(A(s + t), t)$ so that $A'(t) \leq_{st} A(s + t)$. By construction $Z'(t) \geq Z(t)$, and $A'(t) \geq A(t)$. Thus $Z(t) \leq_{st} Z'(t) \sim Z(s + t)$ and $A(t) \leq_{st} A'(t) \leq_{st} A(s + t)$. Therefore, both $A(t)$ and $Z(t)$ are stochastically increasing in $t \geq 0$.

(ii) Recall that a DFR distribution on $[0, \infty)$ is necessarily absolutely continuous on $(0, \infty)$ and possesses a version of its hazard function, $h(x) = f(x)/\bar{F}(x)$, which is decreasing. Now $m(t) = q^{-1}E(h(A(t)))$, where m is the renewal density, $q = \bar{F}(0)$, and $A(t)$ is the renewal age at time t . Since $A(t)$ is stochastically increasing (part (i)), and h is decreasing, it follows that $h(A(t))$ is stochastically decreasing and therefore $E(h(A(t)))$ is decreasing. Thus $m(t)$ is decreasing and therefore converges as $t \rightarrow \infty$. By the elementary renewal theorem ($M(t)/t \rightarrow \mu_1^{-1}$) the limit must be μ_1^{-1} . Concavity of M follows from m decreasing.

(iii) Define process 2 as in (i) above. By construction $V(t) \geq V'(t) \sim V(s + t)$. Thus $V(t)$ is stochastically decreasing in $t \geq 0$. For a Borel set $A \subset (0, \infty)$, $g = I_A$ yields $N(\{A + t\})$ stochastically decreasing. If $0 \in A \subset [0, \infty)$ with $B = A - \{0\}$ then $N(\{A + t\}) = N(\{B + t\})$ a.s. for each $t > 0$, while $N(\{A\}) > N(\{B\})$; thus the result also holds for the case $0 \in A$. Setting $A = (0, h]$ gives $N(t + h) - N(t)$ stochastically decreasing, and taking expectations gives $M(t + h) - M(t) \downarrow$; thus $M(t + h) - M(t)$ has a limit as $t \rightarrow \infty$, which by the elementary renewal theorem must be h/μ_1 .

(iv) Consider an ordinary F renewal process. An observer enters at time s and records renewal epochs in (s, ∞) subtracting s from each one. He then is observing a sequence $\{S'_n, n \geq 0\}$ with $S'_0 \sim Z_s$ and $S'_n \sim Z_s + S_n$ for $n \geq 1$. Similarly if he enters at time t he observes $\{S''_n, n \geq 0\}$ with $S''_0 \sim Z_t$ and $S''_n \sim Z_t + S_n$ for $n \geq 1$. From part (i) we know that for $t > s$, Z_t is stochastically larger than Z_s . It follows (Lehman [16], page 73) that we can construct a pair of random variables (S'_0, S''_0) with $S'_0 \sim Z_s$, $S''_0 \sim Z_t$, and $S'_0 \leq S''_0$ a.s. Next take an i.i.d. sequence $\{X_i, i \geq 1\}$ with distribution F , independent of (S'_0, S''_0) , define $S_n = \sum_{i=1}^n X_i, n \geq 1$, and set $S'_n = S'_0 + S_n, S''_n = S''_0 + S_n, n \geq 1$. We therefore have versions of the future renewal processes viewed from s and t with $S''_n - S'_n = S''_0 - S'_0 \geq 0$ a.s. Therefore, for g decreasing $g(\{S'_n, n \geq 0\}) \geq g(\{S''_n, n \geq 0\})$ a.s., thus $W(s) \geq_{st} W(t)$ and $W(t)$ is stochastically decreasing in $t \geq 0$. \square

REMARK. Suppose that $\{Y_i, i \geq 1\}$ is a sequence of nonnegative random variables independent of $\{S_i, i \geq 0\}$. Consider $R(s) = \sum_{i=1}^\infty Y_i e^{-\lambda(S_{N(s)+i}-s)} = \sum_{i=N(s)+1}^\infty Y_{i-N(s)} e^{-\lambda(S_i-s)}$. For $\lambda > 0$, $R(s)$ is known as the future discounted reward at time s , and λ is called the discount factor. The process $\{R(s), s \geq 0\}$ is of interest in various stochastic optimization models (Ross [23]). It is easy to show

that $\lambda > 0$ and $\sup_i EY_i < \infty$ implies that $ER(t) < \infty$ and thus that $R(t)$ is a proper random variable for all t .

It follows from either part (iii) or part (iv) of Theorem 3, by conditioning on $\{Y_i = y_i, i = 1, \dots\}$, that if F is DFR, then $R(s)$ is stochastically decreasing in $s \geq 0$.

4. Bounds for M(t): IMRL case.

THEOREM 4. (i) *If F is IMRL on $[0, \infty)$, and $\mu_{k+2} < \infty$ for an integer $k \geq 0$ then:*

$$(3) \quad U(t) \geq M(t) \geq U(t) - \min_{0 \leq i \leq k} c_i t^{-i}$$

where $U(t) = t/\mu_1 + \mu_2/2\mu_1^2$, $c_0 = \mu_2/2\mu_1^2 - q^{-1}$ and

$$0 \leq c_i = -i \int_0^\infty s^{i-1} L(s) ds = \int_0^\infty s^i d(M(s) - s/\mu_1) < \infty$$

for $i = 1, \dots, k$.

The term c_i is a function of μ_1, \dots, μ_{i+2} , $i = 1, 2, \dots, k$ which can be recursively computed from:

$$(4) \quad c_i = \gamma_i - \mu_1^{-1} i! \sum_{s=1}^{i-1} (c_s/s!) \lambda_{i+1-s}, \quad i = 1, \dots, k$$

where

$$\gamma_i = [\mu_{i+2}/(i+1)(i+2)\mu_1^2] - [\mu_2\mu_{i+1}/2(i+1)\mu_1^3]$$

$$\lambda_i = \mu_i/i!$$

Equation (4) can be explicitly solved yielding

$$(5) \quad c_i = \gamma_i + i! \sum_{j=1}^{i-1} (\gamma_j/j!) \sum_{l=1}^{i-j} (-\mu_1)^{-l} \sum_{(i_1, \dots, i_l) \in A_{i-j,l}} (\prod_{r=1}^l \lambda_{i_r+1}),$$

$i = 1, \dots, k$

where

$$A_{k,l} = \{(i_1, \dots, i_l): i_r \geq 1, r = 1, \dots, l, \sum_{r=1}^l i_r = k\}$$

(ii) *If F is quasiexponential ($\bar{F}(t) = qe^{-\lambda t}$, $0 < q \leq 1$, $\lambda > 0$, $t \geq 0$) then $c_i = 0$ for all i . If F is IMRL but nonquasiexponential then $c_i > 0$ for all i .*

In the nonquasiexponential case define $v_{-1} = 0$, $v_i = c_{i+1}/c_i$, $i = 0, 1, \dots$ with $\infty/\infty = \infty$. Then v_i is increasing in i and for $v_{j-1} \leq t \leq v_j$, $c_j t^{-j} = \inf_i c_i t^{-i}$.

PROOF. (i) Since $M(t) = U(t) + L(t)$ and $L(t) \leq 0$ (Theorem 2, (i)) $M(t) \leq U(t)$. Since $L(t)$ is increasing (Theorem 2, (i)) $L(t) \geq L(0) = M(0) - \mu_2/2\mu_1^2 = q^{-1} - \mu_2/2\mu_1^2 = -c_0$. Thus $M(t) \geq U(t) - c_0$. Since $L(t)$ is increasing it is at least as big as its average over $[0, t]$ with respect to any probability measure on $[0, t]$. Thus $L(t) \geq it^{-i} \int_0^\infty s^{i-1} L(s) ds$ which since $L \leq 0$ exceeds $it^{-i} \int_0^\infty s^{i-1} L(s) ds = -c_i t^{-i} > -\infty$ by Theorem 2 part (v). The equivalence between $-i \int_0^\infty s^{i-1} L(s) ds$ and $\int_0^\infty s^i d(M(s) - s/\mu_1)$ follows by integration by parts and part (v) of Theorem 2.

To compute the c_i 's start with the identity $M(t) = 1 + \int_0^t M(t-x) dF(x)$, subtract $U(t)$ from both sides, multiply both sides by t^i and use the identity $t^i - (t -$

$x)^i = \sum_{r=0}^{i-1} \binom{i}{r} (t-x)^r x^{i-r}$ This yields:

$$(6) \quad t^i L(t) = \int_0^t (t-x)^i L(t-x) dF(x) + h(t)$$

where $h(t) = h_1(t) + h_2(t) - h_3(t)$, $h_1(t) = \int_0^t [\sum_{r=0}^{i-1} \binom{i}{r} (t-x)^r x^{i-r}] L(t-x) dF(x)$, $h_2(t) = t^i \mu_1^{-1} \int_0^\infty \bar{F}(x) dx$, $h_3(t) = t^i (\mu_2/2\mu_1^2) \bar{F}(t)$. Now (6) is the renewal equation $g = h + g * F = h * M$, with $g(t) = t^i L(t)$. By part (iii) of Theorem 2, if we can show that h is bounded, integrable and that $\lim_{t \rightarrow \infty} h(t) = 0$ then we can conclude that $\lim_{t \rightarrow \infty} t^i L(t) = \mu_1^{-1} \int_0^\infty h(t) dt$. But for $t = 0, \dots, k$, $\lim_{t \rightarrow \infty} t^i L(t) = 0$ by part (v) of Theorem 2. Thus the conclusion will reduce to $\int_0^\infty h(t) dt = 0$ and this identity provides us with expressions for c_i .

To show that h is bounded, integrable and convergent to zero we do so separately for each h_i . By Lemma 2, h_2 and h_3 are convergent to zero with:

$$(7) \quad \int_0^\infty h_2(t) dt = \mu_{i+2} / (i+1)(i+2)\mu_1$$

$$(8) \quad \int_0^\infty h_3(t) dt = \mu_{i+1} \mu_2 / 2(i+1)\mu_1^2$$

The boundedness of h_2 and h_3 follows from the boundedness on finite intervals and the convergence to zero as $t \rightarrow \infty$.

Defining $L(y) = 0$ for $y < 0$ we write $h_1(t) = \int_0^\infty (\sum_{r=0}^{i-1} \binom{i}{r} x^{i-r} ((t-x)^r L(t-x)) dF(x)$. Since $(t-x)^r L(t-x) \rightarrow 0$ as $t \rightarrow \infty$ by Theorem 2 part (v) the integrand converges pointwise to zero. Since $s^r |L(s)|$ is bounded on finite intervals and converges to 0 as $s \rightarrow \infty$, $\sup_s s^r |L(s)| < \infty$, thus the integrand is dominated by the integrable function $\sum_{r=0}^{i-1} \binom{i}{r} (\sup_s s^r |L(s)|) x^{i-r}$ and by the dominated convergence theorem $h_1(t) \rightarrow 0$. The above argument also shows that $|h_1(t)| \leq \sum_{r=0}^{i-1} \binom{i}{r} \mu_{i-r} (\sup_s s^r |L(s)|) < \infty$, thus h_1 is bounded; h_1 is integrable by part (v) of Theorem 2 since $\int_0^\infty |h_1(t)| dt = \sum_{r=0}^{i-1} \binom{i}{r} \mu_{i-r} \int_0^\infty s^r |L(s)| ds < \infty$. Moreover:

$$(9) \quad \int_0^\infty h_1(t) dt = \sum_{r=0}^{i-1} \binom{i}{r} \mu_{i-r} \int_0^\infty s^r L(s) ds = -i! \sum_{r=1}^i (c_r/r!) \lambda_{i+1-r}$$

The identity $\int_0^\infty h(t) dt = 0$ is equivalent, using (7), (8) and (9), to $i! \sum_{r=1}^i (c_r/r!) \lambda_{i+1-r} = \mu_1 \gamma_i$ which reduces to (4).

Define $d_0 = 0$, $d_i = c_i/i!$, $i = 1, \dots, k$, $\delta_0 = 0$, $\delta_i = \gamma_i/i!$, $i = 1, \dots, k$, $\beta_0 = 0$, $\beta_i = -\mu_1^{-1} \lambda_{i+1}$, $i = 1, \dots, k$. Then rewrite (4) as:

$$(10) \quad d_i = \delta_i + \sum_{s=0}^i d_s \beta_{i-s}$$

Now (10) is a discrete renewal equation. Its solution is $d_i = \sum_{j=0}^i \delta_j M_{i-j}$ where $M_i = \sum_{s=0}^\infty \beta_i^{(s)}$, where $\beta_i^{(s)}$ is the s th convolution of β . Since $\beta_0 = 0$, $\beta_i^{(s)} = 0$ for $s > i$. We thus obtain

$$d_i = \sum_{j=1}^i \delta_j M_{i-j} = \delta_i + \sum_{j=1}^{i-1} \delta_j M_{i-j} = \delta_i + \sum_{j=1}^{i-1} \delta_j \sum_{l=1}^{i-j} \sum_{(i_1, \dots, i_l) \in A_{i-j, l}} (\prod_{t=1}^l \beta_{i_t})$$

which is equivalent to (5).

(ii) If F is quasiexponential ($\bar{F}(t) = qe^{-\lambda t}$) then $M(t) = q^{-1} \lambda t + q^{-1} = t/\mu_1 + \mu_2/2\mu_1^2 (\mu_1 = \lambda q, \mu_2 = 2\lambda^2 q)$ thus $L(t) = 0$. If $c_i = -i \int_0^\infty s^{i-1} L(s) ds = 0$ for some

i , then since $L < 0$ and $L \uparrow$ for F IMRL (Theorem 2 (i)), it follows that $L(s) \equiv 0$ or equivalently $M(t) = at + b$ with $b = M(0) \geq 1$. But the quasiexponential distribution with $q = b^{-1} \leq 1$ and $\lambda = ab^{-1}$ has renewal function $M(t) = at + b$ and the renewal function uniquely determines the distribution. ($\psi_M = (1 - \psi_F)^{-1}$ so the Laplace transform of F determines the Laplace transform of M which determines M). Thus for F IMRL but not quasiexponential, $c_i > 0$ for all i .

Since $L \uparrow$ on $[0, \infty)$ (Theorem 2 (i)) we can interpret L as the distribution function of a measure which assigns negative weight to $\{0\}$ but positive weight to all Borel sets in $(0, \infty)$. Consider the Hilbert space of Borel measurable functions on $(0, \infty)$ satisfying $\int_0^\infty f^2 dL < \infty$ with inner product $(f, g) = \int_0^\infty fg dL$. Now:

$$c_j^2 = \left(\int_0^\infty s^j dL(s)\right)^2 = \left(t^{\frac{j+1}{2}}, t^{\frac{j-1}{2}}\right)^2$$

$$\leq \|t^{\frac{j+1}{2}}\|^2 \|t^{\frac{j-1}{2}}\|^2 = c_{j+1}c_{j-1}$$

by the Cauchy-Schwartz inequality. Note that $c_i = \infty \Rightarrow c_{i+1} = \infty$ so that the inequality $c_j^2 \leq c_{j+1}c_{j-1}$ holds even if not all of c_{j-1}, c_j, c_{j+1} are finite. The inequality is equivalent to $v_j = c_{j+1}/c_j \geq c_j/c_{j-1} = v_{j-1}$ (where $\infty/\infty = \infty$). Thus the v_j 's are increasing.

Suppose $t \leq v_j$. Then since $v_j \uparrow, t^m \leq \prod_{i=0}^{m-1} v_{j+i} = c_{j+m}/c_j$ for $m = 1, 2, \dots$. But the inequality $t^m \leq c_{j+m}/c_j$ is equivalent to $c_j t^{-j} \leq c_{j+m} t^{-(j+m)}$. Similarly if $t \geq v_{j-1}, c_j t^{-j} \leq c_{j-m} t^{-(j-m)}$ for $m = 1, \dots, j$. Thus for $t \in [v_{j-1}, v_j], c_j t^{-j} = \inf_i c_i t^{-i}$. \square

THEOREM 5. *If F is IMRL on $[0, \infty)$ and $\psi_F(a_0) = \int_0^\infty e^{as} dF(s) < \infty$ for an $a_0 > 0$, then for $0 < a \leq a_0$:*

(11)

$$U(t) \geq M(t) \geq U(t) - (e^{at} - 1)^{-1} \left[(\mu_1 a)^{-1} - (\mu_2/2\mu_1^2) - (\psi_F(a) - 1)^{-1} \right].$$

PROOF. The proof is very similar to that of Theorem 4. Choose $a \in (0, a_0]$. Using $L(t) \leq 0, L(t) \uparrow$ as in the proof of Theorem 4 we obtain:

(12)
$$U(t) \geq M(t) \geq U(t) + a(e^{at} - 1)^{-1} \int_0^\infty e^{as} L(s) ds$$

where $0 \geq \int_0^\infty e^{as} L(s) ds > -\infty$ by part (v) of Theorem 2.

To evaluate $\psi_L(a) = \int_0^\infty e^{as} L(s) ds$ we start with $M(t) = 1 + \int_0^t M(t-x) dF(x)$, subtract $U(t)$ from each side and multiply both sides by e^{at} . This gives:

(13)
$$e^{at}L(t) = \int_0^t e^{a(t-x)}L(t-x) dF(x) + l(t)$$

where

$$l(t) = l_1(t) + l_2(t) - l_3(t), \quad l_1(t) = \int_0^t (e^{ax} - 1)e^{a(t-x)}L(t-x) dF(x),$$

$$l_2(t) = \mu_1^{-1}e^{at} \int_0^\infty \bar{F}(x) dx, \quad l_3(t) = (\mu_2/2\mu_1^2)e^{at}\bar{F}(t).$$

We verify the conditions of part (iii) of Theorem 2 in a similar manner as in the proof of Theorem 4, making heavy use of Lemma 2 and part (v) of Theorem 2. The

conclusion of part (iii) of Theorem 2, in light of $\lim_{t \rightarrow \infty} e^{at}L(t) = 0$ (part (v) of Theorem 2) gives us:

(14)

$$(\psi_F(a) - 1)\psi_L(a) = (\mu_2/2\mu_1^2)[(\psi_F(a) - 1)/a] - (\mu_1a^2)^{-1}(\psi_F(a) - \mu_1a - 1).$$

Since $\psi_F(a) - 1 \neq 0$ for $a \neq 0$ we can divide both sides of (14) by $\psi_F(a) - 1$ and solve for $\psi_L(a)$. This gives:

$$(15) \quad \psi_L(a) = \mu_2/2\mu_1^2a - (\mu_1a^2)^{-1} + [a(\psi_F(a) - 1)]^{-1}.$$

Substituting (15) into (12) gives us (11). \square

REMARK. (11) and (15) will hold for $a < 0$ whether or not $\psi_F(a) < \infty$ for an $a > 0$. If $\mu_3 < \infty$ and we let $a \uparrow 0$ in (11) then we obtain $M(t) \geq U(t) - c_1t^{-1}$ where c_1 is given in (4). In general $\mu_{k+3} < \infty$ implies $\psi_L^{(k)}(0^-)$ exists and equals $-(k + 1)^{-1}c_{k+1}$. Thus $\psi_L(a)$ can be considered as a generating function for the c_i 's. However, unless the particular form of $\psi_F(a)$ leads to a simple expression for $\psi_L(a)$, expressions (4) and (5) of Theorem 4 will be preferable for computing the c_i 's.

5. Improved bounds when F is DFR. The bounds given in Theorems 4 and 5 for IMRL distributions can be improved for DFR distributions. Define $\alpha_0 = 1$, $\alpha_i = (i/i + 1)^i$, $i \geq 1$, $c_i^* = \alpha_i c_i$ where c_i is given in Theorem 3, and $v_i^* = c_{i+1}^*/c_i^* = (\alpha_{i+1}/\alpha_i)v_i$. Also define $g_a(t) = a(e^{at} - 1)^{-1} \int_0^t s e^{as} ds = (te^{at}/e^{at} - 1) - a^{-1}$, $h_a(t) = (1 + at)(1 - e^{-at})$, and $\psi(a) = -a\psi_L(a) = [(\mu_1a)^{-1} - (\mu_2/2\mu_1^2) - (\psi_F(a) - 1)^{-1}]$.

COROLLARY 1. Assume that F is DFR on $[0, \infty)$. Then:

- (i) If $\mu_{k+2} < \infty$ then $U(t) \geq M(t) \geq U(t) - \min_{0 \leq i \leq k} c_i^* t^{-i}$;
- (ii) $v_i^* \uparrow$ and for $v_{j-1}^* \leq t \leq v_j^*$, $c_j^* t^{-j} = \inf_i c_i^* t^{-i}$; thus for $v_{j-1}^* \leq t \leq v_j^*$ the bound in (i) is given by $U(t) - c_j^* t^{-j}$.
- (iii) If $\psi_F(a_0) < \infty$ for an $a_0 > 0$ then for $0 < a \leq a_0$:

$$U(t) \geq M(t) \geq U(t) - (e^{ag_a^{-1}(t)} - 1)^{-1}\psi(a) \geq U(t) - (e^{h_a(t)} - 1)^{-1}\psi(a).$$

PROOF. (i) L is concave by Theorem 3 part (ii). Thus $L(jt^{-j} \int_0^t s \cdot s^{j-1} ds) = L((j/j + 1)t) \geq jt^{-j} \int_0^t s^{j-1} L(s) ds \geq jt^{-j} \int_0^\infty s^{j-1} L(s) ds = -c_j t^{-j}$. Thus $L(t) \geq -c_j [(j + 1/j)t]^{-j} = -c_j^* t^{-j}$. The argument now proceeds as in Theorem 4.

(ii) A simple differentiation argument shows that $\alpha_{i+1}/\alpha_i \uparrow$. Since $c_{i+1}/c_i \uparrow$ by Theorem 4 and $c_{i+1}^*/c_i^* = (\alpha_{i+1}/\alpha_i)(c_{i+1}/c_i)$, we see that $c_{i+1}^*/c_i^* \uparrow$. The argument now proceeds as in Theorem 4.

(iii) The concavity argument in (i) shows that $L(g_a(t)) \geq a(e^{at} - 1)^{-1} \int_0^t e^{as} L(s) ds \geq -(e^{at} - 1)^{-1}(-a\psi_L(a))$; thus $L(t) \geq -(e^{-ag_a^{-1}(t)} - 1)^{-1}(-a\psi_L(a)) = -(e^{ag_a^{-1}(t)} - 1)^{-1}[(\mu_1a)^{-1} - (\mu_2/2\mu_1^2) - (\psi_F(a) - 1)^{-1}]$. If $s = g_a^{-1}(t)$ then $s = (t + a^{-1})(1 - e^{-as}) \geq (t + a^{-1})(1 - e^{-at}) = a^{-1}h_a(t)$, thus $-(e^{ag_a^{-1}(t)} - 1)^{-1} \geq -(e^{h_a(t)} - 1)^{-1}$. \square

EXAMPLE. Consider $f(x) = (\Gamma(.5))^{-1}x^{-.5}e^{-x}$, $x > 0$; this is the $\Gamma(.5, 1) = \chi_1^2/2$ distribution, which is DFR ([5], page 378). We compute the moment based bounds for $M(t)$, using Theorem 4 and Corollary 1, to illustrate the method, and to see how much the DFR bounds improve upon the IMRL bounds in this case. Using the recursive formula (4) we compute $c_0 = \frac{1}{2}$, $c_1 = c_2 = \frac{1}{8}$, $c_3 = \frac{15}{64}$, $c_4 = \frac{21}{32}$, $c_5 = \frac{315}{128}$, $c_6 = \frac{1485}{128}$. Next $v_{-1} = 0$, $v_0 = \frac{1}{4}$, $v_1 = 1$, $v_2 = \frac{15}{8}$, $v_3 = \frac{14}{5}$, $v_4 = \frac{15}{4}$, $v_5 = \frac{33}{7}$, $v_6 = \frac{91}{16}$. Denoting the lower bound given in Theorem 4 by $B(t) = U(t) - \inf_i c_i t^{-i}$ we obtain:

$$\begin{aligned}
 B(t) &= 2t + 1, & 0 \leq t \leq \frac{1}{4} \\
 &= 2t + \frac{3}{2} - (8t)^{-1}, & \frac{1}{4} \leq t \leq 1 \\
 &= 2t + \frac{3}{2} - (8t^2)^{-1}, & 1 \leq t \leq \frac{15}{8} \\
 &= 2t + \frac{3}{2} - \left(\frac{15}{64t^3}\right), & \frac{15}{8} \leq t \leq \frac{14}{5} \\
 &= 2t + \frac{3}{2} - \left(\frac{21}{32t^4}\right), & \frac{14}{5} \leq t \leq \frac{15}{4} \\
 &= 2t + \frac{3}{2} - \left(\frac{315}{128t^5}\right), & \frac{15}{4} \leq t \leq \frac{33}{7} \\
 &= 2t + \frac{3}{2} - \left(\frac{1485}{128t^6}\right), & \frac{33}{7} \leq t \leq \frac{91}{16},
 \end{aligned}$$

The lower bound, given in Corollary 1, which we denote by $B^*(t) = U(t) - \inf_i c_i^* t^{-i}$ is similarly computed.

The table below gives a few values of t along with the corresponding intervals $[B(t), U(t)]$ and $[B^*(t), U(t)]$ for $M(t)$.

t	$(B(t), U(t))$	$(B^*(t), U(t))$
.1	[1.2, 1.7]	[1.2, 1.7]
.5	[2.25, 2.5]	[2.375, 2.5]
1	[3.375, 3.5]	[3.444, 3.5]
1.5	[4.444, 4.5]	[4.475, 4.5]
2	[5.471, 5.5]	[5.488, 5.5]
3	[7.492, 7.5]	[7.497, 7.5]
4	[9.4976, 9.5]	[9.4990, 9.5]
5	[11.49926, 11.5]	[11.49971, 11.5]

6. Comments and additions.

(1) For IMRL distributions we have the representation (Theorem 1) $X'_0 = \sum_1^N X_i$, where $X_i \sim F$, $X'_0 \sim G$, $G(x) = \mu_1^{-1} \int_0^x \bar{F}(y) dy$, and N is a stopping time. A simple argument shows that $\Pr(N = 1) > 0$. Since G is absolutely continuous, it follows that an IMRL distribution must have an absolutely continuous component. I do not know whether, aside from an atom at $\{0\}$, an IMRL distribution can have a singular component.

(2) If the above mentioned representation is valid for distribution F , with N a stopping time, then a renewal process with interarrival time distribution F will

enjoy many of the properties which were derived for IMRL renewal processes. I do not know whether F IMRL is a necessary condition for such a representation to hold.

(3) By Theorem 2(i), if F is IMRL and $\mu_2 < \infty$, then $L(t) - L(0)$ is the distribution function of a finite positive measure over the Borel sets in $(0, \infty)$. If F is DFR then, in addition, L has a decreasing density (Theorem 3 (ii)). I suspect that L has an improvement with age property for F IMRL, with a strengthened version for F DFR. This is based on the behavior of the moments of L in a few special cases which I have worked out.

(4) If we imitate the construction of Theorem 1 for increasing failure rate (IFR) distributions we obtain the following. The ordinary and delayed renewal processes alternate renewals until a random time at which their renewal epochs coincide; then all future renewal epochs coincide. So far this representation has not yielded much in the way of special properties for DMRL and IFR renewal processes. An example of Berman ([7] page 429) shows that F IFR does not imply $m(t)$ increasing, nor $M(t) - t/\mu_1$ decreasing, nor $EZ(t)$ decreasing. Thus the obvious analogues of the DFR and IMRL results of this paper do not hold for IFR and DMRL distributions.

(5) Identities (4) and (5) may be of independent renewal theory interest. They hold for distributions on $[0, \infty)$ possessing the required moments and having the property that some convolution has an absolutely continuous component. To prove this one would follow the argument of Theorem 4, invoking results of W. L. Smith [25] to justify both $\lim t^k L(t) = 0$, and the applicability of the key renewal argument.

(6) The results $c_i \geq 0$ and $v_i \uparrow$ (Theorem 4) are inequalities among the moments of IMRL distributions which may be of independent interest.

REFERENCES

- [1] BARLOW, R. E. (1965). Bounds on integrals with applications to reliability problems. *Ann. Math. Statist.* **36** 565–574.
- [2] BARLOW, R. E. and MARSHALL, A. W. (1964a). Bounds for distributions with monotone hazard rate, I. *Ann. Math. Statist.* **35** 1237–1257.
- [3] BARLOW, R. E. and MARSHALL, A. W. (1964b). Bounds for distributions with monotone hazard rate, II. *Ann. Math. Statist.* **35** 1258–1274.
- [4] BARLOW, R. E. and PROSCHAN, F. (1964). Comparison of replacement policies, and renewal theory implications. *Ann. Math. Statist.* **35** 577–589.
- [5] BARLOW, R. E. and PROSCHAN, F. (1975). *Statistical Theory of Reliability and Life Testing: Probability Models*. Holt, Rinehart and Winston, New York.
- [6] BARLOW, R. E., MARSHALL, A. W. and PROSCHAN, F. (1963). Properties of probability distributions with monotone hazard rate. *Ann. Math. Statist.* **34** 375–389.
- [7] BERMAN, M. (1978). Regenerative multivariate point processes. *Advances in Appl. Probability* **10** 411–430.
- [8] BRYSON, M. C. and SIDDIQUI, M. M. (1969). Some criteria for aging. *J. Amer. Statist. Assoc.* **64** 472–483.
- [9] DALEY, D. J. (1968). Stochastically monotone Markov chains. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **10** 305–317.

- [10] ESARY, J. D., MARSHALL, A. W. and PROSCHAN, F. (1973). Shock models and wear processes. *Ann. Probability* **1** 627–649.
- [11] FELLER, W. (1966). *An Introduction to Probability Theory and its Applications, Volume II*. Wiley, New York.
- [12] HAINES, A. L. and SINGPURWALLA, N. D. (1974). Some contributions to the stochastic characterization of weak. In *Reliability and Biometry*. (F. Proschan and R. J. Serfling, eds.), 47–80, SIAM 1974.
- [13] KALMYKOV, G. I. (1962). On the partial ordering of one-dimensional Markov processes. *Theor. Probability Appl.* **7** 456–459.
- [14] KEILSON, J. (1975). Systems of independent Markov components and their transient behavior. In *Reliability and Fault Free Analysis*. (R. E. Barlow, J. B. Fussel and N. D. Singpurwalla, eds.) 351–364, SIAM, Philadelphia.
- [15] KEMPERMAN, J. H. B. (1977). Generalized dilations and other moment problems for a measure with given marginals. Unpublished manuscript, Univ. Rochester.
- [16] LEHMAN, E. L. (1959). *Testing Statistical Hypotheses*. Wiley, New York.
- [17] MARSHALL, A. W. and PROSCHAN, F. (1972). Classes of distributions applicable in replacement, with renewal theory implications. In *Proc. Sixth Berkeley Symp. Math. Statist. Probability I* edited by (L. LeCam, J. Neyman and E. L. Scott, eds), 395–415 Univ. California Press.
- [18] MARSHALL, K. T. (1973). Linear bounds on the renewal function. *SIAM J. Appl. Math.* **25** 245–250.
- [19] MILLER, D. R. (1976). Almost sure comparisons of renewal processes and Poisson processes. Univ. Missouri technical report.
- [20] O'BRIEN, G. L. (1975). The comparison method for stochastic processes. *Ann. Probability* **3** 80–88.
- [21] PLEDGER, G. and PROSCHAN, F. (1973). Stochastic comparisons of random processes. *J. Appl. Probability*. **10** 572–585.
- [22] PYKE, R. (1968). Applications of almost surely convergent constructions of weakly convergent processes. In *Proc. Internat. Symp. Probability Inform. Theor.* Springer, Berlin.
- [23] ROSS, S. M. (1970). *Applied Probability Models with Optimization Applications*. Holden Day, San Francisco.
- [24] SMITH, W. L. (1954). Asymptotic renewal theorems. *Proc. Roy. Soc. Edinburgh Sect. A* **64** 9–48.
- [25] SMITH, W. L. (1958). Renewal theory and its ramifications. *J. Roy. Statist. Soc., Ser. B* **20** 243–302.
- [26] STONE, C. R. (1966). On absolutely continuous components and renewal theory. *Ann. Math. Statist.* **37** 271–275.

DEPARTMENT OF STATISTICS
FLORIDA STATE UNIVERSITY
TALLAHASSEE, FLORIDA 32306