

GEOMETRY OF CANONICAL SELF-SIMILAR TILINGS

ERIN P.J. PEARSE AND STEFFEN WINTER

ABSTRACT. We give several different geometric characterizations of the situation in which the parallel set F_ε of a self-similar set F can be described by the inner ε -parallel set $T_{-\varepsilon}$ of the associated canonical tiling \mathcal{T} , in the sense of [15]. For example, $F_\varepsilon = T_{-\varepsilon} \cup C_\varepsilon$ if and only if the boundary of the convex hull C of F is a subset of F , or if the boundary of E , the unbounded portion of the complement of F , is the boundary of a convex set. In the characterized situation, the tiling allows one to obtain a tube formula for F , i.e., an expression for the volume of F_ε as a function of ε . On the way, we clarify some geometric properties of canonical tilings.

Motivated by the search for tube formulas, we give a generalization of the tiling construction which applies to all self-affine sets F having empty interior and satisfying the open set condition. We also characterize the relation between the parallel sets of F and these tilings.

1. Introduction. As the basic object of our study is a self-affine system and its attractor, the associated self-affine set, we begin by defining these terms.

Definition 1.1. For $j = 1, \dots, N$, let $\Phi_j : \mathbf{R}^d \rightarrow \mathbf{R}^d$ be an affine contraction whose eigenvalues λ all satisfy $0 < |\lambda| < 1$. Then $\{\Phi_1, \dots, \Phi_N\}$ is a *self-affine iterated function system*.

Definition 1.2. A *self-similar system* is a self-affine system for which each mapping is a similitude, i.e.,

2010 AMS *Mathematics subject classification*. Primary 28A80, 28A75, 52A20, 52C22, Secondary 52A38, 53C65, 51M25, 49Q15, 60K05, 54F45.

Keywords and phrases. Iterated function system, parallel set, fractal, complex dimensions, zeta function, tube formula, Steiner formula, renewal theorem, convex ring, inradius, Euler characteristic, Euler number, self-affine, self-similar, tiling, curvature measure, generating function, fractal string.

The work of the first author was partially supported by the University of Iowa Department of Mathematics NSF VIGRE grant DMS-0602242. The work of the second author was partially supported by Cornell University and a grant from the German Academic Exchange Service (DAAD).

Received by the editors on February 3, 2009, and in revised form on May 18, 2009.

DOI:10.1216/RMJ-2012-42-4-1327 Copyright ©2012 Rocky Mountain Mathematics Consortium

$$(1.1) \quad \Phi_j(x) := r_j A_j x + a_j,$$

where for $j = 1, \dots, N$, we have $0 < r_j < 1$, $a_j \in \mathbf{R}^d$ and $A_j \in O(d)$, the orthogonal group of d -dimensional Euclidean space \mathbf{R}^d . The numbers r_j are referred to as the *scaling ratios* of $\{\Phi_1, \dots, \Phi_N\}$.

Let F be the self-affine set generated by the mappings Φ_1, \dots, Φ_N , i.e., the unique (nonempty and compact) set satisfying $\Phi(F) = F$ where Φ is the set mapping

$$(1.2) \quad \Phi := \bigcup_{j=1}^N \Phi_j.$$

The existence and uniqueness of set F is ensured by the classic results of Hutchinson in [6]. It is shown in [15] that, when a self-affine system satisfies the tileset condition (TSC) and the nontriviality condition (given here in Definitions 2.4 and 2.9, respectively), then there is a natural tiling of the convex hull $C = [F]$. That is, $\{\Phi_1, \dots, \Phi_N\}$ generates a decomposition of C into open sets $\mathcal{T} = \{R^n : n \in \mathbf{N}\}$, in the sense that

$$\overline{\bigcup_{n=1}^{\infty} R^n} = C \quad \text{and} \quad R^n \cap R^m = \emptyset \text{ for } n \neq m,$$

cf. Definition 3.1. One of our main objectives in this paper is to explore the consequences of these two conditions and characterize some properties of the tilings. In particular, we clarify the relationship between the tileset condition as defined in [15] and the open set condition; in fact, the latter is implied by the former, cf. Proposition 2.5. The nontriviality condition forbids self-similar sets with convex attractors, like the square or interval. Additionally, we show in Proposition 2.10 that the nontriviality condition ensures the existence of tiles in the tiling construction. Under TSC, nontriviality is also equivalent to F having empty interior, see Proposition 2.11. We discuss the boundary of the tiling and its Hausdorff dimension in Proposition 3.5 and Remark 5.12.

In [15], it was noted that tiling \mathcal{T} constitutes the bulk of the nontrivial portion of the complement of F and, consequently, that one may be able

to study the ε -parallel sets (or ε -neighborhoods) of F by considering the inner ε -neighborhoods of the tiling. By the ε -parallel set A_ε of a set $A \subseteq \mathbf{R}^d$, we mean all points not in the interior of A but with distance at most ε to A . (Note that our usage of A_ε differs from the usual one, where the interior points of A are included, but it is more convenient for our purposes.) Similarly, the inner ε -parallel set $A_{-\varepsilon}$ consists of points of the closure of A within distance ε of $\text{bd } A$, see Definition 4.1 for the details. We determine the conditions under which the tiling allows a (almost disjoint, cf. (4.3)) decomposition of F_ε of the following form:

$$(1.3) \quad F_\varepsilon = T_{-\varepsilon} \cup C_\varepsilon.$$

Here $T := \bigcup R^n$ denotes the union of the tiles of \mathcal{T} . In Theorems 4.4 and 4.9, we give eight equivalent conditions which characterize this state of affairs; these results will be collectively referred to as the Compatibility Theorem.

In Section 5 we generalize the tiling construction introduced in [15] and discussed in earlier sections of the present paper. Specifically, we replace the tilename condition with the less restrictive open set condition (see Definition 2.1) and replace the convex hull with an arbitrary feasible open set. Finally, in Section 6 we extend the Compatibility Theorem to the generalized self-similar tilings developed in Section 5. For instance, the tiling generated from a feasible set O is compatible if and only if $\text{bd } \overline{O} \subseteq F$.

Compatibility allows one to employ the tiling to obtain a tube formula for F , and this is the driving motivation for the current paper. By a *tube formula* of a set $A \subset \mathbf{R}^d$, we mean an expression which gives the Lebesgue volume $V(A_\varepsilon)$ of A_ε as a function of ε . Such objects are of considerable interest in spectral geometry; see [8–10], as well as the more general references [5, 18, 20]. In convex geometry, tube formulas are better known as Steiner formulas:

$$(1.4) \quad V(A_\varepsilon) = \sum_{k=0}^{d-1} \varepsilon^{d-k} \kappa_{d-k} C_k(A)$$

For compact convex subsets A of \mathbf{R}^d , $V(A_\varepsilon)$ is a polynomial in ε , and the coefficients $C_k(A)$ are called *total curvatures* or *intrinsic volumes*;

these are important geometric invariants of set A and are related to the integrals of mean curvature, provided the boundary of A is sufficiently smooth. A polynomial expansion similar to (1.4) is known for sets of *positive reach* [4]. Also for polyconvex sets (finite unions of convex sets) and certain unions of sets with positive reach, polynomial expansions are known. However, in these latter cases, the polynomial describes a “weighted” parallel volume which counts points in the parallel sets with different multiplicities given by an index function, cf. [17, 22].

For more singular sets like fractals one cannot expect such polynomial behavior. Tube formulas for subsets A of \mathbf{R} have been extensively studied, see [11] and the references therein, and they have been related to the theory of complex dimensions. Here the tube formulas typically take the form of an infinite sum. In [8] a first attempt was made to generalize this theory to higher dimensions, and tube formulas have been obtained for so called *fractal sprays*. The theory is developed further in [10].

A self-similar tiling \mathcal{T} is a certain kind of fractal spray, and so this theory applies. One can associate a *geometric zeta function* $\zeta_{\mathcal{T}} : \mathbf{C} \times (0, \infty) \rightarrow \mathbf{C}$ which encodes all the geometric information of \mathcal{T} . The *complex dimensions* \mathcal{D} of the tiling are the poles of $\zeta_{\mathcal{T}}$. Then, for $T = \bigcup R^n$, a tube formula (describing the inner ε -parallel volume of the union of the tiles) of the following form holds

$$V(T_{-\varepsilon}) = \sum_{w \in \mathcal{D}} \operatorname{res}(\zeta_{\mathcal{T}}(s, \varepsilon); s = w), \quad \varepsilon > 0,$$

see [8–10] for details. Under mild additional assumptions, a factor ε^{d-w} can be separated from each residue, and the formula takes a form very similar to the Steiner formula:

$$(1.5) \quad V(T_{-\varepsilon}) = \sum_{w \in \mathcal{D}} \varepsilon^{d-w} C_w(\mathcal{T}), \quad \varepsilon > 0,$$

with coefficients C_w independent of ε . Just as in (1.4), it turns out that \mathcal{D} always contains $\{0, 1, 2, \dots, d-1\}$.

In [21], the author develops a theory of fractal curvatures: a family of geometric invariants $C_k^f(F)$, $k = 0, 1, \dots, d$. The fractal Euler characteristic C_0^f was introduced in [12], and C_d^f coincides with the

Minkowski content. These curvatures are defined for certain self-affine fractals and provide a fractal analogue of the coefficients $C_k(A)$ mentioned in (1.4). Indeed, they are even localizable as *curvature measures* in the same way that the coefficients of the Steiner formula are, cf. [18]. However, the fractal analogue of the Steiner formula is absent from the context of [21], and it is a major impetus for this paper to establish such a link. In particular, the methods of the present paper and the theory of fractal curvatures are both applicable when the envelope (introduced in Definition 4.5) is polyconvex. It remains to be determined if the coefficients $C_w(\mathcal{T})$ appearing in (1.5) can thus be interpreted as curvatures and, if so, if they are compatible with the theory of [12, 21]. The Compatibility Theorems of the present paper describe how parallel sets of the tilings are related to parallel sets of F . For compatible sets F , a tube formula for F is obtained from decomposition (1.3) as the sum of the (inner) tube formula of an appropriate tiling \mathcal{T} and a “trivial” part, describing the “outer” parallel volume of the tiled set, i.e., the convex hull C of F :

$$(1.6) \quad V(F_\varepsilon) = V(T_{-\varepsilon}) + V(C_\varepsilon).$$

Here, $V(T_{-\varepsilon})$ is as in (1.5) and $V(C_\varepsilon)$ is as in (1.4). A similar formula holds for generalized tilings when a compatible feasible set exists. The Compatibility Theorems characterize the situation in which decomposition (1.6) holds; they also show the limitations of this approach. We illustrate this with suitable counterexamples (see Proposition 6.3).

2. Tileset condition and nontriviality condition. The open set condition is a classical separation condition for the study of self-similarity, cf. [3].

Definition 2.1. A self-affine system $\{\Phi_1, \dots, \Phi_N\}$ satisfies the *open set condition* (OSC) if and only if there is a nonempty open set $O \subseteq \mathbf{R}^d$ such that

$$(2.1) \quad \Phi_j(O) \subseteq O, \quad j = 1, 2, \dots, N,$$

$$(2.2) \quad \Phi_j(O) \cap \Phi_k(O) = \emptyset \text{ for } j \neq k.$$

In this case, O is called a *feasible open set* for F .

We denote the *convex hull* of a set $A \subseteq \mathbf{R}^d$ (that is, the smallest convex set containing A), by $[A]$. In particular, we denote the convex hull of the attractor F of a system $\{\Phi_1, \dots, \Phi_N\}$ by $C = [F]$.

Remark 2.2. F is always assumed to be embedded in the smallest possible ambient space, i.e., $\mathbf{R}^d = \text{aff } F$ is the affine hull of F , and thus C is of full dimension.

It was a crucial observation in [15] that the convex hull satisfies $\Phi_j(C) \subseteq C$, which implies the nestedness of C under iteration, cf. [15, Theorem 5.1, page 3162]:

Proposition 2.3. $\Phi^{k+1}(C) \subseteq \Phi^k(C) \subseteq C$, for $k = 1, 2, \dots$.

The last proposition is reminiscent of [6, subsection 5.2 (3)]. We recall the conditions introduced in [15] to ensure the existence of a canonical tiling of the convex hull of F , namely, the *tileset condition* and the *nontriviality condition*.

Definition 2.4. A self-affine system $\{\Phi_1, \dots, \Phi_N\}$ (or its attractor F) satisfies the *tileset condition* (TSC) if and only if it satisfies OSC with $\text{int } C$ as a feasible open set.

Proposition 2.5. F satisfies TSC if and only if

$$(2.3) \quad \text{int } \Phi_j(C) \cap \text{int } \Phi_k(C) = \emptyset \quad \text{for } j \neq k.$$

Proof. The if-part is obvious; for the only-if-part, apply Proposition 2.3. \square

Common examples satisfying TSC (and NTC, defined just below in Definition 2.9) include the Sierpinski gasket and carpet, the Cantor set, the Koch snowflake curve and the Menger sponge. It is obvious from the definition that TSC implies OSC. The following examples demonstrate that the converse is not true.




FIGURE 1. A self-similar system which satisfies the open set condition but not the tileset condition; see Example 2.7. The attractor in this example tiles all of \mathbf{R}^2 .

Example 2.6. Let $F \subseteq \mathbf{R}$ be the self-similar set generated by the system $\{\Phi_1, \Phi_2, \Phi_3\}$ where the mappings $\Phi_j : \mathbf{R} \rightarrow \mathbf{R}$ are given by $\Phi_1(x) = (1/3)x$, $\Phi_2(x) = (1/3)x + (2/3)$ and $\Phi_3(x) = (1/9)x + (1/9)$, respectively. Let $O = (0, 1/3) \cup (2/3, 1)$. Clearly, O is a feasible open set for the OSC for F since the images $\Phi_1 O = (0, 1/9) \cup (2/9, 1/3)$, $\Phi_2 O = (2/3, 7/9) \cup (8/9, 1)$ and $\Phi_3 O = (1/9, 4/27) \cup (5/27, 2/9)$ are subsets of O and pairwise disjoint. Thus F satisfies the OSC. On the other hand, the TSC is not satisfied. The convex hull of F is $C = [0, 1]$, and the sets $\Phi_1 C = [0, 1/3]$ and $\Phi_3 C = [1/9, 2/9]$ strongly overlap.

Note that it is even possible for a self-affine set to satisfy the *strong separation condition* (that is, that the images $\Phi_j(F)$ are pairwise disjoint) but not the tileset condition. An example of such a set is obtained, for instance, by replacing the mapping Φ_3 in the above example with the mapping $\Phi'_3(x) = (1/27)x + (4/27)$. The images $\Phi_1 F'$, $\Phi_2 F'$ and $\Phi'_3 F'$ of the corresponding self-similar set F' are pairwise disjoint, while the images of its convex hull have intersecting interiors.

Example 2.7. Consider a system of three similarity mappings, each with scaling ratio $1/\sqrt{3}$ and a clockwise rotation of $\pi/2$. The mappings are illustrated in Figure 1. They form a system which satisfies the open set condition (simply take the interior of the attractor) but not the tileset condition. On the right, the attractor has been shaded for clarity; the dark overlay indicates the intersection of the convex hulls of two first level images of the attractor.

Remark 2.8. After a talk on the topic of the present paper at the conference, *Fractal Geometry and Stochastics IV*, at Greifswald,

Kenneth Falconer asked the following question: “Is there an easy way to decide whether, for a given self-similar set F satisfying OSC, there is a feasible open set that is convex?” The results in this paper provide the following answer.

There is a convex feasible open set for F if and only if F satisfies the tileset condition, i.e., if and only if the interior of the convex hull C of F is feasible. To see this, assume that a feasible open set O exists that is convex. Then its closure \overline{O} is closed and convex and satisfies $F \subseteq \overline{O}$ (cf. Proposition 5.1). It follows that $C \subseteq \overline{O}$ (since the convex hull is the intersection of all closed convex sets containing F) and thus $\text{int } C \subseteq O$. But this implies $\Phi_i(\text{int } C) \cap \Phi_j(\text{int } C) \subseteq \Phi_i O \cap \Phi_j O = \emptyset$. Hence, F satisfies the tileset condition by Proposition 2.5. Thus, it is sufficient to check whether the interior of the convex hull is feasible to decide the above question.

Definition 2.9. We say that $\{\Phi_1, \dots, \Phi_N\}$ satisfies the *nontriviality condition* (NTC) if and only if its attractor F is not convex.

The nontriviality condition is, besides the TSC, the second necessary condition to ensure the existence of a canonical self-affine tiling for F . Proposition 2.10 shows that nontriviality is precisely the condition that ensures the generators of the tiling exist, as will be apparent from Definition 3.3. The following proposition shows that the present usage of “nontriviality” agrees with that of [15].

Proposition 2.10. *A self-affine system $\{\Phi_1, \dots, \Phi_N\}$ is nontrivial if and only if the images $\Phi_j(C)$ of C do not cover $\text{int}(C)$, i.e., the convex hull C satisfies*

$$(2.4) \quad \text{int}(C) \not\subseteq \Phi(C).$$

Proof. First observe that (2.4) is equivalent to

$$(2.5) \quad C \not\subseteq \Phi(C).$$

Indeed, the implication (2.4) \Rightarrow (2.5) is obvious. Conversely, if (2.5) holds, then $C \cap \Phi(C)^c \neq \emptyset$. Hence, some point $x \in C \cap \Phi(C)^c$

exists and, since $\Phi(C)^{\mathbb{G}}$ is open, some $\delta > 0$ such that the ball $B(x, \delta)$ is contained in $\Phi(C)^{\mathbb{G}}$. Now, since C is convex and thus the closure of its interior ($\dim C = d$, cf. Remark 2.2), there is a point $y \in B(x, \delta) \cap \text{int}(C)$. Hence, $\text{int}(C) \cap \Phi(C)^{\mathbb{G}}$ is nonempty, implying (2.4).

Recall that $\Phi(C) \subseteq C$ by nestedness (Proposition 2.3). Therefore, if (2.4) fails, its equivalence with (2.5) immediately implies $C = \Phi(C)$. By uniqueness of the invariant set (with respect to Φ), this means that F is equal to its convex hull C . Obviously, if the nontriviality condition is satisfied, then F is not equal to its convex hull. \square

For self-affine sets satisfying TSC, we give a different characterization of nontriviality. $F \subset \mathbf{R}^d$ is trivial if and only if it has nonempty interior.

Proposition 2.11. *Let F be a self-affine set satisfying TSC. Then F is nontrivial if and only if $\text{int } F = \emptyset$.*

Proof. If F is nontrivial, then the set $T_0 := \text{int}(C \setminus \Phi(C))$ is nonempty, but $T_0 \cap F = \emptyset$, since $F \subseteq \Phi(C)$. Observe that TSC implies $\Phi_i(\text{int } C) \cap \Phi_j(F) = \emptyset$ for $i \neq j$. Therefore, $\Phi_i(T_0) \cap F \subseteq \Phi_i(T_0) \cap \Phi_i(F) = \Phi_i(T_0 \cap F) = \emptyset$, and so $\Phi(T_0) \cap F = \emptyset$. By induction, we get $\Phi^k(T_0) \cap F = \emptyset$ for $k = 0, 1, 2, \dots$. Now let $x \in F$. Since, by the contraction principle, $d_H(F, \Phi^k(\overline{T_0})) = d_H(\Phi^k(F), \Phi^k(\overline{T_0})) \rightarrow 0$ as $k \rightarrow \infty$, a sequence $x_k \rightarrow x$ exists with $x_k \in \Phi^k(\overline{T_0}) = \overline{\Phi^k(T_0)}$. For each x_k there are points in $\Phi^k(T_0)$ arbitrarily close to x_k . Hence, x cannot lie in the interior of F .

For the converse, if F is trivial, then it is convex by Proposition 2.10. In view of Remark 2.2, $\text{int } F \neq \emptyset$. \square

Remark 2.12. The fact that self-affine sets satisfying TSC and NTC have empty interior was used implicitly in [15] without mention. Proposition 2.11 clarifies that this was justified.

Combining Propositions 2.10 and 2.11, we infer that, for self-affine sets satisfying TSC, nonempty interior means convexity. For the special case of self-similar sets, convexity is also equivalent to having full

dimension. This follows from a result of Schief [17, Corollary 2.3] stating that, for self-similar sets $F \subseteq \mathbf{R}^d$ satisfying OSC, $\dim_H F = d$ implies that F has interior points.

Corollary 2.13. *Let $F \subseteq \mathbf{R}^d$ be a self-affine set satisfying TSC. If F has Hausdorff dimension strictly less than d , then F is nontrivial. Moreover, if F is self-similar, then the converse also holds.*

Proof. If F is trivial, then, by Proposition 2.11, F has a nonempty interior which implies $\dim_H F = d$. Now let F be self-similar and satisfy TSC. Assume $\dim_H F = d$. Since TSC implies OSC, by [18, Theorem 2.2 and Corollary 2.3], F has nonempty interior. Therefore, by Proposition 2.11, F is trivial. \square

See also Proposition 5.4 and Corollary 5.6 for analogues of Proposition 2.11 and Corollary 2.13 in the more general context of OSC.

3. Canonical self-affine tilings. Let $\{\Phi_1, \dots, \Phi_N\}$ be a self-affine system with attractor F satisfying both TSC and NTC. In this section, we recall the construction of the so-called *canonical self-affine tiling* of the convex hull C of F introduced in [15, Section 3]. On the way, we prove some foundational results concerning open tilings, thereby clarifying a couple of technical points which were left vague in [15].

Definition 3.1. A sequence $\mathcal{A} = \{A^i\}_{i \in \mathbf{N}}$ of pairwise disjoint open sets $A^i \subseteq \mathbf{R}^d$ is called an *open tiling* of a set $B \subseteq \mathbf{R}^d$ (or a *tiling of B by open sets*) if and only if

$$\overline{B} = \overline{\bigcup_{i=1}^{\infty} A^i}.$$

Sets A^i are called the *tiles*.

Note that Definition 3.1 is weaker than the usual definition of a tiling: no local finiteness is assumed. In other words, a given compact set may be intersected by infinitely many of the tiles. The case that B is tiled by a finite number $m \in \mathbf{N}$ of tiles A^1, \dots, A^m is included by setting $A^i := \emptyset$ for $i > m$. Since here we are more interested in open tilings

of B by an infinite number of sets, the tiles A^i are usually assumed to be nonempty. Also note that each sequence $\{A^i\}$ of disjoint open sets is an open tiling of some set $B \subseteq \mathbf{R}^d$ but this set is not uniquely determined. For instance, if $\{A^i\}$ is an open tiling of B , then it is also an open tiling of $\text{int } B$ and of \overline{B} . Sequence $\{A^i\}$ only determines the closure of B uniquely.

The following observation regarding the boundaries of tiles will be useful in the sequel. In particular, it is used repeatedly in the proof of Theorem 4.4, a central result of this paper. Let $\{A^i\}$ be an open tiling of a set B . Denote by $A = \bigcup_{i=1}^{\infty} A^i$ the union of the tiles. Since the sets A^i are open, A is open as well. The boundary of A (defined in the usual way as $\text{bd } A = \overline{A} \cap \overline{A^c}$ or, since A is open, equivalently by $\text{bd } A = \overline{A} \setminus A$) is characterized by the tiles as follows:

Lemma 3.2.

$$\text{bd } A = \overline{\bigcup_i \text{bd } A^i}.$$

Proof. (\subseteq). Let $x \in \text{bd } A$. Then a sequence $\{x_k\}_{k=1}^{\infty}$ exists in $A = \bigcup_i A^i$ converging to x as $k \rightarrow \infty$. Using $\{x_k\}$, we construct a sequence $\{x'_k\}$ in $\bigcup_i \text{bd } A^i$ in the following way. For each x_k there is a (unique) index $n(k) \in \mathbf{N}$ such that $x_k \in A^{n(k)}$. Since $A^{n(k)}$ is open, $x \notin A^{n(k)}$. Now let x'_k be any point of the set $[x, x_k] \cap \text{bd } A^{n(k)}$, where $[x, x_k]$ is the (closed) line segment between x and x_k . Such a point exists, since $x \in (A^{n(k)})^c$ (but it may not be unique). Then, clearly, $\{x'_k\}$ is a sequence in $\bigcup_i \text{bd } A^i$. Moreover, $x'_k \rightarrow x$ as $k \rightarrow \infty$, since $x_k \rightarrow x$ and $|x - x'_k| < |x - x_k|$. But this implies that $x \in \overline{\bigcup_i \text{bd } A^i}$, proving the inclusion from left to right.

(\supseteq). For a proof of the reversed inclusion, let $x \in \overline{\bigcup_i \text{bd } A^i}$. Then a sequence $\{y_k\} \subseteq \bigcup_i \text{bd } A^i$ exists such that $y_k \rightarrow x$ as $k \rightarrow \infty$. The existence of this sequence (and disjointness of the tiles A^i) imply immediately that $x \notin A$, since an interior point of A cannot be an accumulation point of a sequence in A^c . Furthermore, each y_k is an element of at least one of the sets $\text{bd } A^i$. Let $n(k)$ be an index such that $y_k \in \text{bd } A^{n(k)}$. For each y_k , we find points in $A^{n(k)}$ arbitrarily close to y_k . Choose $y'_k \in A^{n(k)}$ such that $|y_k - y'_k| < 1/k$. Then $|x - y'_k| \leq |x - y_k| + |y_k - y'_k| < |x - y_k| + (1/k) \rightarrow 0$ as $k \rightarrow \infty$. Thus,

$y'_k \rightarrow x$. Recalling that $y'_k \in A^{n(k)}$ and thus $\{y'_k\} \subseteq \bigcup_i A^i = A$, we conclude that $x \in \bar{A}$. Together with $x \notin A$, this yields $x \in \bar{A} \setminus A = \text{bd } A$, completing the proof. \square

Let

$$(3.1) \quad W := \bigcup_{k=0}^{\infty} \{1, \dots, N\}^k$$

denote the set of all finite words formed by the alphabet $\{1, \dots, N\}$. For any word $w = w_1 w_2 \dots w_n \in W$, let $\Phi_w = \Phi_{w_1} \circ \Phi_{w_2} \circ \dots \circ \Phi_{w_n}$. In particular, if $w \in W$ is the *empty word*, then $\Phi_w = \text{Id}$.

Denote by G_1, G_2, \dots the connected components of the open set $T_0 := \text{int}(C \setminus \Phi(C))$; $T_0 = \bigcup_{q \in Q} G_q$. The index set $Q \subseteq \mathbf{N}$ may be infinite, but, since T_0 is open, the number of its connected components is certainly at most countable.

Definition 3.3. The *canonical self-affine tiling* associated with $\{\Phi_1, \dots, \Phi_N\}$ (or with F) is

$$(3.2) \quad \mathcal{T} = \{\Phi_w(G_q) : w \in W, q \in Q\}.$$

The open subsets G_q of C are called the *generators* of \mathcal{T} . It is shown in [15, Theorem 5.16, page 3167] that \mathcal{T} is an open tiling of $C = [F]$ in the sense of Definition 3.1, i.e., the sets $\Phi_w(G_q)$ are pairwise disjoint and

$$C = \overline{\bigcup_{R \in \mathcal{T}} R}.$$

Write $T = \bigcup_{R \in \mathcal{T}} R$ for the union of the tiles of \mathcal{T} and $\text{bd } T$ for the boundary of this set. Clearly, T is open, $\bar{T} = C$ and so $\text{bd } T = \bar{T} \setminus T = C \setminus T$. By Lemma 3.2, we have

$$(3.3) \quad \text{bd } T = \overline{\bigcup_{R \in \mathcal{T}} \text{bd } R}.$$

Note that the closure in representation (3.3) cannot be omitted. One has $F \subseteq \text{bd } T$ (cf. Lemma 3.4), while $F \not\subseteq \bigcup_{R \in \mathcal{T}} \text{bd } R$. If the Hausdorff

dimension $\dim_H F$ is strictly greater than $d - 1$, then taking the closure leads to a jump of dimension. More precisely, one has the equality $\dim_H \text{bd } T = \max\{\dim_H F, d - 1\}$, as is shown in Proposition 3.5. For the proof, it is convenient to work with a slight variation of the tiling described above: it is possible to consider the set T_0 as the generator of a tiling, instead of its connected components G_q . This point of view leads to a different tiling $\mathcal{T}' := \{\Phi_w(T_0) : w \in W\}$ of C whose tiles are not necessarily connected. It is easily seen that \mathcal{T}' is also an open tiling of C in the sense of Definition 3.1. Moreover, for each tile $\Phi_w(T_0) \in \mathcal{T}'$, $\{\Phi_w(G_q) : q \in Q\}$ is an open tiling of $\Phi_w(T_0)$. If $T' = \bigcup_{R \in \mathcal{T}'} R$ is the union of the tiles, then by two applications of Lemma 3.2, the boundaries of both tilings coincide:

$$\text{bd } T' = \overline{\bigcup_{R' \in \mathcal{T}'} \text{bd } R'} = \overline{\bigcup_{w \in W} \text{bd } \Phi_w T_0} = \overline{\bigcup_{w \in W} \bigcup_{q \in Q} \text{bd } \Phi_w G_q} = \text{bd } T.$$

Lemma 3.4.

$$\text{bd } T = F \cup \bigcup_{R \in \mathcal{T}'} \text{bd } R.$$

Proof. (\supseteq). From Lemma 3.2, we have $\bigcup_{R \in \mathcal{T}'} \text{bd } R \subseteq \text{bd } T' = \text{bd } T$. For the inclusion $F \subseteq \text{bd } T$, note that \mathcal{T} is an open tiling of C , and thus $F \subseteq C = \overline{\bigcup_{R \in \mathcal{T}} R} = \overline{T}$. But, by [15, Theorem 5.16, page 3167], $R \cap F = \emptyset$ for all $R \in \mathcal{T}$, i.e., $F \cap T = \emptyset$. Thus, $F \subseteq \overline{T} \setminus T = \text{bd } T$.

(\subseteq). Let $x \in \text{bd } T = \text{bd } T' = \text{bd}(\bigcup_{R \in \mathcal{T}'} R)$. A sequence (x_i) exists of points converging to x such that each x_i is in some tile $R^i \in \mathcal{T}'$. For each of these tiles R^i , there is a word $w(i) \in W$ such that $R^i = \Phi_{w(i)}(T_0)$. Observe that

$$\begin{aligned} d_H(R^i, \Phi_{w(i)}(F)) &= d_H(\Phi_{w(i)}(T_0), \Phi_{w(i)}(F)) \\ &= r_{w(i)} d_H(T_0, F) \\ &\leq r_{w(i)} \text{diam } C, \end{aligned}$$

since both F and T_0 are subsets of C . For the sequence of tiles R^i , there are two possibilities:

- (i) There is a subsequence (i_k) such that $\text{diam}(\Phi_{w(i_k)}(T_0)) \xrightarrow{k \rightarrow \infty} 0$.
- (ii) There is a constant $c > 0$ such that $\text{diam}(\Phi_{w(i)}(T_0)) \geq c$ for each $i \in \mathbf{N}$.

Case (i) implies $r_{w(i_k)} \rightarrow 0$, and hence $d(x_{i_k}, F) \rightarrow 0$, so that $x \in F$.

Case (ii) is when $x \in \text{bd } R$ for some $R \in \mathcal{T}'$. To see this, observe that $\text{diam}(\Phi_w(T_0)) \geq c$ for only finitely many words $w \in W$. Hence, at least one of these words occurs infinitely often in the sequence $(w(i))$, i.e., there is a $w \in W$ and a subsequence (i_k) such that $w(i_k) = w$ for all k . But this implies $x_{i_k} \in \Phi_w(T_0) =: R$ for all k , and thus $x \in \overline{R}$, since $x_{i_k} \rightarrow x$. It follows that $x \in \text{bd } R$, since R is open and $x \in \text{bd } T'$. \square

Proposition 3.5.

$$\dim_H \text{bd } T = \max\{\dim_H F, d - 1\}.$$

Proof. For $T_0 = \text{int}(C \setminus \Phi(C))$, observe that $\text{bd } T_0$ is a subset of $\text{bd } C \cup \bigcup_j \text{bd } \Phi_j C$. Since C and $\{\Phi_j C\}_{j=1}^N$ are convex, their boundary has dimension $d - 1$. It follows that $\dim_H \text{bd } T_0 \leq d - 1$ by stability and monotonicity of \dim_H . For the reverse inequality, note that $\text{bd } T_0$ is the boundary of an open set in \mathbf{R}^d . Hence, $\dim_H \text{bd } T_0 = d - 1$, and so $\dim_H \text{bd } R = d - 1$ for each $R \in \mathcal{T}'$. Now the assertion follows from Lemma 3.4 by countable stability of the Hausdorff dimension. \square

4. Compatibility of the ε -parallel sets F_ε and $T_{-\varepsilon}$. In this section, we clarify the relation between the (*outer*) *parallel sets* of F and the (*inner*) *parallel sets* of the associated tiling \mathcal{T} . We characterize the situation in which these parallel sets essentially coincide, for this allows one to use the tiling and the theory of complex dimensions developed in [8] to obtain a tube formula for F .

Definition 4.1. For any nonempty, bounded set $A \subseteq \mathbf{R}^d$, and $\varepsilon \geq 0$, define the (*outer*) ε -*parallel set* (or ε -*neighborhood*) of A by

$$(4.1) \quad A_\varepsilon := \{x \in \overline{A^{\mathbb{G}}} : d(x, A) \leq \varepsilon\}.$$

Similarly, define the (*inner*) ε -*parallel set* (or *inner* ε -*neighborhood*) of A by

$$(4.2) \quad A_{-\varepsilon} := \{x \in \overline{A} : d(x, A^{\mathbb{G}}) \leq \varepsilon\},$$

or equivalently, by $A_{-\varepsilon} = (A^{\mathbb{G}})_\varepsilon$.

Note that we do not include interior points of A into the outer parallel sets, as is often done. For each $\varepsilon \geq 0$, both A_ε and $A_{-\varepsilon}$ are always closed, bounded and nonempty subsets of \mathbf{R}^d . Moreover, $A_0 = A_{-0} = \text{bd } A \subseteq A_\delta$ for any $\delta \in \mathbf{R}$, and $A_{-\varepsilon} = A$ for $\varepsilon \geq \rho$, where $\rho = \rho(A)$ denotes the inradius of A . In particular, if $\text{int } A = \emptyset$, then $A_{-\varepsilon} = A$ for all $\varepsilon \geq 0$.

For an open tiling $\mathcal{A} = \{A^i\}$ (cf. Definition 3.1), denote by $A_{-\varepsilon}$ the inner ε -parallel set of $A := \bigcup_i A^i$.

Lemma 4.2.

$$A_{-\varepsilon} = \overline{\bigcup_{i=1}^\infty A_{-\varepsilon}^i}$$

Proof. Let $x \in A_{-\varepsilon}$. Then $x \in \text{bd } A$ or there is some $l \in \mathbf{N}$ such that $x \in A^l$. In the first case, by Lemma 3.2, $x \in \text{bd } A = \overline{\bigcup_i \text{bd } A^i} \subseteq \overline{\bigcup_i A_{-\varepsilon}^i}$, since $\text{bd } A^i \subset A_{-\varepsilon}^i$. In the latter case $d(x, (A^l)^\complement) = d(x, (\bigcup_i A^i)^\complement) \leq \varepsilon$, and thus $x \in A_{-\varepsilon}^l \subseteq \overline{\bigcup_{i=1}^\infty A_{-\varepsilon}^i}$. Hence, $A_{-\varepsilon} \subseteq \overline{\bigcup_{i=1}^\infty A_{-\varepsilon}^i}$.

For the reverse inclusion, let $x \in \overline{\bigcup_{i=1}^\infty A_{-\varepsilon}^i}$. Then there exists a sequence $y_j \in \bigcup_{i=1}^\infty A_{-\varepsilon}^i$ with $y_j \rightarrow x$ as $j \rightarrow \infty$. For each j , there is an index $i(j) \in \mathbf{N}$ such that $y_j \in A_{-\varepsilon}^{i(j)}$, i.e., $y_j \in \overline{A^{i(j)}}$ and $d(x, (A^{i(j)})^\complement) \leq \varepsilon$. Since $A^{i(j)} \subseteq A$, we infer $y_j \in \overline{A}$ and $d(x, A^\complement) \leq d(x, (A^{i(j)})^\complement) \leq \varepsilon$, i.e., $y_j \in A_{-\varepsilon}$. But this implies $x \in A_{-\varepsilon}$, since $A_{-\varepsilon}$ is closed. \square

Now let $\{\Phi_1, \dots, \Phi_N\}$ be a self-affine system satisfying TSC and NTC, F its attractor and $\mathcal{T} = \{R^i\}_{i \in \mathbf{N}}$ the associated canonical tiling, as introduced in the previous sections. Write $T := \bigcup_i R^i$ for the union of the tiles of \mathcal{T} . For $\varepsilon \geq 0$, the set $T_{-\varepsilon}$ will be regarded as the *inner ε -parallel set* of the tiling.

Proposition 4.3. *Let F be the self-affine set associated to the system $\{\Phi_1, \dots, \Phi_N\}$ satisfying TSC and NTC, and let \mathcal{T} be the associated canonical self-affine tiling of its convex hull C . Then*

- (i) $F \subseteq \text{bd } T$.
- (ii) $F_\varepsilon \cap C \subseteq T_{-\varepsilon}$ for $\varepsilon \geq 0$.
- (iii) $F_\varepsilon \cap C^\complement \subseteq C_\varepsilon$ for $\varepsilon \geq 0$.




FIGURE 2. The exterior ε -neighborhood of the Sierpinski gasket F is the union of the inner ε -neighborhood of the Sierpinski gasket tiling and the exterior ε -neighborhood of $C = [F]$. This union is disjoint except for the boundary of C .

Proof. (i) This is a corollary of Lemma 3.4.

(ii) Fix $\varepsilon \geq 0$. Let $x \in F_\varepsilon \cap C$. Then, since $x \in C = \overline{T}$, either $x \in \text{bd}T$ or $x \in T$. In the former case $x \in T_{-\varepsilon}$ is obvious, since $\text{bd}T \subseteq T_{-\varepsilon}$. In the latter case a point $y \in F$ exists with $d(x, y) \leq \varepsilon$. By (i), y is in $\text{bd}T$ and so $d(x, \text{bd}T) \leq \varepsilon$. Hence $x \in T_{-\varepsilon}$, completing the proof of (ii).

(iii) is an immediate consequence of the inclusion $F \subseteq C$. □

In Theorems 4.4 and 4.9, we characterize the situation in which one has the helpful disjoint decomposition

$$(4.3) \quad F_\varepsilon = T_{-\varepsilon} \cup (C_\varepsilon \setminus C),$$

see Figure 2. Decomposition (4.3) is ensured by (v) and (vi), and the other conditions (i)–(iv) provide easy to check criteria for when this holds. See also Theorem 4.9 for two more equivalent conditions.

Theorem 4.4 (Compatibility theorem). *Let F be the self-affine set associated to the system $\{\Phi_1, \dots, \Phi_N\}$ which satisfies TSC and NTC. Then the following assertions are equivalent:*

- (i) $\text{bd}T = F$.
- (ii) $\text{bd}C \subseteq F$.
- (iii) $\text{bd}(C \setminus \Phi(C)) \subseteq F$.
- (iv) $\text{bd}G_q \subseteq F$ for all $q \in Q$.
- (v) $F_\varepsilon \cap C = T_{-\varepsilon}$ for all $\varepsilon \geq 0$.
- (vi) $F_\varepsilon \cap C^\circ = C_\varepsilon \setminus \text{bd}C$ for all $\varepsilon \geq 0$.

Proof. We show the inclusions (i) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (iv) \Rightarrow (i), then (i) \Rightarrow (v) \Rightarrow (iv) and (ii) \Leftrightarrow (vi).

(i) \Rightarrow (ii). Observe that $\text{bd } C \subseteq \text{bd } T$.

(ii) \Rightarrow (iii). Assume that $\text{bd } C \subseteq F$. Then also $\text{bd } \Phi(C) \subseteq \Phi(\text{bd } C) \subseteq \Phi(F) = F$ and so $\text{bd } (C \setminus \Phi(C)) \subseteq \text{bd } C \cup \text{bd } \Phi(C) \subseteq F$. (Here we used that, for $A, B \subseteq \mathbf{R}^d$, $\text{bd } (A \cup B) \subseteq \text{bd } A \cup \text{bd } B$ and $\text{bd } (A \setminus B) \subseteq \text{bd } A \cup \text{bd } B$.)

(iii) \Rightarrow (iv). Assume that $\text{bd } (C \setminus \Phi(C)) \subseteq F$. The generators G_q (being the connected components of the open set $\text{int } (C \setminus \Phi(C))$) form an open tiling of the set $\text{int } (C \setminus \Phi(C))$. Therefore, by Lemma 3.2, $\text{bd } G_q \subseteq \bigcup_q \text{bd } G_q \subseteq \overline{\bigcup_q \text{bd } G_q} = \text{bd } (\bigcup_q G_q) = \text{bd } (\text{int } (C \setminus \Phi(C))) \subseteq \text{bd } (C \setminus \Phi(C))$. Hence, $\text{bd } G_q \subseteq \text{bd } (C \setminus \Phi(C)) \subseteq F$ for each q , showing (iv).

(iv) \Rightarrow (i). Let $\text{bd } G_q \subseteq F$ for all $q \in \{1, \dots, Q\}$. It suffices to show that this implies $\text{bd } T \subseteq F$, the reversed inclusion always being true, cf. Proposition 4.3 (i). By definition of the tiles, $R^i = \Phi_w G_q$ for some $w \in W$ and some q , and thus we have $\text{bd } R^i = \text{bd } \Phi_w G_q = \Phi_w \text{bd } G_q \subseteq \Phi_w F \subseteq F$ for each $i \in \mathbf{N}$. But this implies $\overline{\bigcup_i \text{bd } R^i} \subseteq F$, since F is closed. Finally, since, by Lemma 3.2, $\text{bd } T = \overline{\bigcup_i \text{bd } R^i}$, assertion (i) follows.

(i) \Rightarrow (v). By Proposition 4.3 (iv), it suffices to show the inclusion $T_{-\varepsilon} \subseteq F_\varepsilon \cap C$ for each $\varepsilon \geq 0$. So fix $\varepsilon \geq 0$, and let $x \in T_{-\varepsilon}$. Then, clearly, $x \in C$. Moreover, either $x \in \text{bd } T$ or $x \in R^i$ for some $i \in \mathbf{N}$ and $d(x, \text{bd } R^i) \leq \varepsilon$. Both cases imply $x \in F_\varepsilon$, the former since, by (i), $\text{bd } T = F \subseteq F_\varepsilon$, and the latter since $\text{bd } R^i \subseteq \bigcup_j \text{bd } R^j \subseteq \text{bd } T = F$, and so $d(x, F) \leq d(x, \text{bd } R^i) \leq \varepsilon$.

(v) \Rightarrow (iv) (by contraposition). Assume that (iv) is false, i.e., assume there exists some index q and some $x \in \text{bd } G_q$ such that $x \notin F$. Then, since F is closed, there is some number $\delta > 0$ such that $d(x, F) > \delta$ and so $x \notin F_\varepsilon$ for $\varepsilon \leq \delta$. On the other hand, $x \in \text{bd } G_q$ clearly implies $x \in T_{-\varepsilon}$. Hence, the equality in (v) does not hold.

(ii) \Rightarrow (vi). For $\varepsilon = 0$, there is nothing to prove. So let $\varepsilon > 0$ and $x \in C_\varepsilon \setminus \text{bd } C$. Then there exists a point $y \in \text{bd } C$ such that $d(x, y) \leq \varepsilon$. By (ii), $y \in F$, and thus $d(x, F) \leq \varepsilon$, i.e., $x \in F_\varepsilon$. Hence, $C_\varepsilon \subseteq F_\varepsilon \cap C^{\text{cl}}$. The reversed inclusion is always true, cf. Proposition 4.3 (iii), and so assertion (vi) follows.

(vi) \Rightarrow (ii) (by contraposition). Assume (ii) is false, i.e., a point $x \in \text{bd } C$ exists such that $x \notin F$. Let $\delta := d(x, F)$ and fix some $\varepsilon < \delta/2$. Since $x \in \text{bd } C$, there are points in $C^{\mathbb{G}}$ arbitrarily close to x . Choose $y \in C_\varepsilon \cap C^{\mathbb{G}}$. Then $d(y, F) \geq d(x, F) - d(x, y) > \varepsilon$, implying $y \notin F_\varepsilon$. Hence, equality $F_\varepsilon \cap C^{\mathbb{G}} = C_\varepsilon \setminus \text{bd } C$ cannot be true for this ε , i.e., (vi) does not hold. \square

Note that assertions (ii), (iii) and (iv) are very simple and easy to check. So, in particular, Theorem 4.4 states that if one of the assertions (ii), (iii) or (iv) is true for a given self-affine set F , then each of its parallel sets F_ε is the disjoint union of the two sets $T_{-\varepsilon}$ and $C_\varepsilon \setminus C$, cf. (4.3). Moreover, if for some F , it can be shown that one of assertions (ii), (iii) or (iv) is false, then the inner parallel set $T_{-\varepsilon}$ of the tiling does not describe set $F_\varepsilon \cap C$, and also the sets $F_\varepsilon \cap C^{\mathbb{G}}$ and $C_\varepsilon \setminus C$ are different. Thus, for any set F not satisfying the assertions of Theorem 4.4, F_ε does not coincide with $T_{-\varepsilon} \cup C_\varepsilon$, and one cannot use the tiling directly to study the parallel sets F_ε .

The envelope. We consider another hull operation, the *envelope* and show that the conditions of the compatibility theorem are met for a self-affine set F precisely when its envelope coincides with its convex hull. At the end of Section 6, we examine the feasibility of the envelope as a replacement for the convex hull in the tiling construction, cf. Proposition 6.3 and the ensuing discussion. There are many cases where Theorem 4.4 does not apply for the tiling as constructed using the convex hull, but the analogous result (Theorem 6.2) *does* apply when the convex hull is replaced by the envelope.

Definition 4.5. Let $K \subset \mathbf{R}^d$ be a compact set. $K^{\mathbb{G}}$ has a unique unbounded component, which we call U . (For $d = 1$, there are actually two unbounded components in $K^{\mathbb{G}}$, if $+\infty$ and $-\infty$ are not identified. In this case let U be their union.) Then $\text{bd } U$ is the *exterior boundary* of K ; it consists of that portion of (the boundary of) K which is accessible when approaching K from infinity. The *envelope* $E = E(K)$ of K is the complement of U , $E := U^{\mathbb{G}}$.

Example 4.6. The envelope of the Sierpinski gasket is its convex hull, as is the envelope of the Sierpinski carpet. The envelope of the Koch curve is the Koch curve itself, as is the envelope of the attractor depicted in Figure 1. Some more interesting (and nonconvex) envelopes are shown in Figure 3; for a description of these sets, cf. [12].




FIGURE 3. Three self-similar sets and their envelopes (the union of the shaded region and the attractor). F_3 is the attractor of a system $\Phi^{(1)}$ of seven mappings, each with scaling ratio $1/3$ and no rotation. To construct F_2 , we have given two of the mappings a rotation of π (top left and top right). To make F_1 , we have additionally given one of the mappings a rotation of $\pi/2$ (bottom center).

Lemma 4.7. *Let $K \subset \mathbf{R}^d$ be a compact set. The envelope E of K is compact and satisfies $\text{bd } E \subseteq K \subseteq E$. Moreover, $E \subseteq [K]$, where $[K]$ is the convex hull of K .*

The following results indicate that the conditions of Theorem 4.4 are satisfied precisely when a self-affine set F appears convex when seen “from outside.”

Proposition 4.8. *Let K be a compact set in \mathbf{R}^d with envelope E and convex hull $[K]$. Then E is convex if and only if $E = [K]$.*

Proof. If $E = [K]$, then E is obviously convex. For the other implication, assume E is convex. By Lemma 4.7, we have $E \subseteq [K]$ and, moreover, $K \subseteq E$. The latter implies $[K] \subset [E]$ and, since $E = [E]$, the reversed inclusion $[K] \subseteq E$ is also proved. □

Now we have two more “compatibility conditions” to accompany those already established in Theorem 4.4. Let E denote the envelope of F , and let C be its convex hull, as before.

Theorem 4.9. *Each of the following two conditions is equivalent to any of conditions (i)–(vi) of Theorem 4.4:*

- (a) $E = C$.
- (b) E is convex.

Proof. We prove (ii) \Rightarrow (b) and (a) \Rightarrow (ii), where (iv) and (ii) are the conditions from Theorem 4.4. Note that (a) is equivalent to (b) by Proposition 4.8.

(ii) \Rightarrow (b) (by contraposition). If E is not convex, then $\text{int } C \setminus E$ is nonempty and so a point $x \in \text{int } C \cap U$ must exist, i.e., x is in the unbounded connected component U of F^c . Hence, there must be a path in U connecting x to infinity. Since $x \in \text{int } C$, this path crosses $\text{bd } C$, implying the existence of a point $y \in \text{bd } C$ which is not in F . Hence, condition (ii) of Theorem 4.4 does not hold.

(a) \Rightarrow (ii). Note that $E = C$ is true if and only if $\text{bd } U = \text{bd } C$. Since $\text{bd } U \subseteq F$, we have $\text{bd } C \subseteq F$, which is condition (ii). \square

5. Generalization of the tiling construction. While the non-triviality condition is not very restrictive, the tileset condition puts a serious constraint on the class of sets for which the canonical tiling exists. For the purpose of obtaining tube formulas for F , the compatibility conditions need to be satisfied; this limits the applicability of the tiling construction even further. It is natural to ask whether the tiling construction can be modified to work for more general sets. The NTC and TSC are both necessary restrictions, and each is given in terms of the convex hull C of F . While neither condition can be omitted, they *can* be applied to a different initial set for the tiling construction, in place of C . Provided F satisfies OSC, it turns out that any feasible open set O of F can be used as the initial set. In this section, we show that the tiling construction can still be carried out in this generalized setting. In the next section, we examine the analogue of the compatibility theorem for this generalization.

The main result of this section is Theorem 5.7, which can be paraphrased as follows: if F is a self-affine set with empty interior and which satisfies the OSC with feasible set O , then there exists a self-affine tiling of \overline{O} . In other words, we generalize the tiling construction by replacing the convex hull with the set $K = \overline{O}$, where O is an arbitrary feasible open set for F . The open set condition takes the role of the tileset condition, and we obtain an open tiling of K . The canonical

self-affine tiling of the convex hull of F appears as the special case of this construction in case $\text{int } C$ is a feasible open set. We will need the following well-known fact.

Proposition 5.1 (cf. [6, 5.1 (3) (ii)]). *If F satisfies OSC with feasible open set O , then $F \subseteq \overline{O}$.*

Let $\{\Phi_1, \dots, \Phi_N\}$ be a self-affine system satisfying OSC, and let F be its attractor. Let O be any feasible open set for F , i.e., O satisfies (2.1) and (2.2). Set $K := \overline{O}$ from now on. Since $O \subseteq \text{int } (K)$, it is clear that K is the closure of its interior, $K = \overline{\text{int } (K)}$, and that $F \subseteq K$, by Proposition 5.1. It is easily seen that (2.2) implies

$$(5.1) \quad \Phi_w(O) \cap \Phi_v(O) = \emptyset \quad \text{for all } w, v \in W^j, v \neq w, j \in \mathbf{N}.$$

Write $O^k := \Phi^k(O)$ and $K^k := \Phi^k(K)$ for $k = 0, 1, 2, \dots$

Proposition 5.2 (Nestedness). For $k = 0, 1, 2, \dots$, one has $K^{k+1} \subseteq K^k \subseteq K$.

Proof. Note that (2.1) implies $O^{k+1} \subseteq O^k$. □

Proposition 5.2 extends [15, Theorem 5.1] and shows that $K \supseteq K^1 \supseteq K^2 \supseteq \dots$ is a decreasing sequence of sets which converges to F ; note that $F = \bigcap_{k=0}^\infty K^k$, by the contraction principle. In analogy with the tiling construction for the convex hull, the following nontriviality condition is required for a tiling of O to exist:

Definition 5.3. A self-affine set F satisfying OSC is said to be *nontrivial* if there exists a feasible open set O for F such that

$$(5.2) \quad O \not\subseteq \Phi(\overline{O});$$

F is called *trivial* otherwise.

In fact, nontriviality implies that (5.2) holds for all feasible sets O of F . This is a consequence of the following proposition which

characterizes the trivial case: a set F is trivial if and only if it has interior points. Hence, triviality and non triviality are independent of the particular choice of set O .

Proposition 5.4 (Characterization of triviality). *Let F be a self-affine set satisfying OSC. Then the following assertions are equivalent:*

- (i) F is trivial.
- (ii) $\text{int } F \neq \emptyset$.
- (iii) $\overline{O} = F$ for some feasible open set O of F .
- (iv) $\overline{O} = F$ for each feasible open set O of F .

Proof. (i) \Rightarrow (iv). Let O be an arbitrary feasible set for F ; we already have one containment from Proposition 5.1. Assume F is trivial, which means $O \subseteq \Phi(\overline{O})$. Taking the closure, we get $K \subseteq \Phi(K)$. The OSC implies $\Phi(O) \subseteq O$ which, by taking closures again, implies $\Phi(K) \subseteq K$. Hence, $K = \Phi(K)$. By the uniqueness of the invariant set, we infer $F = K$.

(iv) \Rightarrow (iii) and (iii) \Rightarrow (ii) are trivial. For the latter, note that a feasible set O is nonempty.

(ii) \Rightarrow (i) (by contraposition). If F is nontrivial, then there is a feasible set O such that $O \not\subseteq \Phi(K)$. Hence, the set $T_0 := O \setminus \Phi(K)$ is nonempty, but $T_0 \cap F = \emptyset$, since $F \subseteq \Phi(K)$. Observe that OSC implies $\Phi_i(O) \cap \Phi_j(F) = \emptyset$ for $i \neq j$. Therefore, $\Phi_j(T_0) \cap F \subseteq \Phi_j(T_0) \cap \Phi_j(F) = \Phi_j(T_0 \cap F) = \emptyset$, and so $\Phi(T_0) \cap F = \emptyset$. By induction, we get $\Phi^k(T_0) \cap F = \emptyset$ for $k = 0, 1, 2, \dots$. Now let $x \in F$. Since, by the contraction principle, $d_H(F, \Phi^k(\overline{T_0})) = d_H(\Phi^k(F), \Phi^k(\overline{T_0})) \rightarrow 0$ as $k \rightarrow \infty$, a sequence $x_k \rightarrow x$ exists with $x_k \in \Phi^k(\overline{T_0}) = \overline{\Phi^k(T_0)}$. For each x_k there are points in $\Phi^k(T_0)$ arbitrarily close to x_k . Hence, x is not an interior point of F . \square

Remark 5.5. Note that Proposition 5.4 provides an easy criterion to decide whether a self-affine set has interior points. Take an arbitrary feasible open set O of F and check whether O contains a point with positive distance to F . If not, then F has interior points, otherwise $\text{int } F$ is empty. Conversely, if it is known for some F that it has

a nonempty interior, then the search for a feasible open set can be restricted to subsets of F .

For completeness, we note that Corollary 2.13 also generalizes to this more general notion of nontriviality used here. The argument in the proof carries over, when taking Proposition 5.4 into account.

Corollary 5.6. *Let $F \subseteq \mathbf{R}^d$ be a self-affine set satisfying OSC. If F has Hausdorff dimension strictly less than d , then F is nontrivial. Moreover, if F is self-similar, then also the converse holds.*

Now we can state the main result of this section.

Theorem 5.7 (Generalized tiling). *Let F be a self-affine set satisfying $\text{int } F = \emptyset$ and OSC. Let O be an arbitrary feasible open set for F and $K = \overline{O}$. Let G_1, G_2, \dots denote the connected components of the open set $O \setminus \Phi(K)$. Then*

$$\mathcal{T}(O) := \{\Phi_w(G_q) : w \in W, q \in Q\}$$

is an open tiling of K , i.e., the tiles $\Phi_w(G_q)$ are pairwise disjoint and

$$K = \overline{\bigcup_{R \in \mathcal{T}(O)} R}.$$

To prepare the proof, we note the following facts.

Lemma 5.8. *Let $A = A^0 \supseteq A^1 \supseteq A^2 \supseteq \dots$ be a decreasing sequence of sets, and define $B := \bigcap_{k=0}^\infty A^k$. Then we can decompose A as the disjoint union*

$$A = B \cup \bigcup_{k=0}^\infty (A^k \setminus A^{k+1}).$$

Denote $T_0 := \bigcup_{q \in Q} G_q = \text{int}(O \setminus \Phi(O))$, and more generally, set $T_k := O^k \setminus K^{k+1} = \text{int}(K^k \setminus K^{k+1})$ for $k = 1, 2, \dots$. We now adapt [15, Theorem 5.14, page 3165] to the present more general setting.

Lemma 5.9 (Propagation of tilesets). $\Phi(T_k) = T_{k+1}$, for each $k = 0, 1, 2, \dots$.

Proof (\subseteq). Let $x \in \Phi(T_k)$. Choose j so that $x \in \Phi_j(T_k) = \Phi_j(O^k) \setminus \Phi_j(K^{k+1})$. Then $x \in \Phi_j(O^k) \subseteq O^{k+1}$. To see $x \notin K^{k+2}$, suppose it is. Then $x \in \Phi_\ell(K^{k+1})$ for some ℓ . Note that $\ell \neq j$, since, by the choice of j , $x \notin \Phi_j(K^{k+1})$. Now $K^{k+1} \subseteq K^k$ by Proposition 5.1, which implies $x \in \Phi_\ell(K^{k+1}) \subseteq \Phi_\ell(K^k)$, and hence $x \in \Phi_\ell(K^k) \cap \Phi_j(O^k)$. Since $\Phi_j(O^k)$ is open and $\Phi_\ell(K^k)$ the closure of its interior, there must be points of $\Phi_j(O^k)$ in the interior of $\Phi_\ell(K^k)$, i.e., in $\Phi_\ell(O^k)$, contradicting OSC.

(\supseteq). Pick $x \in T_{k+1} = O^{k+1} \setminus K^{k+2}$. Then, $x \in O^{k+1} = \Phi(O^k)$, and so $x \in \Phi_j(O^k)$ for some j . Hence, $x = \Phi_j(y)$ for $y \in O^k$. If $y \in K^{k+1}$, then $x = \Phi_j(y) \in K^{k+2}$, a contradiction to $x \in T_{k+1}$. So $y \notin K^{k+1}$, and hence $y \in O^k \setminus K^{k+1}$. We conclude $x = \Phi_j(y) \in \Phi_j(O^k \setminus K^{k+1}) \subseteq \Phi(O^k \setminus K^{k+1})$, which completes the proof. \square

Proof of Theorem 5.7. Since F has no interior points, by Proposition 5.4, F is nontrivial, i.e., the set T_0 is nonempty. Since $T_0 = \bigcup_{q \in Q} G_q$, Lemma 5.9 immediately implies

$$(5.3) \quad T_k = \bigcup_{\substack{w \in W^k \\ q \in Q}} \Phi_w(G_q).$$

In the following, $T := \bigcup_{R \in \mathcal{T}(O)} R$ denotes the union of all the tiles in $\mathcal{T}(O)$. Since $\Phi_w(G_q) \subseteq O \subset K$, we have $T \subseteq K$, and since K is closed the inclusion $\overline{T} \subseteq K$ is obvious. It remains to show the reversed inclusion. Let $x \in K$. By Lemma 5.8, either $x \in \bigcap_k K^k = F$ or there is some $k \in \mathbf{N}_0$ such that $x \in K^k \setminus K^{k+1} \subseteq \overline{T}_k$. If $x \in \overline{T}_k$, then equation (5.3) implies $x \in \overline{\bigcup_w \Phi_w(G_q)}$, where the union is taken over all $w \in W^k$ and $q \in Q$, and therefore $x \in \overline{T}$. If $x \in F$, then there exists a sequence $(x_i) \in K \setminus F$ converging to x (since F has no interior points). The previous argument shows $x_i \in \overline{T}$, and hence the same holds for $x = \lim x_i$. This shows $K \subseteq \overline{T}$, and hence $K = \overline{T}$.

By Lemma 5.8, the sets $K^k \setminus K^{k+1}$ are pairwise disjoint, and hence so are the sets T_k . Moreover, the union in (5.3) is disjoint, which follows immediately from (5.1) and the fact that the sets G_q are pairwise

disjoint and subsets of O . Hence, the sets $\Phi_w(G_q) \in \mathcal{T}(O)$ are pairwise disjoint. \square

As a corollary to the proof we note the following for later use.

Corollary 5.10. $R \cap F = \emptyset$ for each $R \in \mathcal{T}(O)$.

Proof. Recall that $F \subseteq K^k$ for each k . Hence, $T_k = O^k \setminus K^{k+1}$ has empty intersection with F . By (5.3), each tile $R \in \mathcal{T}(O)$ is contained in one of the sets T_k . \square

Remark 5.11 (Different open sets may yield the same tiling). Note that two feasible open sets O and O' do not necessarily produce different tilings. If $\overline{O} = \overline{O'}$, then the tilings $\mathcal{T}(O)$ and $\mathcal{T}(O')$ coincide. In both cases one obtains an open tiling of the set $K = \overline{O} = \overline{O'}$. Therefore, for most questions it suffices to restrict considerations to feasible sets O satisfying $\text{int } \overline{O} = O$.

Remark 5.12 (The dimension of the boundary of the tiling in the general case). Proposition 3.5 states that $\dim_H \text{bd } T = \max\{\dim_H F, d - 1\}$ for the $\mathcal{T}(C)$ of the convex hull, but this does not extend to the generalized tilings $\mathcal{T}(O)$. In general, $\text{bd } O$ will not be of dimension $d - 1$ and the tiles of $\mathcal{T}(O)$ may still have a fractal boundary. So the statement is slightly different. For the boundary $\text{bd } T$ of a tiling $\mathcal{T}(O)$, one has $\dim_H \text{bd } T = \max\{\dim_H F, \dim_H \text{bd } T_0\}$, where $\dim_H \text{bd } T_0 \geq d - 1$.

6. Generalizing the compatibility theorem. In the previous section we constructed a tiling for each feasible open set of a self-affine set F , provided F is nontrivial in the sense of Definition 5.3. The motivation was to find a tiling which can be used to decompose the parallel sets of F . The theme of this section is the search for feasible open sets that are suitable for this purpose. We revisit the Compatibility Theorem of Section 4 and find conditions on a feasible open set that allow for an analogue of Theorem 4.4.

We start by discussing an appropriate generalization of Proposition 4.3. Throughout, we use the notation of the previous section.

In particular, for a self-affine set F and a feasible open set O , $\mathcal{T}(O)$ is the associated self-affine tiling, G_q are the generators, $T = \bigcup_{R \in \mathcal{T}(O)} R$ is the union of the tiles and $T_{-\varepsilon}$ the inner parallel set of T .

Proposition 6.1. *Let F be the self-affine set associated to the system $\{\Phi_1, \dots, \Phi_N\}$. Assume that F has empty interior and satisfies the OSC with a feasible set O . Let $\mathcal{T}(O)$ be the associated tiling and $K = \overline{O}$. Then:*

- (i) $F \subseteq \text{bd } T$.
- (ii) $F_\varepsilon \cap K \subseteq T_{-\varepsilon}$ for $\varepsilon \geq 0$.
- (iii) $F_\varepsilon \cap K^{\mathbb{C}} \subseteq K_\varepsilon$ for $\varepsilon \geq 0$.

Proof. (i) On the one hand, $\mathcal{T}(O)$ is an open tiling of K , and thus $F \subseteq K = \overline{T}$. On the other hand, by Corollary 5.10, $R \cap F = \emptyset$ for all $R \in \mathcal{T}(O)$, i.e., $F \cap T = \emptyset$. Thus, $F \subseteq \overline{T} \setminus T = \text{bd } T$.

(ii) Fix $\varepsilon \geq 0$. Let $x \in F_\varepsilon \cap K$. Then, since $x \in K = \overline{T}$, either $x \in \text{bd } T$ or $x \in T$.

In the former case $x \in T_{-\varepsilon}$ is obvious, since $\text{bd } T \subseteq T_{-\varepsilon}$. In the latter case there exists a point $y \in F$ with $d(x, y) \leq \varepsilon$. By (i), y is in $\text{bd } T$ and so $d(x, \text{bd } T) \leq \varepsilon$, whence $x \in T_{-\varepsilon}$.

(iii) is an immediate consequence of the inclusion $F \subseteq K$. □

Theorem 6.2 (Generalized Compatibility theorem). *Let F be the self-affine set associated to the system $\{\Phi_1, \dots, \Phi_N\}$. Assume that F has empty interior and satisfies the OSC with a feasible set O . Let $\mathcal{T}(O)$ be the associated tiling of O . Then the following assertions are equivalent:*

- (i) $\text{bd } T = F$.
- (ii) $\text{bd } K \subseteq F$.
- (iii) $\text{bd}(K \setminus \Phi(K)) \subseteq F$.
- (iv) $\text{bd } G_q \subseteq F$ for all $q \in Q$.
- (v) $F_\varepsilon \cap K = T_{-\varepsilon}$ for all $\varepsilon \geq 0$.
- (vi) $F_\varepsilon \cap K^{\mathbb{C}} = K_\varepsilon \cap K^{\mathbb{C}}$ for all $\varepsilon \geq 0$.

Proof. Observe that in the proof of Theorem 4.4 the convexity of set C is not used (just the inclusion $F \subseteq C$, which is also satisfied here by

Proposition 5.1: $F \subseteq \overline{O} = K$). Thus, the proof can be carried over to the new situation by replacing C with K and applying Proposition 6.1 instead of Proposition 4.3 where necessary. \square

Theorem 6.2 does not indicate whether or not one can always find a set O such that the associated tiling can be used to decompose F_ε . That is, one might ask if for any self-affine set F (satisfying OSC and $\text{int } F = \emptyset$) there is always a feasible set O such that the equivalent conditions (i)–(vi) are satisfied. Unfortunately, this is not the case in general. There are sets for which no such O exists, for instance the *Koch curve*. In fact, for fractals with connected complement, Proposition 6.3 shows that the Compatibility Theorem is never satisfied. The class of sets characterized by the connectedness of $F^{\mathbb{C}}$ includes all simple fractal curves (curves with no self-intersection) like the *Koch curve*, all tree-like sets (dendrites) and all totally disconnected sets in \mathbf{R}^d with $d \geq 2$. For $d \geq 3$, it even includes topologically nontrivial sets like the Menger sponge.

Proposition 6.3. *Let F be a self-affine set satisfying OSC and $\text{int } F = \emptyset$. If the complement of F is connected, then there is no feasible open set O such that $\text{bd } K \subseteq F$.*

Proof. It suffices to consider feasible open sets O satisfying $\text{int } \overline{O} = O$ (cf. Remark 5.11), which implies $\text{bd } K = \text{bd } O$. Since F has no interior points, we have $O \cap F^{\mathbb{C}} \neq \emptyset$. Let $x \in O \cap F^{\mathbb{C}}$. Since O is not the whole set $F^{\mathbb{C}}$, there is also a point $y \in F^{\mathbb{C}} \setminus O$. Since $F^{\mathbb{C}}$ is connected by assumption, it is also path connected. Hence, there is a path from x to y in $F^{\mathbb{C}}$, and it must cross the boundary $\text{bd } O = \text{bd } K$ somewhere. Hence, $\text{bd } K$ is not completely contained in F . \square

From the proof of Proposition 6.3, it is clear that any feasible set O satisfying compatibility must be a subset of envelope E of F , the complement of the unbounded component of $F^{\mathbb{C}}$ (cf. Definition 4.5). For many self-affine sets F the (interior of the) envelope itself is compatible; note that envelope E always satisfies the compatibility condition $\text{bd } E \subset F$, by definition. If E is feasible, then a tiling exists that can be used to describe the parallel sets of F .

Corollary 6.4. *Let F be a self-affine set with $\text{int } F = \emptyset$ and satisfying OSC. If $\text{int } E$ is a feasible open set for F , then the self-affine*




FIGURE 4. A self-similar set which satisfies OSC but for which the envelope is not feasible, see Example 6.5.

tiling $\mathcal{T} = \mathcal{T}(\text{int } E)$ allows a decomposition of F_ε ,

$$(6.1) \quad F_\varepsilon = T_{-\varepsilon} \cup E_\varepsilon,$$

which is disjoint but for the null set $\text{bd } E$.

However, one should not be overoptimistic; the set $F \subset \mathbf{R}$ in Example 2.6 shows that $\text{int } E$ is not always a feasible open set (in this case, envelope E coincides with the convex hull). We also provide the following example.

Example 6.5. Let F be the attractor of system $\{\Phi_1, \dots, \Phi_4\}$ of four similarities, where Φ_1, Φ_2, Φ_3 are the usual mappings used for the Sierpinski gasket and Φ_4 scales the initial triangle by a factor $1/4$, rotates it by π and translates it by $[(1/4, \sqrt{3}/8)]$ such that it fits in the largest hole in $\Phi_1 F$ (cf. Figure 4). This set satisfies OSC, for instance, the set $O := \bigcup_{j=1}^4 \Phi_j C$ is feasible. The envelope of F coincides with the convex hull of F , $E = C$, but E is not feasible since $\Phi_1 E \cap \Phi_4 E$ is not empty.

We believe that, if F° is not connected, i.e., if envelope E of F has nonempty interior, then there always exists a subset O of E which is both compatible and feasible. So far we have not been able to prove this.

Note added in proof. In a recent joint work by the second author with Dušan Pokorný, cf. [16], the following statement was obtained as a side result: Let $F \subset \mathbf{R}^d$ be a self-similar set satisfying OSC and $\text{int } F = \emptyset$. If the complement of F has a bounded connected component, then a

feasible open set exists such that $\text{bd } K \subseteq F$. For self-similar sets, this allows one to strengthen Proposition 6.3 to an if-and-only-if statement. The proof requires the existence of a feasible open set O which satisfies the additional requirement $O \cap F \neq \emptyset$, known as the strong open set condition (SOSC).¹ In conclusion, we have the following result:

Theorem 6.3. *Let $F \subset \mathbf{R}^d$ be a self-affine set satisfying SOSC and $\text{int } F = \emptyset$. Then a feasible open set O exists such that $\text{bd } K \subseteq F$ if and only if the complement of F is not connected.*

7. Concluding comments and remarks.

Remark 7.1. Corollary 5.6 says that the trivial self-similar sets in \mathbf{R}^d are precisely those which have full Hausdorff dimension. Hence, all self-similar sets for which self-similar tilings can be constructed have Hausdorff dimension (and thus Minkowski dimension) strictly less than d . In [10], tube formulas are obtained for a class of fractal sprays in \mathbf{R}^d , provided these sprays satisfy the same condition on the Minkowski dimension of their boundary. So Corollary 5.6 ensures that the latter condition does not impose any restrictions on the applicability of the tube formula results to the self-similar case.

In the self-affine case, however, it remains open whether there exists a nontrivial $F \subset \mathbf{R}^d$ (i.e., one with empty interior) satisfying OSC which has full Hausdorff dimension. On the other hand, tube formulas are not available yet in this more general setting. The results obtained for fractal sprays do not apply in this case.

Remark 7.2 (Relation to tilings of \mathbf{R}^d). There is another notion of self-similar (self-affine) tilings which has been studied at length, namely tilings of the plane or, more generally, of \mathbf{R}^d . In this approach copies of self-similar (or self-affine) sets $F \subseteq \mathbf{R}^d$ are used as tiles to tile the whole of \mathbf{R}^d , which is very different from our approach, where a feasible set of F is tiled and where the tiles are not copies of F but subsets of F^c . See [1, 7, 13, 19], for example.

However, there are interesting relations between both concepts. Firstly, the open set condition is a natural requirement in both approaches. Secondly, the concepts are in a way complementary to each

other. Tilings of \mathbf{R}^d require self-similar sets F to have full dimension, while the tilings of feasible sets require F to have dimension strictly less than d .

Proposition 7.3. *Let F be a self-similar set satisfying OSC. Then there is a dichotomy:*

- (i) $\text{int } F = \emptyset$, in which case there is a self-similar tiling of any feasible open set O of F , or
- (ii) $\text{int } F \neq \emptyset$, in which case F gives a self-similar tiling of \mathbf{R}^d .

Proof. (i) is a corollary of Theorem 5.7; (ii) is [1, Theorem 9.1] with unit tile F . \square

The dichotomy of Proposition 7.3 extends to the self-affine case, see [7, Theorem 1.2] and [13, Lemma 2.3] for case (ii). The latter result is formulated for subsets of \mathbf{R}^2 (and more general contractions) but the same argument works in \mathbf{R}^d .

Acknowledgments. The authors are grateful for helpful discussions, friendly advice, and useful references from Christoph Bandt, Kenneth Falconer, John Hutchinson, Michel Lapidus, Mathias Mesing, Sze-Man Ngai, Bob Strichartz, Luke Rogers, Huo-Jun Ruan, Sasha Teplyaev, Yang Wang, and Martina Zähle.

ENDNOTES

1. While, in the self-similar case, SOSC is known to be equivalent to OSC, cf. [17], this condition has to be imposed in the self-affine setting in order to carry over the proof.

REFERENCES

1. C. Bandt, *Self-similar tilings and patterns described by mappings*, in *The mathematics of long-range aperiodic order*, NATO Adv. Sci. Inst. Math. Phys. Sci. **489** (1997), 45–83.
2. C. Bandt, N. Hung and H. Rao, *On the open set condition for self-similar fractals*, Proc. Amer. Math. Soc. **134** (2005), 1369–1374.
3. K.J. Falconer, *Fractal geometry—Mathematical foundations and applications*, John Wiley, Chichester, 1990.
4. H. Federer, *Curvature measures*, Trans. Amer. Math. Soc. **93** (1959), 418–491.

5. A. Gray, *Tubes* (2nd edition), Progress Math. **221**, Birkhauser, Boston, 2004.
6. J.E. Hutchinson, *Fractals and self-similarity*, Indiana Univ. Math. J. **30** (1981), 713–747.
7. J.C. Lagarias and Y. Wang, *Self-affine tiles in \mathbf{R}^n* , Adv. Math. **121** (1996), 21–49.
8. Michel L. Lapidus and Erin P.J. Pearse, *Tube formulas and complex dimensions of self-similar tilings*, Acta Appl. Math. **112** (2010), 91–137.
9. ———, *Tube formulas of self-similar fractals*, in *Analysis on graphs and its applications*, Proc. Symp. Pure Math. **77**, P. Exner, et al., eds., American Mathematical Society, Providence, RI, 2008.
10. Michel L. Lapidus, Erin P.J. Pearse and Steffen Winter, *Pointwise tube formulas for fractal sprays and self-similar tilings with arbitrary generators*, Adv. Math. **227** (2011), 1349–1398.
11. Michel L. Lapidus and Machiel van Frankenhuysen, *Fractal geometry, complex dimensions and zeta functions: Geometry and spectra of fractal strings*, Springer Mono. Math., Springer-Verlag, New York, 2006.
12. M. Llorente and S. Winter, *A notion of Euler characteristic for fractals*, Math. Nachr. **280** (2007), 152–170.
13. S.-M. Ngai and T.-M. Tang, *Topology of connected self-similar tiles in the plane with disconnected interiors*, Topol. Appl. **150** (2005), 139–155.
14. E.P.J. Pearse, *Complex dimensions of self-similar systems*, Ph.D. dissertation, University of California, Riverside, June 2006.
15. ———, *Canonical self-affine tilings by iterated function systems*, Indiana Univ. Math. J. **56** (2007), 3151–3169.
16. D. Pokorný and S. Winter, *Scaling exponents of curvature measures*, in preparation.
17. A. Schief, *Separation properties for self-similar sets*, Proc. Amer. Math. Soc. **122** (1994), 111–115.
18. R. Schneider, *Convex bodies: The Brunn-Minkowski theory*, Cambridge University Press, Cambridge, 1993.
19. R.S. Strichartz and Y. Wang, *Geometry of self-affine tiles I*, Indiana Univ. Math. J. **48** (1999), 1–23.
20. H. Weyl, *On the volume of tubes*, Amer. J. Math. **61** (1939), 461–472.
21. S. Winter, *Curvature measures and fractals*, Dissert. Math. **453** (2008).
22. M. Zähle, *Curvatures and currents for unions of sets with positive reach*, Geom. Dedi. **23** (1987), 155–171.

DEPARTMENT OF MATHEMATICS, CALIFORNIA POLYTECHNIC STATE UNIVERSITY,
 SAN LUIS OBISPO, CA 93407
Email address: epearse@calpoly.edu

INSTITUTE OF STOCHASTICS, DEPARTMENT OF MATHEMATICS, KARLSRUHE INSTITUTE OF TECHNOLOGY, 76133 KARLSRUHE, GERMANY
Email address: steffen.winter@kit.edu