

EIGENVALUES OF THE LAPLACIAN OF RIEMANNIAN MANIFOLDS

SHÛKICHI TANNO

(Received December 19, 1972)

1. Introduction. Let (M, g) be a compact orientable Riemannian manifold (connected and C^∞ , $\dim M = m$) with metric tensor g . By ∇ we denote the Riemannian connection and by Δ we denote the Laplacian acting on p -forms, $0 \leq p \leq m$. Let $\lambda_{\alpha,p}$ be eigenvalues of Δ and put

$$(1.1) \quad \text{Spec}(M, g; p) = \{0 \leq \lambda_{0,p} \leq \lambda_{1,p} \leq \lambda_{2,p} \leq \cdots \downarrow -\infty\}.$$

For $p = 0$, we denote $\text{Spec}(M, g; 0) = \text{Spec}(M, g)$ and $\lambda_{\alpha,0} = \lambda_\alpha$. Relations between $\text{Spec}(M, g)$ and Riemannian structures have been studied by Berger [1], McKean-Singer [6], Patodi [10], Sakai [11], etc. Some results are listed in the first part of §3. A useful tool is a formula of Minakshisundaram:

$$(1.2) \quad \sum_{\alpha=0}^{\infty} e^{\lambda_\alpha t} \underset{t \rightarrow 0}{\sim} \left(\frac{1}{4\pi t}\right)^{m/2} \sum_{\beta=0}^{\infty} a_\beta t^\beta.$$

a_0, a_1, a_2 were calculated by Berger [1] and McKean-Singer [6]; and a_3 was calculated by Sakai [11]. In the following Theorems A, B, and D, the assumption on $\text{Spec}(\cdot)$ is also replaced by $\sum e^{\lambda_\alpha t}$, and more precisely in terms of a_β ($\beta = 0, 1, 2, 3$).

THEOREM A. *Let (M, g) and (M', g') be compact orientable Riemannian manifolds. Assume that $\text{Spec}(M, g) = \text{Spec}(M', g')$. Then $m = m'$ and*

(1) *for $2 \leq m \leq 5$, (M, g) is of constant curvature K , if and only if (M', g') is of constant curvature $K' = K$,*

(2) *for $m = 6$, (2-1) (M, g) is conformally flat and the scalar curvature S is constant, if and only if (M', g') is conformally flat and the scalar curvature S' is constant, $S' = S$,*

(2-2) *(M, g) is of constant curvature $K > 0$, if and only if (M', g') is of constant curvature $K' = K > 0$.*

Theorem A for $m = 2, 3$ was proved by Berger [1]. For $m = 4$, Berger's Theorem 8.1 in [1] requires an additional condition $\chi(M) = \chi(M')$, where $\chi(M)$ denotes the Euler-Poincaré characteristic of M . Our result

generalizes this and furthermore it is valid for $m = 5$.

By $S^m(c)$ or $(S^m(c), g_0)$ we denote a Euclidean m -sphere with constant curvature $c > 0$, and by $H^m(-c)$ we denote a hyperbolic m -space with constant curvature $-c < 0$.

THEOREM B. *Let (M, g) be a compact orientable Riemannian manifold, $2 \leq m \leq 6$. If $\text{Spec}(M, g) = \text{Spec}(S^m(c), g_0)$, then (M, g) is isometric to $(S^m(c), g_0)$.*

This follows from Theorem A (1) and (2-2).

Theorem B has some aspect related to Obata's theorem [8] on the first non-zero eigenvalue on Einstein spaces.

THEOREM C. *Let (M, g) be a compact orientable Riemannian manifold with $m = 6$. In (1.2), $a_2 = a_3 = 0$ holds, if and only if (M, g) is either*

(1) E^6/Γ_1 , where Γ_1 is some discontinuous group of translations of the 6-dimensional Euclidean space E^6 , or

(2) $[S^3(c) \times H^3(-c)]/\Gamma_2$, where Γ_2 is some discontinuous group of isometries of $S^3(c) \times H^3(-c)$.

For $m \leq 5$, McKean-Singer [6] and Patodi [10] showed that (1) is the only case. For $m \geq 7$, see Proposition 7 (3).

Kählerian analogues are also true. Corresponding to Theorem A, we have Theorem E in §4. Corresponding to Theorem B, we have

THEOREM D. *Let (M, g, J) be a compact Kählerian manifold, $m = 2n \leq 12$. Let $(CP^n(H), g_0, J_0)$ be a complex n -dimensional projective space with the Fubini-Study metric of constant holomorphic sectional curvature H . If $\text{Spec}(M, g, J) = \text{Spec}(CP^n(H), g_0, J_0)$, then (M, g, J) is holomorphically isometric to $(CP^n(H), g_0, J_0)$.*

2. Preliminaries. By $R = (R^i_{jkl})$ we denote the Riemannian curvature tensor: $R^i_{jkl}\partial/\partial x^j = R(\partial/\partial x^k, \partial/\partial x^l)\partial/\partial x^i$ and $R(X, Y)Z = \nabla_{[X, Y]}Z - [\nabla_X, \nabla_Y]Z$, $i, j, k, l = 1, \dots, m = \dim M$. By $R_1 = (R_{jk}) = (R^r_{jkr})$ we denote the Ricci curvature tensor. By $S = (g^{jk}R_{jk})$ we denote the scalar curvature. For a tensor field $T = (T_{ijk})$, for example, we denote $|T|^2 = (T_{ijk}T^{ijk})$. Then we have (cf. [1], [2], [6], [11])

$$(2.1) \quad a_0 = \text{Vol}(M) = \int dM,$$

$$(2.2) \quad a_1 = \frac{1}{6} \int S dM,$$

$$(2.3) \quad a_2 = \frac{1}{360} \int [2|R|^2 - 2|R_1|^2 + 5S^2] dM,$$

$$\begin{aligned}
 (2.4) \quad \alpha_3 = & \frac{1}{6!} \iint \left[-\frac{1}{9} |\nabla R|^2 - \frac{26}{63} |\nabla R_1|^2 - \frac{142}{63} |\nabla S|^2 \right. \\
 & - \frac{8}{21} R^{ij}{}_{kl} R^{kl}{}_{rs} R^{rs}{}_{ij} - \frac{8}{63} R^{rs} R_r{}^{jkl} R_{s,jkl} + \frac{2}{3} S |R|^2 \\
 & \left. - \frac{20}{63} R^{ik} R^{jl} R_{ijkl} - \frac{4}{7} R^i{}_j R^j{}_k R^k{}_i - \frac{2}{3} S |R_1|^2 + \frac{5}{9} S^3 \right] dM.
 \end{aligned}$$

The following are also useful.

$$(2.5) \quad |R|^2 - \frac{2}{m-1} |R_1|^2 \geq 0,$$

$$(2.6) \quad |R_1|^2 - \frac{1}{m} S^2 \geq 0.$$

The equality in (2.5) on M implies that (M, g) is of constant curvature, and the equality in (2.6) on M implies that (M, g) is an Einstein space (cf. [1], or [2]).

The Weyl's conformal curvature tensor $C = (C^i{}_{jkl})$, $C_{ijkl} = g_{ir} C^r{}_{jkl}$, is given by (for $m \geq 4$)

$$\begin{aligned}
 (2.7) \quad C_{ijkl} = & R_{ijkl} - \frac{1}{m-2} (R_{jk}g_{il} - R_{jl}g_{ik} + g_{jk}R_{il} - g_{jl}R_{ik}) \\
 & + \frac{1}{(m-1)(m-2)} (g_{jk}g_{il} - g_{jl}g_{ik})S.
 \end{aligned}$$

Then we have (cf. [14], [15])

$$(2.8) \quad |C|^2 = |R|^2 - \frac{4}{m-2} |R_1|^2 + \frac{2}{(m-1)(m-2)} S^2 \geq 0.$$

By (2.8) we have

$$(2.9) \quad 2|R|^2 - 2|R_1|^2 + 5S^2 = 2|C|^2 + \frac{2(6-m)}{m-2} |R_1|^2 + \frac{5m(m-3) + 6}{(m-1)(m-2)} S^2.$$

3. Geometry reflected by the spectrum. Let (M, g) and (M', g') be compact orientable Riemannian manifolds. The following are known.

[i] $\text{Spec}(M, g) = \text{Spec}(M', g')$ implies $m = m'$, $\text{Vol}(M) = \text{Vol}(M')$ ([2], p. 215).

[ii] For $m = m' = 2$, if $a_\beta = a'_\beta$ ($\beta = 0, 1, 2$) and $S = \text{constant}$, then S' is also constant and $S = S'$ (Berger: [2], p. 226).

[iii] For $m = m' = 3$, if $a_\beta = a'_\beta$ ($\beta = 0, 1, 2$) and if (M, g) is of constant curvature K , then (M', g') is also of constant curvature K (Berger: [2], p. 228).

[iv] For $m = m' = 4$, if $a_\beta = a'_\beta$ ($\beta = 0, 1, 2$), $\chi(M) = \chi(M')$ and if (M, g) is of constant curvature K , then (M', g') is of constant curvature

K (Berger: [2], p. 229).

[v] Assume that (M, g) and (M', g') are Einstein spaces and $a_\beta = a'_\beta$ ($\beta = 0, 1, 2$). Then (M, g) is of constant curvature K , if and only if (M', g') is of constant curvature K (Sakai [11], p. 599).

[vi] For $m = m' = 6$, assume that (M, g) and (M', g') are Einstein spaces. If $a_\beta = a'_\beta$ ($\beta = 0, 1, 2, 3$) and $\chi(M) = \chi(M')$, then (M, g) is locally symmetric if and only if (M', g') is locally symmetric (Sakai [11], p. 601).

[vii] For $m \leq 5$, assume $a_\beta = 0$ ($\beta = 1, 2$). Then (M, g) is locally flat (for $m \leq 3$, Mckean-Singer [6]; for $m \leq 5$, Patodi [10]).

Concerning [iii], [iv] and [v], we have

THEOREM A'. *Let (M, g) and (M', g') be compact orientable Riemannian manifolds. Assume $a_\beta = a'_\beta$ for $\beta = 0, 1, 2$. Then,*

(1) *for $m = m' \leq 5$, (M, g) is of constant curvature K , if and only if (M', g') is of constant curvature K ,*

(2) *for $m = m' = 6$, (M, g) is conformally flat and S is constant, if and only if (M', g') is conformally flat and S' is constant, $S = S'$.*

PROOF. Since the case $m = 2, 3$ was proved in [1], assume $m \geq 4$. By (2.3) and (2.9), $a_2 = a'_2$ is written as

$$\begin{aligned}
 (3.1) \quad & \int \left[2|C|^2 + \frac{2(6-m)}{m-2} \left(|R_1|^2 - \frac{1}{m} S^2 \right) \right. \\
 & \quad \left. + \left(\frac{2(6-m)}{m(m-2)} + \frac{5m(m-3)+6}{(m-1)(m-2)} \right) S^2 \right] dM \\
 & = \int \left[2|C'|^2 + \frac{2(6-m)}{m-2} \left(|R'_1|^2 - \frac{1}{m} S'^2 \right) \right. \\
 & \quad \left. + \left(\frac{2(6-m)}{m(m-2)} + \frac{5m(m-3)+6}{(m-1)(m-2)} \right) S'^2 \right] dM'.
 \end{aligned}$$

First assume $4 \leq m \leq 5$ and (M', g') is of constant curvature K' . Then it is conformally flat ($C' = 0$) and is an Einstein space ($|R'_1|^2 = S'^2/m$). Since S' is constant, $a_0 = a'_0$ and $a_1 = a'_1$ imply $\int S^2 dM \geq \int S'^2 dM'$. In fact, using Schwarz inequality, we have

$$\begin{aligned}
 \left(\int dM \right) \left(\int S^2 dM \right) & \geq \left(\int S dM \right)^2 = \left(\int S' dM' \right)^2 \\
 & = S'^2 \left(\int dM' \right)^2 = S'^2 \left(\int dM \right) \left(\int dM' \right) \\
 & = \left(\int dM \right) \left(\int S'^2 dM' \right).
 \end{aligned}$$

Hence, (3.1) gives $C = 0$, $|R_1|^2 = S^2/m$ and $\int S^2 dM = \int S'^2 dM'$. Consequently,

(M, g) is of constant curvature $K = K'$.

Next, assume $m = 6, C' = 0$ and $S' = \text{constant}$. Then, using (3.1), similarly we have $C = 0$ and $S = \text{constant}$.

REMARK. (3.1) gives a simple proof of [v].

Concerning [vii], a simple proof is (2.3) and (2.9). Patodi [10] gives a counter-example for $m = 6$. For $m = 6$, under an additional condition $a_3 = 0$, we determine (M, g) .

LEMMA 1. *If $a_2 = 0$ and $m = 6$, then (M, g) is conformally flat and the scalar curvature S is vanishing.*

PROOF. This follows from (2.3) and (2.9).

Now we denote $(R_1^3) = (R^i_j R^j_k R^k_i)$.

LEMMA 2. *If (M, g) is conformally flat, $m \geq 4$, and $S = \text{constant}$, then*

$$(3.2) \quad \int |\nabla R_1|^2 = \int \left[\frac{-m}{m-2} (R_1^3) + \frac{2m-1}{(m-1)(m-2)} S |R_1|^2 - \frac{1}{(m-1)(m-2)} S^3 \right] dM.$$

PROOF. By (2.7) and $C = 0, R_{ijkl}$ is expressed by R_{jk}, g_{jk} , and S . By the second Bianchi identity for R_{ijkl} we have

$$(3.3) \quad \nabla_k R_{ij} = \nabla_j R_{ik}.$$

Then we have

$$(3.4) \quad \begin{aligned} |\nabla R_1|^2 &= \nabla_k R_{ij} \nabla^k R^{ij} = \nabla_j R_{ik} \nabla^k R^{ij} \\ &= \nabla^k (\nabla_j R_{ik} \cdot R^{ij}) - \nabla^k \nabla_j R_{ik} \cdot R^{ij}, \end{aligned}$$

where $\nabla^k \nabla_j R_{ik}$ is calculated by the Ricci identity:

$$(3.5) \quad g^{kr} \nabla_r \nabla_j R_{ik} = g^{kr} (\nabla_j \nabla_r R_{ik} - R^s_{ijr} R_{sk} - R^s_{kjr} R_{is}).$$

Noticing that $g^{kr} \nabla_r R_{ik} = (1/2) \nabla_i S = 0$ and $g^{kr} R^s_{kjr} = -R^s_j$, we simplify (3.5). Putting the result into (3.4) and integrating, we have (3.2).

LEMMA 3. *If (M, g) is conformally flat, $m \geq 4$, and $S = \text{constant}$, then*

$$(3.6) \quad \begin{aligned} a_3 &= \frac{1}{6!} \int \left[\frac{24 - 26m}{63(m-2)} |\nabla R_1|^2 + \frac{1}{63(m-2)^3} [192(m-4) \right. \\ &\quad \left. - 16(m-2)(m-4) - 40(m-2)^2 - 36(m-2)^3] (R^3) \right] \end{aligned}$$

$$\begin{aligned}
 & + \frac{1}{63(m-1)(m-2)^3} [576 - 16(m+1)(m-2) \\
 & + 4(m-2)^2(52m-47) - 42(m-1)(m-2)^2] S |R_1|^2 \\
 & + \frac{1}{63(m-1)^2(m-2)^3} [-96m + 16(m-1)(m-2) \\
 & - 104(m-1)(m-2)^2 + 35(m-1)^2(m-2)^3] S^3 \Big] dM .
 \end{aligned}$$

PROOF. By (2.7) and $C = 0$, we have

$$(3.7) \quad |\nabla R|^2 = \frac{4}{m-2} |\nabla R_1|^2 ,$$

$$(3.8) \quad S |R|^2 = \frac{4}{m-2} S |R_1|^2 - \frac{2}{(m-1)(m-2)} S^3 ,$$

$$(3.9) \quad R^{ik} R^{jl} R_{ijkl} = \frac{2}{m-2} (R_1^3) - \frac{2m-1}{(m-1)(m-2)} S |R_1|^2 + \frac{1}{(m-1)(m-2)} S^3 ,$$

$$(3.10) \quad R^{rs} R_r^{jkl} R_{sjkl} = \frac{2(m-4)}{(m-2)^2} (R_1^3) + \frac{2(m+1)}{(m-1)(m-2)^2} S |R_1|^2 - \frac{2}{(m-1)(m-2)^2} S^3 ,$$

$$(3.11) \quad R^{ij}_{kl} R^{kl}_{rs} R^{rs}_{ij} = [\text{replacing } R^{ij}_{kl} \text{ by (2.7)}] R^{kl}_{rs} R^{rs}_{ij} \\
 = \frac{-4}{m-2} R^{rs} R_r^{jkl} R_{sjkl} + \frac{2}{(m-1)(m-2)} S |R|^2 \\
 = -\frac{8(m-4)}{(m-2)^3} (R_1^3) - \frac{24}{(m-1)(m-2)^3} S |R_1|^2 \\
 + \frac{4m}{(m-1)^2(m-2)^3} S^3 .$$

Substituting these into (2.4), we get (3.6).

PROPOSITION 4. *Let (M, g) and (M', g') be compact orientable Riemannian manifolds, $m = m' = 6$. Assume $\alpha_\beta = \alpha'_\beta$ ($\beta = 0, 1, 2, 3$). Then (M', g') is of constant curvature $K' > 0$, if and only if (M, g) is also of constant curvature $K = K' > 0$.*

PROOF. Assume that (M', g') is of constant curvature $K' > 0$. By Theorem A' we have $C = 0$ and $S = \text{constant}$. By (3.6) we have

$$(3.12) \quad \alpha_3 = \frac{1}{6!} \int \left[-\frac{11}{21} |\nabla R_1|^2 - \frac{2}{3} (R_1^3) + \frac{19}{105} S |R_1|^2 + (*) S^3 \right] dM ,$$

where $(*)$ denotes the coefficient of S^3 . By (3.2) we have

$$(3.13) \quad a_3 = \frac{1}{6!} \iint \left[-\frac{5}{63} |\nabla R_1|^2 - \frac{4}{63} S \left(|R_1|^2 - \frac{1}{6} S^2 \right) + (**) S^3 \right] dM.$$

Since (M', g') is of constant curvature, we have $a'_3 = (1/6!) \int (**) S'^3 dM'$. Since $S = S' > 0$, $a_3 = a'_3$ implies that $\nabla R_1 = 0$ and $|R_1|^2 = S^2/6$. Hence, (M, g) is of constant curvature $K = K' > 0$. q.e.d.

Theorem A' and Proposition 4 give a proof of Theorem A.

PROPOSITION 5. *If (M, g) is conformally flat and $S = 0$, and if $m \geq 4$ ($m \neq 8$) and $a_3 = 0$, then $\nabla R_1 = 0$, and hence (M, g) is either*

- (1) *locally flat, or*
- (2) *locally Riemannian product $S^{m/2}(c) \times H^{m/2}(-c)$.*

PROOF. By (3.2) and (3.6), we have

$$a_3 = \frac{1}{6!} \int \frac{2(m-8)(5m^2-2m-48)}{63m(m-2)^2} |\nabla R_1|^2 dM.$$

Since $5m^2 - 2m - 48 > 0$ for $m \geq 4$, if $m \neq 8$, we have $\nabla R_1 = 0$. If (M, g) is irreducible, then (M, g) is an Einstein space. $S = 0$ implies that (M, g) is locally flat. This is a contradiction. If (M, g) is reducible, then it is locally Riemannian product $[E^1 \times S^{m-1}(c)$, or $E^1 \times H^{m-1}(-c)$, or E^m , or $S^r(c) \times H^{m-r}(-c)$] (cf. Kurita [4]). $S = 0$ implies that (M, g) is locally E^m or locally $S^{m/2}(c) \times H^{m/2}(-c)$.

THEOREM C'. *Let (M, g) be a compact orientable Riemannian manifold with $m = 6$. If $a_\beta = 0$ for $\beta = 2, 3$, then (M, g) is either*

- (1) E^6/Γ_1 , or
- (2) $[S^3(c) \times H^3(-c)]/\Gamma_2$.

PROOF. This follows from Lemma 1 and Proposition 5.

REMARK. As for $\sum e^{\lambda \alpha t}$ for $S^3(c) \times [H^3(-c)/\Gamma^*]$, cf. [10], p. 283~285.

Concerning [vii] for $m \geq 7$, we can state

PROPOSITION 6. *Let (M, g) be a compact orientable Riemannian manifold with $a_2 = 0$, $m \geq 7$. If*

$$(3.14) \quad |R_1|^2 \leq \frac{5m(m-3) + 6}{2(m-1)(m-6)} S^2$$

holds on M , then (M, g) is conformally flat, and the equality holds in (3.14) on M .

PROOF. This follows from (2.3) and (2.9).

Hence, as for a_2 , we can summarize the above as follows.

PROPOSITION 7. *Let (M, g) be a compact orientable Riemannian manifold.*

(1) *For $2 \leq m \leq 5, a_2 \geq 0$ holds, equality is only for locally flat (M, g) .*

(2) *For $m = 6, a_2 \geq 0$ holds; (if $a_3 = 0$) equality is only for locally flat (M, g) or locally Riemannian product $S^3(c) \times H^3(-c)$.*

(3) *For $m \geq 7$, if Ricci curvatures are non-negative (or non-positive) on M , then $a_2 \geq 0$ holds; equality is only for locally flat (M, g) .*

PROOF. (1) is [vii] or (2.9). (2) is Theorem C'. We show (3). By the assumption of Ricci curvatures we have

$$(3.15) \quad |R_1|^2 \leq S^2 \leq \frac{5m(m-3)+6}{2(m-1)(m-6)} S^2.$$

Hence, (3.14) holds. Therefore $C = 0$ and (3.15) must be equalities. Thus $S = 0$ and $|R_1|^2 = 0$. This implies $R = 0$.

Next, we show

PROPOSITION 8. *Let (M, g) and (M', g') be compact orientable Riemannian manifold, $m = m' = 4$. Assume $a_\beta = a'_\beta$ for $\beta = 0, 1$, and 2. If (M', g') is an Einstein space, then*

$$(3.16) \quad \chi(M) \leq \chi(M')$$

holds. The equality holds, if and only if (M, g) is also an Einstein space.

PROOF. By Gauss-Bonnet formula we have (cf. [1], (8.1))

$$(3.17) \quad a_2 = \frac{8\pi^2}{45} \chi(M) + \frac{1}{120} \int [2|R_1|^2 + S^2] dM.$$

Since (M', g') is an Einstein space, $a_2 = a'_2$ is written as

$$\begin{aligned} \frac{8\pi^2}{45} \chi(M) + \frac{1}{120} \int \left[2\left(|R_1|^2 - \frac{1}{4} S^2\right) + \frac{3}{2} S^2 \right] dM \\ = \frac{8\pi^2}{45} \chi(M') + \frac{1}{120} \int \frac{3}{2} S'^2 dM'. \end{aligned}$$

Since $\int S^2 dM \geq \int S'^2 dM'$ (cf. proof of Theorem A'), we have (3.16). The equality implies $|R_1|^2 = S^2/4$.

REMARK. In connection with Theorem A' (2) and Proposition 4, the role of $a_3 = a'_3$ may be replaced by the fundamental group of M . Namely we have

PROPOSITION 9. *Let (M, g) and (M', g') be compact orientable Riemannian manifolds, $m = m' = 6$. Assume that $a_\beta = a'_\beta$ holds for $\beta = 0, 1, 2$*

and assume that (M', g') is of constant curvature K' . If the fundamental group $\pi_1(M)$ of M is finite, then (M, g) is also of constant curvature K' and $K' > 0$.

In particular, if $(M', g') = (S^0(c), g_0)$, then (M, g) is isometric to $(S^0(c), g_0)$.

This follows from Theorem A' (2) and the following fact: Let (M, g) be a compact conformally flat Riemannian manifold with finite $\pi_1(M)$, $m \geq 3$; if S is constant, then (M, g) is of positive constant curvature (cf. Tanno [13]).

4. Kählerian manifolds. Let (M, g, J) be a Kählerian manifold with almost complex structure tensor $J = (J_j^i)$ and Kählerian metric tensor $g = (g_{ij})$. The complex dimension of M is $n = m/2$. Then,

$$(4.1) \quad g_{rs} J_i^r J_j^s = g_{ij}, \quad J_r^i J_j^r = -\delta_j^i$$

and $\nabla_k J_j^i = 0$. $J_{ij} = g_{ir} J_j^r$ is skew-symmetric. The Ricci curvature tensor satisfies

$$(4.2) \quad R_{ij} J_r^i J_s^j = R_{rs}, \quad R_{ir} J_j^r = -R_{jr} J_i^r.$$

The Bochner curvature tensor $B = (B^i_{jkl})$, $B_{ijkl} = g_{ir} B^r_{jkl}$, is given by (cf. [15], etc.)

$$(4.3) \quad B_{ijkl} = R_{ijkl} - \frac{1}{m+4} (R_{jk} g_{il} - R_{jl} g_{ik} + g_{jk} R_{il} - g_{jl} R_{ik} + R_{jr} J_k^r J_{il} - R_{jr} J_i^r J_{jk} + J_{jk} R_{ir} J_l^r - J_{jl} R_{ir} J_k^r - 2R_{kr} J_l^r J_{ij} - 2R_{ir} J_j^r J_{kl}) + \frac{1}{(m+2)(m+4)} (g_{jk} g_{il} - g_{jl} g_{ik} + J_{jk} J_{il} - J_{jl} J_{ik} - 2J_{kl} J_{ij}) S.$$

$|B|$ is given by (cf. [15])

$$(4.4) \quad |B|^2 = |R|^2 - \frac{16}{m+4} |R_1|^2 + \frac{8}{(m+2)(m+4)} S^2.$$

A Kählerian manifold (M, g, J) , $m \geq 4$, is of constant holomorphic sectional curvature H , if and only if

$$(4.5) \quad R_{ijkl} = \frac{H}{4} (g_{jk} g_{il} - g_{jl} g_{ik} + J_{jk} J_{il} - J_{ik} J_{jl} - 2J_{ij} J_{kl})$$

holds. Then R_{jk} and S are given by

$$(4.6) \quad R_{jk} = \frac{m+2}{4} H g_{jk}, \quad S = \frac{m(m+2)}{4} H.$$

A Kählerian manifold (M, g, J) is of constant holomorphic sectional curvature, if and only if $B = 0$ and $|R_1|^2 = S^2/m$.

PROPOSITION 10. *Let (M, g, J) and (M', g', J') be compact Kählerian manifolds. Assume $a_\beta = a'_\beta$ for $\beta = 0, 1,$ and 2 . Then,*

(1) *for $m = m' \leq 10$, (M, g, J) is of constant holomorphic sectional curvature, if and only if (M', g', J') is of constant holomorphic sectional curvature, $H = H'$,*

(2) *for $m = m' = 12$, $B = 0$ and $S =$ constant, if and only if $B' = 0$ and $S' =$ constant, $S = S'$.*

PROOF. By (4.4), we have

$$(4.7) \quad a_2 = \frac{1}{360} \int \left[2|B|^2 + \frac{2(12-m)}{m+4} \left(|R_1|^2 - \frac{1}{m} S^2 \right) + \frac{5m^2 + 8m + 12}{m(m+2)} S^2 \right] dM.$$

Then the proof is completed in a way similar to that of Theorem A' .

LEMMA 11. (Matsumoto [5]) *If the Bochner curvature tensor $B = 0$ (more generally, parallel) and $S =$ constant, then the Ricci curvature tensor R_i is parallel (and (M, g) is locally symmetric).*

LEMMA 12. *If $\nabla R_1 = 0$, then*

$$(4.8) \quad R_{rjks}R^{rs} = R_{jr}R^r_k.$$

PROOF. By $0 = (\nabla_s \nabla_k R_{rj} - \nabla_k \nabla_s R_{rj})g^{rs}$ and the Ricci identity, we have (4.8).

LEMMA 13. *If $B = 0$ and $\nabla R_1 = 0$, then*

$$(4.9) \quad m(R_i^3) = \frac{2(m+1)}{m+2} S |R_1|^2 - \frac{1}{m+2} S^3.$$

PROOF. Transvect (4.8) with R^{jk} and use (4.3). Then, using (4.2), we have (4.9).

LEMMA 14. *If $B = 0$ and $\nabla R_1 = 0$, then*

$$(4.10) \quad S |R|^2 = \frac{16}{m+4} S |R_1|^2 - \frac{8}{(m+2)(m+4)} S^3,$$

$$(4.11) \quad R^{ik}R^{jl}R_{ijkl} = -(R_1^3),$$

$$(4.12) \quad R^{rs}R_{rjkl}R_s^{jkl} = \frac{16}{m+4} (R_1^3) - \frac{8}{(m+2)(m+4)} S |R_1|^2,$$

$$(4.13) \quad R^{ij}_{kl}R^{kl}_{rs}R^{rs}_{ij} = -\frac{16(m+12)}{(m+4)^2}(R_1^3) + \frac{8(m+20)}{(m+2)(m+4)^2}S|R_1|^2 - \frac{32}{(m+2)^2(m+4)^2}S^3.$$

PROOF. (4.10) follows from (4.4). (4.11) follows from (4.8). (4.12) is calculated as follows: First write down $R^{rs}R_{rjkl}$ using (4.3). Next, transvect it with R_s^{jkl} and use (4.2) and well known identities:

$$(4.14) \quad R_{ijkl}J^{kl} = 2J_i^r R_{rj}, \quad R_{ijkl}J_r^k J_s^l = R_{ijrs}, \quad \text{etc.}$$

For example, $R_s^{jkl}R_{kt}J_l^t R_s^h J_j^h (= -2(R_1^3))$ is calculated by the first Bianchi identity and the above relations.

(4.13) is calculated as follows (using (4.2), (4.8), (4.14))

$$\begin{aligned} & \text{[replacing } R_{ijkl} \text{ by (4.3)] } R^{ij}_{rs}R^{rskl} \\ &= \frac{1}{m+4} [-8R^{rs}R_{rjkl}R_s^{jkl} - 16(R_1^3)] \\ & \quad - \frac{1}{(m+2)(m+4)} [-4|R|^2 - 8|R_1|^2]S. \end{aligned}$$

Substituting (4.10) and (4.12), we have (4.13).

LEMMA 15. *If $B = 0$ and $\nabla R_1 = 0$, then*

$$(4.15) \quad a_3 = \frac{1}{6!} \int \left[\left(\frac{128(3m^2 + 40m + 64)}{63m(m+2)(m+4)^2} - \frac{32(m+1)}{63m(m+2)} + \frac{2(12-m)}{3(m+4)} \right) S|R_1|^2 + \langle * \rangle S^3 \right] dM.$$

PROOF. In (2.4) we substitute (4.10), ..., (4.13) and next eliminate (R_1^3) using (4.9).

PROPOSITION 16. *Let (M, g, J) and (M', g', J') be compact Kählerian manifolds, $m = m' = 12$. Assume $a_\beta = a'_\beta$ for $\beta = 0, 1, 2$, and 3. Then (M', g', J') is of constant holomorphic sectional curvature $H' \neq 0$, if and only if (M, g, J) is of constant holomorphic sectional curvature $H = H'$.*

PROOF. Assume that (M', g', J') is of constant holomorphic sectional curvature $H' \neq 0$. By proposition 10 and Lemmas 11 and 15, $a_3 = a'_3$ is written as

$$(4.16) \quad \begin{aligned} & \frac{1}{6!} \int \left[\frac{1}{147} S \left(|R_1|^2 - \frac{1}{12} S^2 \right) + \langle ** \rangle S^3 \right] dM \\ &= \frac{1}{6!} \int \langle ** \rangle S'^3 dM'. \end{aligned}$$

$S = S' \neq 0$ gives $|R_1|^2 = S^2/12$. Hence, (M, g, J) is of constant holomorphic sectional curvature H and $H = H'$.

By $CD^n(-H)$ we denote a simply connected complex space form (of constant holomorphic sectional curvature $-H < 0$) of complex dimension n .

PROPOSITION 17. *Let (M, g, J) be a compact Kählerian manifold with dimension $m = 2n \leq 12$. $a_2 = 0$ holds good, if and only if (M, g, J) is either*

(1) CE^n/Γ_3 , where Γ_3 is some discontinuous group of automorphisms of the complex n -dimensional Euclidean space CE^n , or

(2) $m = 2n = 12$ and $[CP^3(H) \times CD^3(-H)]/\Gamma_4$, where Γ_4 is some discontinuous group of automorphisms of $CP^3(H) \times CD^3(-H)$.

PROOF. The case $m = 2n \leq 10$ is clear from (4.7). For $m = 2n = 12$, by (4.7) we have $B = 0$ and $S = 0$. By Lemma 11, we have $\forall R_1 = 0$. Since $S = 0$, (M, g, J) is not irreducible. Hence, it is reducible and locally $[CE^6$ or $CP^r(H) \times CD^s(-H)$, $r + s = 6$] (cf. Takagi-Watanabe [12]). $S = 0$ gives $r = s = 3$.

Finally we combine Proposition 10 and Proposition 16.

THEOREM E. *Let (M, g, J) and (M', g', J') be compact Kählerian manifolds, $m = m' \leq 12$. Assume $\text{Spec}(M, g, J) = \text{Spec}(M', g', J')$.*

(1) *For $m \leq 10$, (M, g, J) is of constant holomorphic sectional curvature H , if and only if (M', g', J') is of constant holomorphic sectional curvature $H' = H$.*

(2) *For $m = 12$, (M, g, J) is of constant holomorphic sectional curvature $H \neq 0$, if and only if (M', g', J') is of constant holomorphic sectional curvature $H' = H$.*

By Theorem E we have Theorem D in the introduction.

REFERENCES

- [1] M. BERGER, Le spectre des variétés riemanniennes, Rev. Roum. Math. Pure et Appl., 13 (1969), 915-931.
- [2] M. BERGER, P. GAUDUCHON et E. MAZET, Le Spectre d'une Variété riemannienne, Lect. Notes in Math. Springer-Verlag, 194.
- [3] M. KAC, Can one hear the shape of a drum? Amer. Math. Monthly, 73 (April, 1966), 1-23.
- [4] M. KURITA, On the holonomy group of the conformally flat Riemannian manifold, Nagoya Math. J., 9 (1955), 161-171.
- [5] M. MATSUMOTO, On Kählerian spaces with parallel or vanishing Bochner curvature tensor, Tensor (N.S.), 20 (1969), 25-28.
- [6] H. F. MCKEAN AND I. M. SINGER, Curvature and the eigenvalues of the Laplacian, J. Diff. Geometry, 1 (1967), 43-69.

- [7] J. MILNOR, Eigenvalues of the Laplace operator on certain manifolds, Proc. Nat. Sci. U.S.A., 51 (1964), 542.
- [8] M. OBATA, Certain conditions for a Riemannian manifold to be isometric with a sphere, J. Math. Soc. Japan, 14 (1962), 333-340.
- [9] V. K. PATODI, Curvature and the eigenforms of the Laplace operator, J. Diff. Geometry, 5 (1971), 233-249.
- [10] V. K. PATODI, Curvature and the fundamental solution of the heat operator, J. Indian Math. Soc., 34 (1970), 269-285.
- [11] T. SAKAI, On eigenvalues of Laplacian and curvature of Riemannian manifold, Tôhoku Math. J., 23 (1971), 589-603.
- [12] H. TAKAGI AND Y. WATANABE, On the holonomy groups of Kählerian manifolds with vanishing Bochner curvature tensor, Tôhoku Math. J.,
- [13] S. TANNO, Compact conformally flat Riemannian manifolds, to appear in J. Diff. Geometry, 8 (1973).
- [14] S. TANNO, Euler-Poincaré characteristics and curvature tensors, Tôhoku Math. J., 25 (1973), 33-52.
- [15] S. TANNO, An inequality for 4-dimensional Kählerian manifolds, Proc. Japan Acad., 49 (1973), 257-261.

MATHEMATICAL INSTITUTE
TÔHOKU UNIVERSITY

