

# Relative Property (T) and the Vanishing of the first $\ell^2$ -Betti number

Talia Fernós

## Abstract

In this paper, we show that certain families with relative property (T) have trivial first  $\ell^2$ -Betti number. We apply this to the elementary matrix group  $EL_n(\mathcal{R})$  where  $\mathcal{R}$  is any countable unital ring of characteristic 0.

## 1 Introduction

The notion of  $\ell^2$ -Betti numbers has been studied for a variety of objects from groups to measurable foliations [Ati76], [CG86], [Con79]. They were studied by D. Gaboriau who showed that if the (probability measure preserving and free) actions of two groups are orbit equivalent then their  $\ell^2$ -Betti numbers are the same [Gab02]. Namely,  $\ell^2$ -Betti numbers are an invariant of orbit equivalence.

A generalization of the usual Betti numbers,  $\ell^2$ -Betti numbers count the “module-dimension” of certain infinite dimensional cohomology groups. See [Gab00] for a brief introduction to  $\ell^2$ -Betti numbers for groups and equivalence relations.

Our objective here is to prove the following:

**Theorem 1.1.** *Let  $(\Gamma, A)$  be a group pair with relative property (T). If  $A$  is infinite and weakly normal or weakly quasi-normal in  $\Gamma$  then  $\beta_1(\Gamma) = 0$ .*

**Remark:** In the case when  $A$  is normal in  $\Gamma$  the above theorem follows from Proposition 3.5.1 in Florian Martin’s thesis [Mar03]. The proposition reads: if  $G$  is locally compact and  $N$  is a closed normal subgroup such that  $(G, N)$  has relative (T), then  $H^1(G, \pi) = 0$  for every representation  $\pi$  without  $N$ -invariant vectors.

Recall the following two definitions:

---

Received by the editors September 2009.

Communicated by A. Valette.

**Definition 1.2.** A subgroup  $A \leq \Gamma$  is said to be weakly normal if there exists a well ordered (by inclusion) family of intermediate subgroups  $\{N_t : 0 \leq t \leq \tau\}$  of  $\Gamma$  with  $N_0 = A$  and  $N_\tau = \Gamma$  and such that for each  $0 < t \leq \tau$  the following inclusion is normal:

$$\bigcup_{s < t} N_s \trianglelefteq N_t$$

**Definition 1.3** (Popa). A subgroup  $A \leq \Gamma$  is said to be weakly quasi-normal if there exists a well ordered (by inclusion) family of intermediate subgroups  $\{N_t : 0 \leq t \leq \tau\}$  of  $\Gamma$  with  $N_0 = A$  and  $N_\tau = \Gamma$  and

- for each non-limit ordinal  $0 < t \leq \tau$  we have  $N_t = \langle \gamma : |N_{t-1} \cap \gamma N_{t-1} \gamma^{-1}| = \infty \rangle$ ;
- for each limit ordinal  $0 < t \leq \tau$  we have  $N_t = \bigcup_{s < t} N_s$ .

Since many of the family of examples of nontrivial group pairs with relative property (T) are of the form  $(\Gamma, A)$  where  $A$  is normal and amenable [Val05], [Fer06a], Theorem 1.1 can be seen as a natural extension of the following:

**Theorem 1.4.** [BV97, Corollary 6, Theorem D] If  $\Gamma$  is a group satisfying either of the following two conditions then  $\beta_1(\Gamma) = 0$ :

1.  $\Gamma$  has property (T);
2.  $\Gamma$  is finitely generated and contains a normal amenable subgroup.

Examples of groups having  $\beta_1 > 0$  are lattices (both uniform and nonuniform) in  $\mathrm{PSL}_2(\mathbb{R})$  [BV97] Example p. 318. For a large class of examples see [JP07].

I would like to thank Alain Valette for reading this and making useful suggestions. This work was part of my doctoral thesis.

## 2 Some preliminaries

The celebrated theorem of Delorme and Guichardet says that property (T) is equivalent to property FH. The same is true for relative property (T) and it is through this fact that we will prove our theorem.

**Definition 2.1.** A group pair  $(\Gamma, A)$  has relative property FH if every representation  $\alpha : \Gamma \rightarrow \mathrm{Isom}(\mathcal{H})$  on a Hilbert space  $\mathcal{H}$ , has an  $A$ -fixed point  $v \in \mathcal{H}$ .

We will be thinking of the affine Hilbert spaces as vector spaces with an affine structure. By this we mean that we choose once and for all, an origin  $0 \in \mathcal{H}$ . This is equivalent to choosing an isomorphism  $\mathrm{Isom}(\mathcal{H}) \cong \mathcal{U}(\mathcal{H}) \ltimes \mathcal{H}$ .

**Theorem 2.2** ([Jol05],[dC06],[Fer06b]). A group pair  $(\Gamma, A)$  has relative property (T) if and only if it has relative property FH.

## 2.1 Affine Actions and Relative Cohomology

Let  $\mathcal{H}$  be a Hilbert space and  $\pi: \Gamma \rightarrow \mathcal{U}(\mathcal{H})$  be a unitary representation of  $\Gamma$ . We will need to recall some definitions:

**Definition 2.3.** 1. The space of  $\pi$ -cocycles is:

$$Z^1(\Gamma, \pi) := \{b: \Gamma \rightarrow \mathcal{H} \mid b(\gamma_1\gamma_2) = b(\gamma_1) + \pi(\gamma_1)b(\gamma_2)\}$$

2. The space of  $\pi$ -relative-coboundaries is:

$$B^1(\Gamma, A, \pi) := \{b \in Z^1 \mid b(a) = \pi(a)v - v \text{ for some } v \in \mathcal{H} \text{ and for all } a \in A\}.$$

3. The first cohomology is the quotient:

$$H^1(\Gamma, \pi) := Z^1(\Gamma, \pi) / B^1(\Gamma, \Gamma, \pi).$$

4. The relative first cohomology is the quotient:

$$H^1(\Gamma, A, \pi) := Z^1(\Gamma, \pi) / B^1(\Gamma, A, \pi).$$

Observe that to each cocycle  $b \in Z^1(\Gamma, \pi)$  there corresponds an isometric action

$$\alpha_b(\gamma)(v) := \pi(\gamma)v + b(\gamma).$$

The cocycle identity assures that this is an action. We will refer to this action as the one induced by  $b$  and will often drop the subscript if there is no ambiguity. Conversely, having fixed an origin  $0 \in \mathcal{H}$ , any isometric action  $\alpha: \Gamma \rightarrow \text{Isom}(\mathcal{H})$  gives rise to the map  $b_\alpha(\gamma) := \alpha(\gamma)(0)$  which is guaranteed to be a cocycle by the fact that the action is affine.

The first cohomology  $H^1(\Gamma, \pi)$  is developed in order to understand what kinds of affine  $\Gamma$ -actions can arise with linear part  $\pi$ , modulo the trivial ones (i.e. those arising out of the coboundaries in  $B^1(\Gamma, \Gamma, \pi)$ ). Analogously, one may understand  $H^1(\Gamma, A, \pi)$  as the possible ways of assembling an isometric action of  $\Gamma$ , with linear part  $\pi$  which are trivial relative to  $A$ . It is an easy exercise to show, employing the Lemma of The Center [BdlHV08], that a cocycle is a coboundary on  $A$  exactly when it is bounded on  $A$ .

We can now rephrase Theorem 2.2 in terms of relative cohomology:

**Theorem 2.4.** A group pair  $(\Gamma, A)$  has relative property (T) if and only if  $H^1(\Gamma, A, \pi) = 0$  for every unitary  $\Gamma$ -representation  $\pi$ .

### 3 Proof of Theorem 1.1

Theorem 1.1 follows from three results<sup>1</sup>:

**Theorem 3.1** ([BV97]Corollary 4). *Let  $\lambda_\Gamma: \Gamma \rightarrow \mathcal{U}(\ell^2(\Gamma))$  be the left regular representation of  $\Gamma$ . If  $\Gamma$  is nonamenable and  $H^1(\Gamma, \lambda_\Gamma) = 0$  then  $\beta_1(\Gamma) = 0$ .*

**Proposition 3.2.** *Let  $(\Gamma, A)$  be a group-subgroup pair, and  $A$  infinite and weakly normal in  $\Gamma$ . If  $\pi: \Gamma \rightarrow \mathcal{U}(\mathcal{H})$  is a unitary representation with no nontrivial  $A$ -invariant vectors and  $H^1(\Gamma, A, \pi) = 0$  then  $H^1(\Gamma, \pi) = 0$ .*

**Proposition 3.3.** *Let  $(\Gamma, A)$  be a group-subgroup pair, and  $A$  infinite and weakly quasi-normal in  $\Gamma$ . If  $\pi: \Gamma \rightarrow \mathcal{U}(\mathcal{H})$  is a  $C_0$ -unitary representation and  $H^1(\Gamma, A, \pi) = 0$  then  $H^1(\Gamma, \pi) = 0$ .*

We now prove these propositions, first in the case where  $A$  is weakly normal and then when it is weakly quasi-normal in  $\Gamma$ .

#### 3.1 $A$ weakly normal

Under the same assumptions as Proposition 3.2, we prove the following two lemmas, the last of which proves the proposition.

**Lemma 3.4.** *If  $A \leq N \leq \Gamma$  and  $b \in B^1(\Gamma, N, \pi)$  then  $b \in B^1(\Gamma, N', \pi)$  for any subgroup  $N'$  of the normalizer of  $N$  in  $\Gamma$ .*

*Proof.* Suppose that  $b \in B^1(\Gamma, N, \pi)$  and  $v \in \mathcal{H}$  such that  $b(a) = \pi(a)v - v$  for every  $a \in N$ . Now, fix  $\gamma \in N'$  and we have:

$$\begin{aligned} b(\gamma^{-1}a\gamma) &= b(\gamma^{-1}a) + \pi(\gamma^{-1}a)b(\gamma) \\ &= b(\gamma^{-1}) + \pi(\gamma^{-1})b(a) + \pi(\gamma^{-1}a)b(\gamma) \\ = \pi(\gamma^{-1}a\gamma)v - v &= b(\gamma^{-1}) + \pi(\gamma^{-1})(\pi(a)v - v) + \pi(\gamma^{-1}a)b(\gamma) \end{aligned}$$

Recalling the fact that  $b(\gamma^{-1}) = -\pi(\gamma^{-1})b(\gamma)$  is a consequence of the cocycle identity, we multiply the last line above by  $\pi(\gamma)$  and get:

$$\pi(a)(v + b(\gamma) - \pi(\gamma)v) = v + b(\gamma) - \pi(\gamma)v.$$

Since  $\pi$  has no nontrivial  $A$ -invariant vectors, and the last equation holds for every  $a \in N \geq A$  it follows that  $v + b(\gamma) - \pi(\gamma)v = 0$ , that is to say  $b(\gamma) = \pi(\gamma)v - v$  and therefore,  $b \in B^1(\Gamma, N', \pi)$ . ■

**Lemma 3.5.** *For every  $b \in Z^1(\Gamma, \pi)$  either  $b \in B^1(\Gamma, \Gamma, \pi)$  or  $b \notin B^1(\Gamma, A, \pi)$ .*

*Proof.* Suppose that  $b \notin B^1(\Gamma, \Gamma, \pi)$ . Then the set  $\{t: b \notin B^1(\Gamma, N_t, \pi)\}$  contains  $\tau$  (recall that  $N_\tau = \Gamma$ ) and therefore the set is nonempty and must have a minimal element  $t_0$ . If  $t_0 = 0$  then we are done. Otherwise,  $b \in B^1(\Gamma, \bigcup_{s < t_0} N_s, \pi)$  but  $b \notin B^1(\Gamma, N_{t_0}, \pi)$  which contradicts the previous lemma since  $\bigcup_{s < t_0} N_s \triangleleft N_{t_0}$ . ■

<sup>1</sup>It is well known that for  $(\Gamma, A)$  to have relative property (T), with  $A$  non-compact, it is necessary for  $\Gamma$  to be nonamenable. See for example [Fer06a].

### 3.2 wq-Normality

Under the same assumptions, the following proves Proposition 3.3.

**Lemma 3.6.** *Let  $A \leq \Gamma$  be a wq-normal subgroup, with  $A$  infinite and  $\pi: \Gamma \rightarrow \mathcal{U}(\mathcal{H})$  a  $C_0$ -representation. Then for every  $b \in Z^1(\Gamma, \pi)$  either  $b \in B^1(\Gamma, \Gamma, \pi)$  or  $b \notin B^1(\Gamma, A, \pi)$ .*

*Proof.* Let  $\{N_t : 0 \leq t \leq \tau\}$  be the well ordered collection of subgroups guaranteed by the definition of weak quasi-normality for  $A \leq \Gamma$ .

Let  $b \notin B^1(\Gamma, \Gamma, \pi)$ . We aim to show that  $b \notin B^1(\Gamma, A, \pi)$ . Consider the set  $\{t : b \notin B^1(\Gamma, N_t, \pi)\}$ . As this set contains  $\tau$  it is not empty and hence contains a minimal element  $t_0$ . If  $t_0 = 0$ , then we are done. By contradiction suppose that  $t_0 > 0$ .

We claim that  $t_0$  is not a limit ordinal. If  $t_0$  is a limit ordinal, then  $N_{t_0} = \bigcup_{t < t_0} N_t$ .

Let  $v_t \in \mathcal{H}$  be the vector making  $b|_{N_t}(\gamma) = \pi(\gamma)v_t - v_t$  a relative coboundary on each  $N_t$ . Then  $v_t - v_0$  is an invariant vector for the infinite subgroup  $N_0 = A$  and since the representation  $\pi$  is  $C_0$  it must be that  $v_t = v_0$ , which shows that the restriction of  $b$  to  $N_{t_0}$  is a coboundary, contradicting the minimality of  $t_0$ .

Therefore  $t_0$  is not a limit ordinal and by minimality, the restriction of  $b$  to  $N_{t_0-1}$  must be a coboundary.

Recall that  $N_{t_0}$  is generated by a set  $S_{t_0}$  where every  $h \in S_{t_0}$  has the property that

$$|N_{t_0-1} \cap hN_{t_0-1}h^{-1}| = \infty.$$

We now show that there is a  $v \in \mathcal{H}$  such that  $b(h) = \pi(h)v - v$  for each  $h \in S_{t_0}$ . The cocycle relation then implies that  $b \in B^1(\Gamma, N_{t_0}, \pi)$  contradicting the minimality of  $t_0$ .

Let  $v \in \mathcal{H}$  such that  $b|_{N_{t_0-1}}(\gamma) = \pi(\gamma)v - v$ . For  $h \in S_{t_0}$  consider  $\bar{\gamma} \in hN_{t_0-1}h^{-1} \cap N_{t_0-1}$ . Then,  $\bar{\gamma} = h\gamma h^{-1}$  for some  $\gamma \in N_{t_0-1}$  and so

$$\begin{aligned} \pi(\bar{\gamma})v - v &= b(\bar{\gamma}) = b(h\gamma h^{-1}) \\ &= b(h) + \pi(h)(\pi(\gamma)v - v) - \pi(h)(\pi(\gamma)\pi(h^{-1})b(h)) \\ &= b(h) + \pi(\bar{\gamma})\pi(h)v - \pi(h)v - \pi(\bar{\gamma})b(h). \end{aligned}$$

Rearranging terms we have that

$$\pi(\bar{\gamma})(\pi(h)v - v - b(h)) = \pi(h)v - v - b(h).$$

Of course this holds for infinitely many  $\bar{\gamma} \in hN_{t_0-1}h^{-1} \cap N_{t_0-1}$  and since  $\pi$  is  $C_0$  we conclude that  $b(h) = \pi(h)v - v$ .  $\blacksquare$

#### 4 Application to $\mathrm{EL}_n(\mathcal{R})$ for $n \geq 3$

Let  $\mathcal{R}$  be a countable (not necessarily commutative) ring with unit and characteristic 0. Consider the group  $\mathrm{EL}_n(\mathcal{R})$  generated by elementary matrices and  $n \geq 3$ . We have many options here for a weakly quasi-normal subgroup. For example, since  $1 \in \mathcal{R}$  it follows that  $\mathrm{EL}_n(\mathcal{R}) \geq \mathrm{SL}_n(\mathbb{Z})$ . Instead, we choose  $A = \langle E_{i,n}(1) : i = 1, \dots, n-1 \rangle \cong \mathbb{Z}^{n-1}$  to be the subgroup generated by the elementary matrices corresponding to the last column. Of course,  $A$  is contained in a copy of  $\mathrm{SL}_{n-1}(\mathbb{Z}) \times A$  inside  $\mathrm{EL}_n(\mathcal{R})$  and therefore,  $(\mathrm{EL}_n(\mathcal{R}), A)$  has relative property (T).

Consider

$$N_1 = \langle E_{i,j}(r) : r \in \mathcal{R}, i = 1, \dots, n-1, j = 1, \dots, n, \text{ and } i \neq j \rangle \cong \mathrm{EL}_{n-1}(\mathcal{R}) \times \mathcal{R}^{n-1}$$

and

$$N_2 = \langle E_{i,j}(r) : r \in \mathcal{R}, i = 1, \dots, n, j = 1, \dots, n, \text{ and } i \neq j \rangle \cong \mathrm{EL}_n(\mathcal{R}).$$

For each  $i_0$ , we see that the infinite cyclic subgroup  $A_{i_0} = \langle E_{i_0,n}(1) \rangle$  is centralized by

$$C_{i_0} := \{E_{i_0,j}(r) : r \in \mathcal{R}, j = 1, \dots, n-1, \text{ and } j \neq i_0\} \cup \{E_{i,n}(r) : r \in \mathcal{R} \text{ and } i \neq i_0\}.$$

This means that  $\gamma A \gamma^{-1} \cap A \supset A_{i_0}$  for each  $\gamma \in C_{i_0}$  and hence these intersections are infinite.

Now, as  $N_2$  is generated by  $N_1$  together with  $\{E_{n,j}(r) : r \in \mathcal{R} \text{ and } j = 1, \dots, n-1\}$  we need only see that  $E_{n,j}(\cdot)$  centralizes some infinite subgroup of  $N_1$  for each  $j \neq n$ . But of course, as before,  $E_{n,j}(\cdot)$  commutes with  $E_{i,j}(\cdot)$  for any  $i \neq n, j$ .

**Corollary 4.1.** *For  $n \geq 3$  and  $\mathcal{R}$  a unital ring of characteristic 0, the first  $\ell^2$ -Betti number  $\beta_1(\mathrm{EL}_n(\mathcal{R})) = 0$ .*

#### References

- [Ati76] Michael F. Atiyah. Elliptic operators, discrete groups and von Neumann algebras. pages 43–72. *Astérisque*, No. 32–33, 1976.
- [BdlHV08] Bachir Bekka, Pierre de la Harpe, and Alain Valette. *Kazhdan's property (T)*, volume 11 of *New Mathematical Monographs*. Cambridge University Press, Cambridge, 2008.
- [BV97] Mohammed E. B. Bekka and Alain Valette. Group cohomology, harmonic functions and the first  $L^2$ -Betti number. *Potential Anal.*, 6(4):313–326, 1997.
- [CG86] Jeff Cheeger and Mikhael Gromov.  $L_2$ -cohomology and group cohomology. *Topology*, 25(2):189–215, 1986.
- [Con79] Alain Connes. Sur la théorie non commutative de l'intégration. 725:19–143, 1979.

- [dC06] Yves de Cornulier. Relative Kazhdan property. *Ann. Sci. École Norm. Sup. (4)*, 39(2):301–333, 2006.
- [Fer06a] Talia Fernós. Relative property (T) and linear groups. *Ann. Inst. Fourier (Grenoble)*, 56(6):1767–1804, 2006.
- [Fer06b] Talia Fernós. *Relative Property (T), Linear Groups, and Applications*. PhD thesis, University of Illinois, Chicago, 2006.
- [Gab00] Damien Gaboriau. On Orbit Equivalence of Measure Preserving Actions. *Rigidity in Dynamics and Geometry*, pages 167–186, 2000.
- [Gab02] Damien Gaboriau. Invariants  $\ell^2$  de relations d'équivalence et de groupes. *Publ. math., Inst. Hautes tud. Sci.*, (95):93–150, 2002.
- [Jol05] Paul Jolissaint. On property (T) for pairs of topological groups. *Enseign. Math. (2)*, 51(1-2):31–45, 2005.
- [JP07] Andreas Thom Jesse Peterson. Group cocycles and the ring of affiliated operators. *arXiv:0708.4327v1 [math.GR]*, 2007.
- [Mar03] Florian Martin. *Analyse harmonique et 1-cohomologie réduite des groupes localement compacts*. PhD thesis, Univ. Neuchatel, 2003.
- [Val05] Alain Valette. Group pairs with property (T), from arithmetic lattices. *Geom. Dedicata*, 112:183–196, 2005.

Hebrew University, Jerusalem Israel  
and Mathematics & Statistics  
The University of North Carolina at Greensboro  
116 Petty Building  
Greensboro, NC 27402-6170  
email:fernos@math.huji.ac.il