

## THE RANGE OF A CONTRACTIVE PROJECTION ON AN $L_p$ -SPACE

S. J. BERNAU AND H. ELTON LACEY

Suppose  $(X, \Sigma, \mu)$  is a measure space,  $1 \leq p < \infty$  and  $p \neq 2$ . Let  $L_p = L_p(X, \Sigma, \mu)$  be the usual space of equivalence classes of  $\Sigma$ -measurable functions  $f$  such that  $|f|^p$  is integrable. A contractive projection on  $L_p$  is a linear operator  $P: L_p \rightarrow L_p$  such that  $P^2 = P$  and  $\|P\| \leq 1$ . In this paper we give a complete description of such contractive projections in terms of conditional expectation operators. We also show that a closed subspace  $M$  of  $L_p$  is the range of a contractive projection if and only if  $M$  is isometrically isomorphic to another  $L_p$ -space. Another sufficient condition shows, in particular, that every closed vector sublattice of an  $L_p$ -space is the range of a positive contractive projection.

Most of our results are known. The case of finite  $\mu$  was treated, for  $p = 1$ , by Douglas [2] and for  $1 < p < \infty$  by Ando [1] who showed how to reduce this case to that of  $p = 1$ . These authors obtained our necessary and sufficient condition. Grothendieck [4] considered  $p = 1$  and general  $\mu$  and showed that the range of a contractive projection on  $L_1$  is isometrically isomorphic to another  $L_1$ -space. Wulbert [11] showed that a positive contractive projection on  $L_1$  which is also  $L_\infty$  contractive is a conditional expectation, and pointed out that his proofs applied for  $p > 1$ . Tzafriri [10] showed that for general  $\mu$  the range of a contractive projection on  $L_p$  is isometrically isomorphic to another  $L_p$ -space. In [5] we gave an outline, based on Tzafriri's, of another proof of this fact.

We obtain complete generalizations of the Douglas-Ando results to the case of an arbitrary measure  $\mu$ . We have chosen to give our proofs in detail. It seems easier not to reduce the case  $p > 1$  to the case  $p = 1$ . The proofs for  $p > 1$  often use duality arguments which are just not available for  $p = 1$ . By giving such proofs, generalizations to reflexive Banach function spaces may be possible. Some such generalizations have been tried by Rao [8] but his reduction from arbitrary norms to the  $L_1$  case is faulty and his Theorem 2.7 is false in general (see Remark 4.4). Duplissey [3] considers Banach function spaces but requires  $\|Pf\|_\infty \leq \|f\|_\infty$  as well as  $P$  contractive. We also avoid reducing to the case of finite measures. This device turns out to be unnecessary, and needlessly complicated.

We have deliberately omitted the cases  $0 < p < 1$ , except in the appendix, and the case  $p = 2$ . A contractive projection on Hilbert

space is an orthogonal projection and every closed subspace is the range of a unique one. For  $0 < p < 1$  the arguments for  $p = 1$  will work or can be modified to work. We no longer have a norm, however, and it seemed best to ignore this case.

We have included a section in which we discuss the proof of the famous theorem that if  $1 \leq p < \infty$ , a Banach space is an  $L_p$ -space, if and only if it is an  $\mathcal{L}_{p,\lambda}$  for all  $\lambda > 1$ , if and only if it contains an increasing set of finite dimensional subspaces whose union is dense and each of which is isometrically isomorphic to a finite dimensional  $l_p$ -space of appropriate dimension. This result is a combination of work of Zippin [12] and of Lindenstrauss and Pełczyński [7]. We discussed the real case in [5]. There seems to some value in going over the results again here because both [5] and [7] really consider only the real case. The extensions to the complex case are technically more difficult than is admitted in [7]. Also we have had many questions about some of the details omitted in [5].

In our final appendix we have given two technical results used by Ando [1] and Tzafriri [10]. Our proofs seem a little easier and Ando's result has been generalized to arbitrary measure spaces.

1. Notation and definitions. We consider complex  $L_p$ -spaces throughout. Our proofs are valid, with obvious modifications in the real case too. We use, for complex  $z$ , the version of the signum function,  $\operatorname{sgn} z$  defined by

$$\operatorname{sgn} z = \begin{cases} z/|z| & \text{if } z \neq 0 \\ 0 & \text{if } z = 0. \end{cases}$$

We modify some standard vector lattice terminology to apply in the complex case. A *closed vector sublattice* of  $L_p$  is a closed subspace  $M$  such that if  $f \in M$ ,  $\operatorname{Re} f \in M$ , and if  $f \in M$  and  $f$  is real-valued,  $f^+ = f \vee 0 \in M$ .

If  $f \in L_p$  write  $S(f) = \{x \in X: f(x) \neq 0\}$  and call  $S(f)$  the *support* of  $f$ . This only determines the support of  $f$  to a set of  $\mu$ -measure zero. However, this will either not matter, or we will want all possible determinations for the support of  $f$ . If  $M \subset L_p$ , the *polar* of  $M$ ,  $M^\perp$ , is defined by

$$M^\perp = \{g \in L_p: |g| \wedge |m| = 0 (m \in M)\}.$$

(By  $|g| \wedge |m| = 0$  we mean  $\mu$ -almost everywhere of course.) If  $M = M^{\perp\perp}$  we call  $M$  a *band* (or *polar subspace*). If  $M$  is a band  $L_p = M \oplus M^\perp$ , and the natural, *band projection*  $J_M$  of  $L_p$  onto  $M$  is given, for positive  $h \in L_p$ , by

$$J_M h = \sup \{g \in M: 0 \leq g \leq h\} .$$

If  $f \in L_p$ , and  $M = f^{\perp\perp}$ , we write  $J_f$  for the band projection on  $f^{\perp\perp}$  and note that, if  $0 \leq h \in L_p$

$$J_f h = \sup \{h \wedge n|f|: n = 1, 2, \dots\} ,$$

(indeed, by dominated convergence,  $h \wedge n|f| \rightarrow J_f h$  in  $L_p$ -norm) while for any  $h \in L_p$ ,  $J_f h = \chi_{S(f)} h$ . The following lemma is easy to prove.

LEMMA 1.1. *If  $M$  is a subspace of  $L_p(X, \Sigma, \mu)$ ,  $h \in L_p$ , and  $J$  is the band projection on  $M^{\perp\perp}$ , then there is a sequence  $(f_n)$  in  $M$  such that  $Jh = \lim \chi_{S(f_n)} h$ .*

*Proof.* Choose a sequence  $(f_n)$  in  $M$  such that

$$\|\chi_{S(f_n)} h\|_p \longrightarrow \sup \{\|\chi_{S(f)} h\|_p: f \in M\} .$$

We omit the remaining details.

REMARK 1.2. This lemma can be strengthened, in case  $M$  is closed, to say that for each  $h \in L_p$  there exists  $f \in M$  such that  $Jh = J_f h = \chi_{S(f)} h$ . This depends essentially on the fact that the set of supports of functions whose equivalence classes are in  $M$  is closed under countable union. This is proved by Ando [1, Lemma 3] for finite  $\mu$ , and we give a rather easier alternative proof in our appendix.

2. Preliminary results. In this section the cases  $p = 1$ , and  $1 < p < \infty$ ,  $p \neq 2$ , are treated separately. Our first lemma is based on an argument of Douglas [2, p. 452].

LEMMA 2.1. *Let  $P$  be a contractive projection on  $L_1(X, \Sigma, \mu)$  and suppose  $f \in \mathcal{R}(P)$ ; then*

- (i)  $PJ_f = J_f P J_f$ ;
- (ii)  $P(h \operatorname{sgn} f) = |P(h \operatorname{sgn} f)| \operatorname{sgn} f$  ( $0 \leq h \in L_1$ );
- (iii)  $\|P(h \operatorname{sgn} f)\| = \|J_f h\|$  ( $0 \leq h \in L_1$ ).

*Proof.* Suppose  $0 \leq h \leq |f|$ , then

$$\begin{aligned} \|f\| - \|h \operatorname{sgn} f\| &= \|f - h \operatorname{sgn} f\| \\ &\geq \|P(f - h \operatorname{sgn} f)\| \\ &= \|f - P(h \operatorname{sgn} f)\| \\ &\geq \|f\| - \|P(h \operatorname{sgn} f)\| \\ &\geq \|f\| - \|h \operatorname{sgn} f\| . \end{aligned}$$

This gives equality throughout so (iii) is valid for  $0 \leq h \leq |f|$ . In

addition we have  $0 \leq |f - P(h \operatorname{sgn} f)| = |f| - |P(h \operatorname{sgn} f)|$   $\mu$ -almost everywhere, and (ii) also follows for  $0 \leq h \leq |f|$ . We extend immediately to  $h \in L_1$  such that  $0 \leq h \leq n|f|$  for some  $n$ , and since linear combinations of such  $h$  are dense in  $f^{\perp\perp}$  we have (ii) and (iii) for  $0 \leq h \in f^{\perp\perp}$ . If  $h \in L_1$  and  $h \geq 0$ ,  $(J_f h) \operatorname{sgn} f = h \operatorname{sgn} f$  so (ii) and (iii) are proved.

For (i) take  $g \in L_1$  and put  $h = (\operatorname{Re}(g \operatorname{sgn} \bar{f}))^+ \operatorname{sgn} f$ , by (ii)  $Ph \in f^{\perp\perp}$  so  $Ph = J_f Ph$ . We conclude easily that

$$P(J_f g) = P((g \operatorname{sgn} \bar{f}) \operatorname{sgn} f) = J_f P J_f g$$

and (i) is proved.

Suppose  $1 < p < \infty$ ; then identify the dual of  $L_p(X, \Sigma, \mu)$  with  $L_q(X, \Sigma, \mu)$  in the usual way ( $1/p + 1/q = 1$ ). Let  $P$  be a contractive projection on  $L_p$ . The conjugate operator  $P^*$  is defined uniquely on  $L_q$  by the equation

$$\int P f \cdot g d\mu = \int f \cdot P^* g d\mu \quad (f \in L_p, g \in L_q).$$

Clearly  $P^*$  is a contractive projection on  $L_q$ .

**LEMMA 2.2.** [1, Lemma 1]. *Suppose  $1 < p < \infty$  and let  $P$  be a contractive projection on  $L_p(X, \Sigma, \mu)$ , then  $f \in \mathcal{R}(P)$  if and only if  $|f|^{p-1} \operatorname{sgn} \bar{f} \in \mathcal{R}(P^*)$ .*

*Proof.* Suppose  $f \in \mathcal{R}(P)$ ; by Hölder's inequality

$$\begin{aligned} \|f\|_p^p &= \int |f|^p d\mu = \int P f \cdot |f|^{p-1} \operatorname{sgn} \bar{f} d\mu \\ &= \int f \cdot P^*(|f|^{p-1} \operatorname{sgn} \bar{f}) d\mu \\ &\leq \|f\|_p \|P^*(|f|^{p-1} \operatorname{sgn} \bar{f})\|_q \\ &\leq \|f\|_p \| |f|^{p-1} \operatorname{sgn} \bar{f} \|_q \\ &= \|f\|_p \|f\|_p^{p/q} \\ &= \|f\|_p^p. \end{aligned}$$

The conditions for equality in Hölder's inequality lead to

$$P^*(|f|^{p-1} \operatorname{sgn} \bar{f}) = |f|^{p-1} \operatorname{sgn} \bar{f}$$

as required. This proves necessity. Sufficiency follows dually.

We next generalize an argument in Ando's Theorem 1 [1].

**LEMMA 2.3.** *Suppose  $1 < p < \infty$ ,  $p \neq 2$ ; and let  $P$  be a contractive projection on  $L_p(X, \Sigma, \mu)$ ; if  $f \in \mathcal{R}(P)$  then,*

- (i)  $|f| \operatorname{sgn} g \in \mathcal{R}(P)$  ( $g \in \mathcal{R}(P)$ ),
- (ii)  $P J_f = J_f P$ ,

$$(iii) \quad P(h \operatorname{sgn} f) = |P(h \operatorname{sgn} f)| \operatorname{sgn} f \quad (0 \leq h \in L_p).$$

*Proof.* (i) Suppose first that  $p > 2$ , let  $\lambda \in R$ ,  $0 < |\lambda| < 1$ , and let  $g \in \mathcal{R}(P)$ . By Lemma 2.2,

$$g_\lambda = \lambda^{-1}(|f + \lambda g|^{p-1} \operatorname{sgn} \overline{(f + \lambda g)} - |f|^{p-1} \operatorname{sgn} \bar{f}) \in \mathcal{R}(P^*).$$

Since  $p > 2$ ,

$$\begin{aligned} g_\lambda &= \lambda^{-1}[ (|f + \lambda g|^{p-2} - |f|^{p-2}) \overline{(f + \lambda g)} + |f|^{p-2} \cdot \lambda \bar{g} ] \\ &= \lambda^{-1}[ (|f + \lambda g|^{p-2} - |f|^{p-2}) \overline{(f + \lambda g)} ] + |f|^{p-2} \bar{g}. \end{aligned}$$

Recall, that for real  $\lambda$  and complex  $w, z$ ,  $d/d\lambda |w + \lambda z| = \operatorname{Re}[z \operatorname{sgn} \overline{(w + \lambda z)}]$ , provided  $w + \lambda z \neq 0$ . It follows that as  $\lambda \rightarrow 0$ ,

$$g_\lambda \longrightarrow (p-2) |f|^{p-3} \operatorname{Re}(g \operatorname{sgn} \bar{f}) \cdot \bar{f} + |f|^{p-2} \bar{g}$$

at all points of  $X$  where  $f \neq 0$ .

If  $2|\lambda g| < |f|$  we have  $|f|/2 < |f + \theta \lambda g| < 2|f|$  if  $0 < \theta < 1$ ; and, by the mean value theorem there exists  $\theta$ ,  $0 < \theta < 1$  such that

$$\begin{aligned} |g_\lambda| &\leq (p-2) |f + \theta \lambda g|^{p-3} |\operatorname{Re}(g \operatorname{sgn} \overline{(f + \theta \lambda g)})| |f + \lambda g| + |f|^{p-2} |g| \\ &\leq (p-2) 2^{p-3} |f|^{p-3} |g| 2 |f| + |f|^{p-2} |g| \\ &\leq ((p-2)2^p + 1) |f|^{p-2} |g| \in L_q. \end{aligned}$$

If  $2|\lambda g| \geq |f|$ ,  $|f + \lambda g| \leq 3|\lambda g|$  and

$$\begin{aligned} |g_\lambda| &\leq \lambda^{-1} [(3|\lambda g|)^{p-1} + (2|\lambda g|)^{p-1}] \\ &= (3^{p-1} + 2^{p-1}) |g|^{p-1} |\lambda|^{p-2} \\ &\leq (3^{p-1} + 2^{p-1}) |g|^{p-1} \in L_q. \end{aligned}$$

The penultimate line above shows that  $g_\lambda \rightarrow 0$  ( $\lambda \rightarrow 0$ ) if  $f = 0$ .

This shows that  $g_\lambda$  converges to

$$g_0 = (p-2) |f|^{p-2} \operatorname{sgn} \bar{f} \operatorname{Re}(g \operatorname{sgn} \bar{f}) + |f|^{p-2} \bar{g},$$

pointwise almost everywhere on  $X$  and that the convergence is dominated by an element of  $L_q$ . Hence  $\|g_\lambda - g_0\|_q \rightarrow 0$  and  $g_0 \in \mathcal{R}(P^*)$  because  $\mathcal{R}(P^*)$  is closed.

By the same argument, applied to  $-ig$ , we have, using  $\operatorname{Re} - iz = \operatorname{Im} z$ ,

$$k_0 = (p-2) |f|^{p-2} \operatorname{sgn} \bar{f} \operatorname{Im}(g \operatorname{sgn} \bar{f}) + i |f|^{p-2} \bar{g} \in \mathcal{R}(P^*).$$

Now,

$$\begin{aligned} g_0 - ik_0 &= (p-2) |f|^{p-2} \operatorname{sgn} \bar{f} \cdot \overline{(g \operatorname{sgn} \bar{f})} + 2 |f|^{p-2} \bar{g} \\ &= (p-2) |f|^{p-2} \operatorname{sgn} \bar{f} \cdot \bar{g} \cdot \operatorname{sgn} f + 2 |f|^{p-2} \bar{g} \\ &= p |f|^{p-2} \cdot \bar{g} \in \mathcal{R}(P^*). \end{aligned}$$

(Note that this last is valid in the real case too.)

Using Lemma 2.2 again, we conclude that  $\| |f|^{p-2} \bar{g} |^{q-1} \operatorname{sgn} \overline{|f|^{p-2} \bar{g}} = |f|^{1-(q-1)} |g|^{q-1} \operatorname{sgn} g \in \mathcal{R}(P)$ . Set

$$k_n = |f|^{1-(q-1)^n} |g|^{(q-1)^n} \operatorname{sgn} g \quad (n = 1, 2, \dots).$$

We have just shown that  $k_1 \in \mathcal{R}(P)$  and the same method, applied inductively, gives  $k_n \in \mathcal{R}(P)$  for all  $n$ . Since  $0 < q - 1 < 1$ ,

$$|k_n| \leq \max\{|f|, |g|\} \leq |f| + |g| \in L_p,$$

so  $(k_n)$  is dominated in  $L_p$ . Since  $k_n \rightarrow |f| \operatorname{sgn} g$   $\mu$ -almost everywhere on  $X$ , we have  $\|k_n - |f| \operatorname{sgn} g\|_p \rightarrow 0$  and since  $\mathcal{R}(P)$  is closed  $|f| \operatorname{sgn} g \in \mathcal{R}(P)$  which proves (i) for  $p > 2$ .

Suppose  $1 < p < 2$ ; as we have already stated  $P^*$  is a contractive projection on  $L_q$ , and  $q > 2$ . By Lemma 2.2,  $f_1 = |f|^{p-1} \operatorname{sgn} \bar{f}$  and  $g_1 = |g|^{p-1} \operatorname{sgn} \bar{g}$  are in  $\mathcal{R}(P^*)$ . By our proof above  $|f_1| \operatorname{sgn} g_1 = |f|^{p-1} \operatorname{sgn} \bar{g} \in \mathcal{R}(P^*)$ , and, by Lemma 2.2 again,  $|f| \operatorname{sgn} g \in \mathcal{R}(P)$ .

This completes the proof of (i).

For (ii) we have by (i), that  $|f| \operatorname{sgn} Pk \in \mathcal{R}(P)$  ( $k \in L_p$ ). By (i) again,

$$J_f Pk = |Pk| \operatorname{sgn} (|f| \operatorname{sgn} Pk) \in \mathcal{R}(P).$$

Thus  $J_f P = P J_f P$ . Further, since  $P^*$  is a contractive projection on  $L_q$ , and  $|f|^{p-1} \operatorname{sgn} \bar{f} \in \mathcal{R}(P^*)$  we have  $J_g P^* = P^* J_g P^*$  with

$$g = |f|^{p-1} \operatorname{sgn} \bar{f}.$$

In addition  $J_g = J_g^*$ , since  $J_g$  and  $J_f$  are each multiplication by the same characteristic function. We conclude

$$J_f P = P J_f P = (P^* J_f^* P^*)^* = (P^* J_g P^*)^* = (J_g P^*)^* = P J_f,$$

which is (ii).

(iii) The proof is like the proof of Lemma 2.1(ii). Suppose  $0 \leq h \leq |f|$ . By (i),  $|f| \operatorname{sgn} P(h \operatorname{sgn} f) \in \mathcal{R}(P)$ , so by Lemma 2.2,

$$|f|^{p-1} \operatorname{sgn} \overline{P(h \operatorname{sgn} f)} \in \mathcal{R}(P^*).$$

Hence,

$$\begin{aligned} \int |P(h \operatorname{sgn} f)| |f|^{p-1} d\mu &= \int P(h \operatorname{sgn} f) \cdot |f|^{p-1} \operatorname{sgn} \overline{P(h \operatorname{sgn} f)} d\mu \\ &= \int h \operatorname{sgn} f \cdot |f|^{p-1} \operatorname{sgn} \overline{P(h \operatorname{sgn} f)} d\mu \\ &\leq \int h |f|^{p-1} d\mu. \end{aligned}$$

Also  $0 \leq |f - h \operatorname{sgn} f| = |f| - h \leq |f|$ .

Hence,

$$\begin{aligned}
\|f\|_p^p &= \int |P(|f| \operatorname{sgn} f)| |f|^{p-1} d\mu \\
&= \int |P(h \operatorname{sgn} f) + P(|f| - h) \operatorname{sgn} f| |f|^{p-1} d\mu \\
&\leq \int |P(h \operatorname{sgn} f)| |f|^{p-1} d\mu + \int |P(|f| - h) \operatorname{sgn} f| |f|^{p-1} d\mu \\
&\leq \int h |f|^{p-1} d\mu + \int (|f| - h) |f|^{p-1} d\mu \\
&= \|f\|_p^p.
\end{aligned}$$

We have equality at each stage and hence, ( $\mu$ -almost everywhere),

$$|f| = |P(|f| \operatorname{sgn} f)| = |P(h \operatorname{sgn} f)| + |f - P(h \operatorname{sgn} f)|.$$

This proves (iii) for  $0 \leq h \leq |f|$ . The extension to  $0 \leq h \in L_p$  is the same as in the proof of Lemma 2.1(ii) and (iii) so we are done.

**3. Contractive projections and conditional expectations.** In this section we describe the contractive projections on  $L_p(X, \Sigma, \mu)$  ( $1 \leq p < \infty, p \neq 2$ ) in terms of conditional expectation.

We first need the necessary  $\sigma$ -subring.

**LEMMA 3.1.** *Suppose  $1 \leq p < \infty, p \neq 2$ , and let  $P$  be a contractive projection on  $L_p(X, \Sigma, \mu)$ . Define  $\Sigma_0$  to be the set of supports of all functions whose equivalence classes are in  $\mathcal{R}(P)$ ; then*

- (i)  $PJ_g f = J_g f$  ( $f, g \in \mathcal{R}(P)$ );
- (ii)  $\Sigma_0$  is a  $\sigma$ -subring of  $\Sigma$ .

*Proof.* (i) By Lemma 2.3(ii), (i) is valid if  $p \neq 1$ . We give a proof that uses only the identity  $J_g P J_g = P J_g$  valid for  $1 \leq p < \infty, p \neq 2$  (Lemma 2.1(i) or 2.3(ii) weakened). Since  $f - J_g f \in g^\perp$  and  $J_g f - P J_g f \in g^{\perp\perp}$ , we have

$$\begin{aligned}
\|P(f - J_g f)\|^p &= \|f - P J_g f\|^p \\
&= \|f - J_g f\|^p + \|J_g f - P J_g f\|^p \\
&\geq \|P(f - J_g f)\|^p + \|J_g f - P J_g f\|^p.
\end{aligned}$$

Thus  $P J_g f = J_g f$  which is (i).

(ii) By (i),  $S(f) \sim S(g) = S(f - J_g f) = S(P(f - J_g f)) \in \Sigma_0$ . Thus  $\Sigma_0$  is closed under differences. If  $(f_n)$  is a sequence of nonzero elements in  $\mathcal{R}(P)$  such that  $S(f_n) \cap S(f_m) = \emptyset$  ( $m \neq n$ ) then

$$f = \Sigma 2^{-n} \|f_n\|^{-1} f_n \in \mathcal{R}(P)$$

and  $S(f) = \bigcup S(f_n)$ . This proves (ii).

**COROLLARY 3.2.** *Let  $P$  be a contractive projection on  $L_p(X, \Sigma, \mu)$  ( $1 \leq p < \infty, p \neq 2$ ). If  $h \in \mathcal{R}(P)^{\perp\perp}$  there exists  $f \in \mathcal{R}(P)$  such that  $h \in f^{\perp\perp}$ .*

*Proof.* By Lemma 1.1 there is a sequence  $(f_n)$  in  $\mathcal{R}(P)$  such that  $h = \lim_{n \rightarrow \infty} \chi_{S(f_n)} h$ . Choose  $f \in \mathcal{R}(P)$  such that  $S(f) = \bigcup S(f_n)$ , then  $h \in f^{\perp\perp}$ .

Observe now that if  $f \in L_p$  the measure  $|f|^p \mu$  restricted to any  $\sigma$ -subring,  $\Sigma_0$ , of  $\Sigma$ , is finite. By the Radon-Nikodym theorem we may define the *conditional expectation operator*,  $\mathcal{E}_f = \mathcal{E}(\Sigma_0, |f|^p)$ , for the measure  $|f|^p \mu$  relative to  $\Sigma_0$ .  $\mathcal{E}_f$  is uniquely determined by the equation

$$\int_A h |f|^p d\mu = \int_A (\mathcal{E}_f h) |f|^p d\mu \quad (A \in \Sigma_0)$$

for  $h \in L_1(X, \Sigma, |f|^p d\mu)$ , and the condition that  $\mathcal{E}_f h$  is  $\Sigma_0$ -measurable.

**LEMMA 3.3.** *Suppose  $1 \leq p < \infty, p \neq 2$ ; let  $P$  be a contractive projection on  $L_p(X, \Sigma, \mu)$  and let  $\Sigma_0$  be the  $\sigma$ -subring of  $\Sigma$ , consisting of supports of functions in  $\mathcal{R}(P)$ . If  $M_f = f^{-1} J_f \mathcal{R}(P) = \{f^{-1} J_f g : g \in \mathcal{R}(P)\}$  then  $M_f = L_p(S(f), \Sigma_0 | S(f), |f|^p \mu)$  where  $\Sigma_0 | S(f) = \{A \in \Sigma_0 : A \subset S(f)\}$  and we make the obvious identification of functions on  $S(f)$  and functions on  $X$  which vanish off  $S(f)$ . In addition the map  $h \rightarrow f^{-1} h$  is an isometric isomorphism between  $J_f \mathcal{R}(P)$  and  $L_p(S(f), \Sigma_0 | S(f), |f|^p \mu)$ .*

*Proof.* Observe that  $|f|^p \mu$  is finite on  $S(f)$ , and that the isometry claim is obviously true. If  $A \in \Sigma_0 | S(f)$  then  $A = S(g)$  for some  $g \in \mathcal{R}(P)$ . By Lemmas 2.1 and 3.1 (if  $p = 1$ ) or 2.3 (if  $p > 1$ ) we have  $J_g f = P J_g f$  so that  $\chi_A = f^{-1} J_g f \in M_f$ . Let  $h$  be a simple function with respect to  $\Sigma_0 | S(f)$ . Then  $h \in M_f$  and  $h f \in \mathcal{R}(P)$ . In addition

$$\int_{S(f)} |h|^p \cdot |f|^p d\mu = \int_X |h f|^p d\mu.$$

We conclude that

$$M_f \supset L_p(S(f), \Sigma_0 | S(f), |f|^p \mu).$$

Conversely, let  $h \in M_f$ , then  $h \in L_p(S(f), \Sigma | S(f), |f|^p \mu)$  and it is enough to show that  $h$  is  $\Sigma_0$ -measurable. Let  $g = (\operatorname{Re} h)^+$ , then  $g f \in L_p(X, \Sigma, \mu)$ . By Lemma 2.1(ii) or 2.3(iii)

$$P(gf) = P(|gf| \operatorname{sgn} f) = |P(|gf| \operatorname{sgn} f)| \operatorname{sgn} f$$

so  $f^{-1} P(gf) = |f|^{-1} |P(|gf| \operatorname{sgn} f)| \in M_f$ . It follows that

$$\operatorname{Re} h = f^{-1}P((\operatorname{Re} h)^+ f) - f^{-1}P((\operatorname{Re} h)^- f) \in M_f .$$

Since each of these functions is nonnegative it is sufficient to consider  $0 \leq h \in M_f$ . Suppose  $\alpha > 0$  and put  $k = h \vee \alpha \chi_{S(f)}$ . Arguing as above, we have  $f^{-1}P(kf) \geq h$  and  $f^{-1}P(kf) \geq \alpha \chi_{S(f)}$  so that  $f^{-1}P(kf) \geq k \geq 0$ . Since  $P$  is contractive we have

$$\begin{aligned} \|kf\|^p &\geq \|P(kf)\|^p = \|P(kf) - kf + kf\|^p \\ &\geq \|P(kf) - kf\|^p + \|kf\|^p . \end{aligned}$$

This gives  $P(kf) = kf$ , so that  $k \in M_f$ . This shows, incidently, that  $M_f$  is a lattice. For our purpose, however, we have

$$\begin{aligned} \{t \in S(f): h(t) > \alpha\} &= \{t \in S(f): (k - \alpha \chi_{S(f)})(t) \neq 0\} \\ &= S(kf - \alpha f) \in \Sigma_0 . \end{aligned}$$

Thus  $M_f$  consists of  $\Sigma_0$ -measurable functions and we are done.

**THEOREM 3.4.** *Suppose  $1 \leq p < \infty$ ,  $p \neq 2$  and that  $P$  is a contractive projection on  $L_p(X, \Sigma, \mu)$ . If  $f \in \mathcal{R}(P)$  and  $h \in f^{\perp}$  then*

$$Ph = f \mathcal{E}(\Sigma_0, |f|^p)(hf^{-1}) .$$

*Proof.* Since  $f^{-1}Ph \in M_f$  we know  $f^{-1}Ph$  is  $\Sigma_0$ -measurable. Thus we have only to show

$$\int_A f^{-1}Ph |f|^p d\mu = \int_A hf^{-1} \cdot |f|^p d\mu \quad (A \in \Sigma_0) .$$

Choose  $g \in \mathcal{R}(P)$  such that  $A = S(g)$ . By Lemma 3.1(i),  $u = J_g f \in \mathcal{R}(P)$ .

Suppose  $p = 1$  and  $0 \leq k \in L_1$ . By Lemma 2.1(ii) and (iii),

$$\begin{aligned} \int_A k \operatorname{sgn} f \cdot f^{-1} |f| d\mu &= \int_{A \cap S(f)} k d\mu = \|J_u k\| = \|P(k \operatorname{sgn} u)\| \\ &= \| |P(J_g k \operatorname{sgn} f)| \operatorname{sgn} f \| \\ &= \int_A f^{-1}P(J_g k \operatorname{sgn} f) \cdot |f| d\mu . \end{aligned}$$

Putting  $v = f - u = f - J_g f \in \mathcal{R}(P)$ , we have, by Lemma 2.1(i),

$$P(k \operatorname{sgn} f) = J_u P(J_u k \operatorname{sgn} f) + J_v P(J_v k \operatorname{sgn} f) .$$

Hence

$$\int_A f^{-1}P(J_g k \operatorname{sgn} f) \cdot |f| d\mu = \int_A f^{-1}P(k \operatorname{sgn} f) \cdot |f| d\mu .$$

We conclude that

$$\int_A h f^{-1} \cdot |f| d\mu = \int_A f^{-1} P h \cdot |f| d\mu$$

for all  $h \in f^{\perp\perp}$  and all  $A \in \Sigma_0$  so we are finished for  $p = 1$ .

If  $p > 1$  we have  $PJ_g = J_g P$  by Lemma 2.3(ii) and  $|f|^{p-1} \operatorname{sgn} \bar{f} \in \mathcal{R}(P^*)$  by Lemma 2.2. Hence,

$$\begin{aligned} \int_A h f^{-1} \cdot |f|^p d\mu &= \int_X J_g h \cdot |f|^{p-1} \operatorname{sgn} \bar{f} d\mu \\ &= \int_X J_g h \cdot P^*(|f|^{p-1} \operatorname{sgn} \bar{f}) d\mu \\ &= \int_X P J_g h \cdot |f|^{p-1} \operatorname{sgn} \bar{f} d\mu \\ &= \int_X J_g P h \cdot f^{-1} |f|^p d\mu \\ &= \int_A f^{-1} P h \cdot |f|^p d\mu \quad (A \in \Sigma_0). \end{aligned}$$

Thus

$$P h = f^{-1} \mathcal{E}(\Sigma_0, |f|^p)(h f^{-1}) \quad (h \in f^{\perp\perp})$$

as claimed.

Our theorem has useful consequences.

**THEOREM 3.5.** *Suppose  $1 \leq p < \infty$ ,  $p \neq 2$ , let  $P$  be a contractive projection on  $L_p(X, \Sigma, \mu)$  and let  $J$  be the band projection on  $\mathcal{R}(P)^{\perp\perp}$ ; then  $PJ$  is the unique contractive projection on  $L_p$  which satisfies  $\mathcal{R}(PJ) = \mathcal{R}(P)$  and  $PJ\mathcal{R}(P)^\perp = \{0\}$ . If  $p \neq 1$ ,  $P = PJ$  so  $P$  is uniquely determined by its range. If  $p = 1$ , and  $A$  is a linear contraction on  $L_1$  which satisfies  $PA = A$  and  $AJ = 0$ , then  $PJ + A$  is a contractive projection on  $L_1$  with the same range as  $P$ .*

*Proof.* Let  $Q$  be a contractive projection on  $L_p$  such that  $\mathcal{R}(Q) = \mathcal{R}(P)$  and  $Q\mathcal{R}(P)^\perp = \{0\}$ . Then  $Q = QJ$  and if  $h \in L_p$  there exists, by Corollary 3.2,  $f \in \mathcal{R}(P) = \mathcal{R}(Q)$  such that  $Jh = J_f h$ . By Theorem 3.4,  $Qh = QJh = f^{-1} \mathcal{E}(\Sigma_0, |f|^p)(Jh \cdot f^{-1}) = PJh$ . Thus  $Q = PJ$ . (It is clear that  $PJ$  satisfies the stated conditions.)

If  $p \neq 1$  take  $h, f$  as above and put  $u = Ph - PJh = Ph - PJ_f h = Ph - J_f Ph$ , by Lemma 2.3(ii). Since band projections commute and  $u \in \mathcal{R}(P) \cap f^\perp$ ,  $J_u h = J_u Jh = J_u J_f h = 0$ . By Lemma 2.3(ii) again,

$$u = J_u u = J_u Ph - J_u PJ_f h = PJ_u h - J_u J_f Ph = 0 - 0 = 0.$$

Hence  $P = PJ$  as required.

If  $p = 1$ ,  $PA = A$ , and  $AJ = 0$ , we have  $AP = AJP = 0$  and  $A^2 =$

$APA = 0$ . Also  $(PJ + A)^2 = PJPJ + PJA + APJ + A^2 = PPJ + PJPA + 0 + 0 = PJ + A$ . Thus  $PJ + A$  is a projection. Observe that

$$\begin{aligned} \mathcal{R}(PJ + A) &= \mathcal{R}(PJ + PA) \subset \mathcal{R}(P) = \mathcal{R}(PJP + AP) \\ &= \mathcal{R}((PJ + A)P) \subset \mathcal{R}(PJ + A). \end{aligned}$$

It remains to show that if  $A$  is contractive,  $PJ + A$  is contractive. If  $h \in L_1$ ,

$$\begin{aligned} \|(PJ + A)h\|_1 &= \|PJh + A(h - Jh)\|_1 \\ &\leq \|PJh\|_1 + \|A(h - Jh)\|_1 \\ &\leq \|Jh\|_1 + \|h - Jh\|_1 \\ &= \|Jh + h - Jh\|_1 \\ &= \|h\|_1. \end{aligned}$$

**4. Contractive projections and isometric isomorphisms.** In this section we prove the equivalence of various conditions on a subspace of  $L_p$  so that it is the range of a contractive projection.

Let  $\mathcal{S}(X, \Sigma)$  denote the set of  $\Sigma$ -measurable functions  $h$  such that  $S(h)$  is  $\sigma$ -finite. By a *multiplication operator* on  $\mathcal{S}(X, \Sigma)$  we mean a map  $h \rightarrow kh$  defined for functions  $h$  in some subset of  $\mathcal{S}(X, \Sigma)$  and some fixed  $\Sigma$ -measurable function  $k$ . If  $k$  satisfies  $|k| = 1$  on  $S(k)$  we will call  $k$  a *unitary multiplication*.

A multiplication operator on  $\mathcal{S}(X, \Sigma)$  preserves equality almost everywhere and hence induces a multiplication operator on each  $L_p(X, \Sigma, \mu)$  into  $\mathcal{S}(X, \Sigma)$  modulo null functions ( $1 \leq p < \infty$ ). Further,  $k_1$  and  $k_2$  will induce the same such multiplication operator on  $L_p$  if  $k_1$  and  $k_2$  agree locally almost everywhere.

Suppose that  $\mathcal{K}$  is a set of  $\Sigma$ -measurable functions such that if  $k_1, k_2 \in \mathcal{K}$  and  $k_1 \neq k_2$ ,  $\mu(S(k_1) \cap S(k_2)) = 0$ . If  $f \in \mathcal{S}(X, \Sigma)$  then, because  $S(f)$  has  $\sigma$ -finite measure,  $S(f)$  meets at most countably many  $S(k)$ , with  $k \in \mathcal{K}$ , in a set of positive measure. Enumerate these as  $(k_n)$ , then there is a unique set  $N \in \Sigma$  such that,  $N \subset S(f)$  and each  $t \in S(f) \sim N$  lies in at most one set  $S(k_n)$ . (In fact  $N = \bigcup_{1 \leq n < m < \infty} (S(k_n) \cap S(k_m))$ .) On  $S(f) \sim N$  the series  $\sum_{n=1}^{\infty} f(t)k_n(t)$  has at most one nonzero term. Thus  $\mathcal{K}$  determines a map  $U_{\mathcal{K}} : \mathcal{S}(X, \Sigma) \rightarrow \mathcal{S}(X, \Sigma)$  by taking, for  $f$  as above,  $U_{\mathcal{K}}f(t) = \sum_{n=1}^{\infty} f(t)k_n(t)$  for  $t \in S(f) \sim N$  and  $U_{\mathcal{K}}f(t) = 0$  elsewhere. We call  $U_{\mathcal{K}}$  the *direct sum* of the (disjoint) multiplication operators induced by the elements of  $\mathcal{K}$ . If  $U_{\mathcal{K}}$  maps  $L_p$  to  $L_p$  ( $1 \leq p < \infty$ ) it is not hard to check that the net of finite sums of the multiplication operators in  $\mathcal{K}$  is strongly convergent to  $U_{\mathcal{K}}$ .

We can now state our theorem. The equivalence of (i) and (ii) generalizes [1, Theorem 4] and extends [10, Theorem 6].

**THEOREM 4.1.** *Suppose  $1 \leq p < \infty$  and  $p \neq 2$  and let  $M$  be a subspace of  $L_p(X, \Sigma, \mu)$ . The following conditions on  $M$  are equivalent.*

- (i)  *$M$  is the range of a contractive projection on  $L_p$ .*
- (ii) *There is a measure space  $(\Omega, \mathcal{E}, \lambda)$  such that  $M$  is isometrically isomorphic to  $L_p(\Omega, \mathcal{E}, \lambda)$ .*
- (iii) *There is a direct sum of unitary multiplication operators  $U: L_p(X, \Sigma, \mu) \rightarrow L_p(X, \Sigma, \mu)$  such that  $U$  is an isometry and  $UM$  is a closed vector sublattice of  $L_p(X, \Sigma, \mu)$ .*

*Furthermore, in (ii) we can always choose  $\Omega = X$ ,  $\mathcal{E}$  a  $\sigma$ -subring of  $\Sigma$ ,  $\lambda$  absolutely continuous with respect to  $\mu$ , and the isometry a direct sum of multiplication operators.*

*If  $\mu$  is  $\sigma$ -finite the direct sums of multiplication operators can be taken to be ordinary multiplications.*

*Proof.* Assume (i). By Zorn's lemma there is a maximal subset  $\mathcal{H}$  of  $M$  consisting of functions  $f \in M$ , such that  $\mu(S(f_1) \cap S(f_2)) = 0$  if  $f_1 \neq f_2$ . If  $g \in M$ ,  $S(g)$  is  $\sigma$ -finite and there is countable subset  $\{f_n\}$  of  $\mathcal{H}$  such that if  $f \in \mathcal{H} \sim \{f_n\}$ ,  $\mu(S(f) \cap S(g)) = 0$ . By Lemma 3.1,  $\Sigma_0$  is a  $\sigma$ -ring so, there exists  $h \in M$  such that  $S(h) = S(g) \sim \bigcup S(f_n)$  and by maximality of  $\mathcal{H}$ ,  $h = 0$ . Define a measure  $\lambda$  on  $\Sigma_0$  by  $\lambda A = \sum_{f \in \mathcal{H}} \int_A |f|^p d\mu$ . This definition is meaningful since  $A$  has  $\sigma$ -finite  $\mu$ -measure and at most countably many of the integrals are nonzero. For  $f \in \mathcal{H}$  define  $f^{-1}$  by

$$f^{-1}(t) = \begin{cases} 1/f(t) & t \in S(f) \\ 0 & t \notin S(f) \end{cases},$$

and let  $V$  be the direct sum of the multiplications  $f^{-1}(f \in \mathcal{H})$ . By Lemma 3.3  $J_f h \rightarrow f^{-1}h (h \in M)$  is an isometric isomorphism of  $J_f M$  with  $L_p(S(f), \Sigma_0|S(f), |f|^p \mu)$ . It is routine to check that  $V$  is an isometric isomorphism of  $M$  with  $L_p(X, \Sigma_0, \lambda)$ . ( $M$  is the direct sum of its subspaces  $J_f M (f \in \mathcal{H})$  and similarly for the  $L_p$ -spaces.)

It  $\mu$  is  $\sigma$ -finite  $\mathcal{H}$  will be countable, say  $\mathcal{H} = \{f_n\}$  and we can find  $f \in M$  such that  $S(f) = \bigcup S(f_n)$ . Then  $\Sigma_0$  consists entirely of subsets of  $S(f)$  and sets of measure zero so that  $M_f = L_p(X, \Sigma_0, |f|^p \mu)$ ,  $J_f M = M$ , and  $V$  can be multiplication by  $f^{-1}$ .

Assume (ii) and let  $T: L_p(\Omega, \mathcal{E}, \lambda) \rightarrow L_p(X, \Sigma, \mu)$  be a linear isometry with range  $M$ . Suppose  $a, b \in L_p(\Omega, \mathcal{E}, \lambda)$  and  $|a| \wedge |b| = 0$ , we claim that  $|Ta| \wedge |Tb| = 0$ . This is essentially proved by Lamperti [6]. Since  $|a| \wedge |b| = 0$ ,  $\|a + b\|^p + \|a - b\|^p = 2\|a\|^p + 2\|b\|^p$ . Since  $T$  is an isometry,  $\|Ta + Tb\|^p + \|Ta - Tb\|^p = 2\|Ta\|^p + 2\|Tb\|^p$ . Since  $p \neq 2$ , the equality condition for Clarkson's inequality [6, Corollary 2.1] shows that  $|Ta| \wedge |Tb| = 0$ .

Take a maximal subset of  $\mathcal{E}$  consisting of sets of nonzero finite

$\lambda$ -measure which intersect pairwise in sets of  $\lambda$ -measure zero and let  $\mathcal{K}$  be the corresponding set of characteristic functions. Let  $a \in \mathcal{K}$  and suppose  $B \in \mathcal{E}$  and  $B \subset S(a)$ . Write  $b = \chi_B$ , then  $T(a - b)$ ,  $Tb$  are disjoint in  $M$  so we have  $Tb = |Tb| \operatorname{sgn} Ta$ . This extends to non-negative simple functions  $b$  in  $a^{\perp\perp}$  and then to all nonnegative  $b \in a^{\perp\perp}$ . Define  $U: L_p(X, \Sigma, \mu) \rightarrow L_p(X, \Sigma, \mu)$  to be the direct sum of the unitary multiplications  $\operatorname{sgn} \overline{Ta}$  ( $a \in \mathcal{K}$ ). It is easy to see that  $U$  is an isometry of  $M$  such that  $UT$  is positive and hence  $UM = UT L_p(\Omega, \mathcal{E}, \lambda)$  is a closed vector sublattice of  $L_p(X, \Sigma, \mu)$  (compare the proof in Lemma 3.3 where we showed that functions in  $M_f$  were  $\Sigma_0$ -measurable).

Assume (iii) and let  $\Sigma_0$  be the set of supports of functions (whose equivalence classes are) in  $M$ . Then  $\Sigma_0$  is a  $\sigma$ -subring of  $\Sigma$ . (If  $(f_n)$  is a sequence in  $M$ ,  $S(f_n) = S(Uf_n) = S(|Uf_n|)$  so

$$\bigcup S(f_n) = S(U^{-1}\Sigma 2^{-n} \|f_n\|^{-1} |Uf_n|).$$

If  $f, g \in M$ ,  $J_g = J_{Ug}$ ;  $J_g|Uf| = \lim |Uf| \wedge n|Ug| \in UM$  and  $S(f) \sim S(g) = S(U^{-1}(|Uf| - J_g|Uf|))$ . Let  $f, g \in UM$  and suppose  $f$  is real,  $g \geq 0$  and  $f \in g^{\perp\perp}$ , then  $\{t \in X: (f/g)(t) > \alpha\} = S((f - \alpha g)^+) \in \Sigma_0$ . Thus  $f/g$  is  $\Sigma_0$ -measurable. This extends to all  $f \in UM \cap g^{\perp\perp}$  and hence  $J_g f/g$  is  $\Sigma_0$ -measurable if  $f, g \in UM$  and  $g \geq 0$ . This now extends to all  $f, g \in UM$  and, since  $U^{-1}J_g f/U^{-1}g = J_g f/g$ , we have  $f/g$ ,  $\Sigma_0$ -measurable for  $f, g \in M$  and  $f \in g^{\perp\perp}$ . It follows that  $M$  is the set of all elements in  $L_p(X, \Sigma, \mu)$  which can be written in the form  $hf$  with  $h, \Sigma_0$ -measurable and  $f \in M$ . (If  $h = \chi_{S(g)}$  with  $g \in M$ ,  $hf = J_g f = U^{-1}J_{Ug} Uf \in U^{-1}(UM) = M$ .)

Let  $J$  be the band projection on  $M^{\perp\perp}$ , let  $h \in L_p(X, \Sigma, \mu)$ , choose  $f \in M$  such that  $Jh = J_f h$ , (such an  $f$  exists by the arguments used in Corollary 3.2) and define

$$Ph = f \mathcal{E}(\Sigma_0, |f|^p)(hf^{-1}).$$

Then  $Ph \in M$  and this definition is independent of the choice of  $f$  in  $M$  such that  $h \in f^{\perp\perp}$ . To see this suppose  $g \in M$  and  $h \in g^{\perp\perp}$ . Then  $h$  is zero outside  $S(f) \cap S(g) \in \Sigma_0$  and so is  $\mathcal{E}(\Sigma_0, |f|^p)(hf^{-1})$ ,  $\mu$ -almost everywhere. Let  $B = S(f) \cap S(g)$ , then  $f_1 = \chi_B f \in M$  and

$$\int_A hf^{-1}|f|^p d\mu = \int_{A \cap B} hf^{-1}|f|^p d\mu = \int_A hf_1^{-1}|f_1|^p d\mu \quad (A \in \Sigma_0),$$

so that  $f \mathcal{E}(\Sigma_0, |f|^p)(hf^{-1}) = f_1 \mathcal{E}(\Sigma_0, |f_1|^p)(hf_1^{-1})$ . Thus we may assume  $S(f) = S(g)$ . Now

$$g^{-1} f \mathcal{E}(\Sigma_0, |f|^p)(hf^{-1}) \in L_1(X, \Sigma_0, |g|^p \mu),$$

so we have, for  $A \in \Sigma_0$ ,

$$\begin{aligned} & \int_A g^{-1} f \mathcal{E}(\Sigma_0, |f|^p)(hf^{-1}) |g|^p d\mu \\ &= \int_A g^{-1} f |f^{-1} g|^p \mathcal{E}(\Sigma_0, |f|^p)(hf^{-1}) |f|^p d\mu. \end{aligned}$$

Because  $g^{-1}f$  and  $f^{-1}g$  are  $\Sigma_0$ -measurable and the integrals are finite, the second integral is

$$\int_A g^{-1} f |f^{-1} g|^p h f^{-1} |f|^p d\mu = \int_A h g^{-1} |g|^p d\mu.$$

Thus

$$f \mathcal{E}(\Sigma_0, |f|^p)(hf^{-1}) = g \mathcal{E}(\Sigma_0, |g|^p)(hg^{-1})$$

and our definition of  $Ph$  is unambiguous. If  $h_1, h_2 \in L_p$  we can take  $f \in M$  such that  $Jh_1 = J_f h_1$  and  $Jh_2 = J_f h_2$ . Thus  $P$  is linear. Since  $f^{-1}Ph = \mathcal{E}(\Sigma_0, |f|^p)(hf^{-1})$  we see  $P^2 = P$ . Finally, if  $p > 1$ , write  $u = \mathcal{E}(\Sigma_0, |f|^p)(hf^{-1})$ , we have

$$\|Ph\|_p^p = \int |u|^{p-1} \operatorname{sgn} \bar{u} \cdot \mathcal{E}(\Sigma_0, |f|^p)(hf^{-1}) |f|^p d\mu.$$

Since  $u$  is  $\Sigma_0$ -measurable, this is

$$\begin{aligned} \int |u|^{p-1} \operatorname{sgn} \bar{u} \cdot h f^{-1} |f|^p d\mu &= \int |Ph|^{p-1} \operatorname{sgn} \bar{f} \bar{u} \cdot h d\mu \\ &\leq \| |Ph|^{p-1} \|_q \|h\|_p \\ &= \|Ph\|_p^{p/q} \|h\|_p. \end{aligned}$$

(We used Hölder's inequality and  $q$  for the conjugate index to  $p$ .) We conclude that  $\|Ph\|_p \leq \|h\|_p$ .

Since  $Ph = h(h \in M)$  we have shown that  $M$  is the range of the contractive projection  $P$ .

**REMARK 4.2.** The results (iii) implies (i) (with the same proof) and (i) is equivalent to (ii) are valid if  $p = 2$ ; in fact (i) and (ii) are equivalent for any Hilbert space. If we assume the projection  $P$ , is positive as well as contractive the proof in Lemma 3.3 that  $M_f$  is a lattice shows  $\mathcal{R}(P)$  is a sublattice of  $L_2$  and Theorem 4.1 is valid for  $L_2$  with the projection and the isometry both required to be positive and in (iii)  $M$  required to be a closed vector sublattice. We use this remark in our next result.

**COROLLARY 4.3.** *If  $M$  is a closed vector sublattice of  $L_p$  ( $1 \leq p < \infty$ ) then  $M$  is the range of a positive contractive projection.*

*Proof.* Clearly  $M$  satisfies condition (iii) with  $U = I$ . In the definition of  $Ph$  we may always choose a positive  $f \in M$  such that  $h \in f^{\perp 1}$ . Positivity of  $P$  follows from positivity of conditional expectation.

REMARK 4.4. In the introduction we referred to Rao's paper [8] and claimed that its treatment of contractive projections contained errors. In particular, his Theorem II. 2.7 asserts that if  $M$  is the range of a contractive projection  $P$  on a Banach function space  $L^p(\Sigma)$  there is, under suitable conditions, a unitary multiplication  $U$  such that  $UPU^{-1}$  is a positive contractive projection.

The conditions are all satisfied if  $M$  is the subspace of  $l^3(3) = C^3$  spanned by  $(1, 1, 1)$  and  $(1, 2, -3)$ . Rao's theorem now claims the existence of a unitary multiplication, say by  $u = (\lambda_1, \lambda_2, \lambda_3)$ , such that  $uM$  is a vector sublattice of  $C^3$ . This is impossible, as we show. First,  $uM$  contains the elements  $(0, \lambda_2, -4\lambda_3)$ ,  $(\lambda_1, 0, 5\lambda_3)$ , and  $(4\lambda_1, 5\lambda_2, 0)$ . If  $\text{Re } \lambda_2 \bar{\lambda}_3 = 0$  we have  $\lambda_2 \lambda_3^{-1} = \lambda_2 \bar{\lambda}_3 = \pm i$  and  $uM$  contains  $\text{Im}(0, \lambda_2 \bar{\lambda}_3, -4) = (0, \pm 1, 0)$ ; so that  $(0, 1, 0) \in uM$ , and  $uM = C^3$ . If all  $\text{Re } \lambda_i \bar{\lambda}_j \neq 0$  ( $i \neq j$ ), then  $uM$  contains  $\text{Re}(0, 1, -4\lambda_3 \bar{\lambda}_2)$  and  $\text{Re}(1, 0, 5\lambda_3 \bar{\lambda}_1)$ ; hence, taking a multiple of their infimum,  $(0, 0, 1) \in uM$  and again  $uM = C^3$ .

Exactly the same counterexample vitiates the proof of Rao's Theorem II. 2.8 see p. 177 lines -15 to -11.

The error in both cases seems to be the reduction of the general case of  $L^p(\Sigma)$  to the  $L_1$  situation. Vital to this reduction, but invalid, is the assertion that if  $L^p(\Sigma) \subset L^1(\Sigma, G)$  and  $\|\cdot\|_{1,G} \leq \rho(\cdot)$  then a contraction on  $L^p(\Sigma)$  for the  $\rho$ -norm can be extended to the closure of  $L^p(\Sigma)$  in  $L^1(\Sigma, G)$  with the 1,  $G$ -norm and that the extension is contractive for the 1,  $G$ -norm.

5. The theorem of Lindenstrauss, Pelczynski, and Zippin. We begin by recalling some definitions.

If  $E, F$  are isomorphic Banach spaces,  $d(E, F) = \inf \{\|L\| \|L^{-1}\| : L \text{ is a linear isomorphism between } E \text{ and } F\}$ .

A Banach space  $E$  is an  $\mathcal{L}_{p,\lambda}$  space (for  $1 \leq p \leq \infty$  and  $\lambda \geq 1$ ) if for each finite dimensional subspace  $F$  of  $E$  there is a finite dimensional subspace  $G$  of  $E$  such that  $F \subset G$  and  $d(G, l_p(\dim G)) \leq \lambda$ .

We shall say that a Banach space  $E$  is a  $Z_p$ -space (for  $1 \leq p \leq \infty$ ) if there exists a set  $\mathcal{X}$  of finite dimensional subspaces of  $E$  such that:

- (i)  $\mathcal{X}$  is upwards directed by set inclusion;
- (ii)  $\text{cl } \cup \mathcal{X} = E$ ;
- (iii) each  $F \in \mathcal{X}$  is linearly isometric to  $l_p(\dim F)$ .

Our definitions apply, of course, over the real or complex number

fields.

We now state the theorem of Lindenstrauss-Pelczynski-Zippin, [5], [7], [12].

**THEOREM 5.1.** *Let  $E$  be a Banach space and suppose  $1 \leq p < \infty$ , then the following are equivalent.*

(1) *There is a measure  $(X, \Sigma, \mu)$  such that  $E$  is isometrically isomorphic to  $L_p(X, \Sigma, \mu)$ .*

(2)  *$E$  is a  $Z_p$  space.*

(3)  *$E$  is an  $\mathcal{L}_{p,\lambda}$ -space for all  $\lambda > 1$ .*

As outlined in the introduction we discuss some details of the proof for the complex case.

Observe first that (3) is a trivial consequence of (1). Simply identify  $E$  with  $L_p(X, \Sigma, \mu)$  and take for  $\mathcal{X}$  the subspaces spanned by finite sets of ( $p$ th power)-integrable characteristic functions.

*The proof that (3) implies (2).* This result is certainly part of the folklore. It can be obtained quite efficiently as follows.

**LEMMA 5.2.** *Let  $x_1, \dots, x_n$  be  $n$  linearly independent elements of a normed space  $E$  then there exists  $\varepsilon > 0$  such that if  $y_i \in E$ , and  $\|x_i - y_i\| < \varepsilon (i = 1, 2, \dots, n)$  then  $\{y_1, \dots, y_n\}$  is a linearly independent subset of  $E$ .*

*Proof.* (This is standard but our proof may be novel.) Let  $K$  denote the scalar field and  $S$  the unit sphere in  $K^n$ ,  $S = \{\lambda \in K^n: \|\lambda\| = 1\}$ . The map  $g: S \times E^n \rightarrow E$  defined by  $g((\lambda_1, \dots, \lambda_n), (y_1, \dots, y_n)) = \lambda_1 y_1 + \dots + \lambda_n y_n$  is continuous. By linear independence, the compact set  $S \times (x_1, \dots, x_n)$  does not meet the closed set  $g^{-1}(0)$ . Hence there are open neighborhoods  $U_i$  of  $x_i$ ,  $i = 1, \dots, n$ , such that  $(S \times U_1 \times \dots \times U_n) \cap g^{-1}(0) = \emptyset$ . If  $y_i \in U_i (i = 1, \dots, n)$  it follows that  $\{y_1, \dots, y_n\}$  is linearly independent.

**LEMMA 5.3.** *Let  $E$  be a  $Z_p$ -space, then  $E$  is an  $\mathcal{L}_{p,\lambda}$ -space for every  $\lambda > 1$ .*

*Proof.* Let  $F$  be a finite dimensional subspace of  $E$ . Let  $\{x_1, \dots, x_n\}$  be a basis for  $F$ , such that  $\|x_i\| = 1 (i = 1, \dots, n)$ . Let  $x_i^*, \dots, x_n^* \in E^*$  be such that  $x_i^*(x_j) = \delta_{ij}$ , and let  $M = \sum_{i=1}^n \|x_i^*\|$ . Choose  $\varepsilon > 0$  such that  $M\varepsilon < 1$  and  $\|x_i - y_i\| < \varepsilon$  for  $i = 1, \dots, n$  implies that  $\{y_1, \dots, y_n\}$  is linearly independent. By the  $Z_p$ -hypothesis there is a finite dimensional subspace  $H$  of  $E$  and points  $y_1, \dots, y_n$  in  $H$ , such that  $H$  is isometrically isomorphic to  $l_p(\dim H)$ , and  $\|x_i - y_i\| < \varepsilon (i = 1, \dots, n)$ . Then  $\{y_1, \dots, y_n\}$  is a linearly independent subset of

H. If

$$\sum_{i=1}^n \alpha_i y_i \in \bigcap_{i=1}^n \mathcal{N}(x_i^*),$$

then

$$\begin{aligned} \sum_{j=1}^n |\alpha_j| &= \sum_{j=1}^n \left| x_j^* \left( \sum_{i=1}^n \alpha_i x_i \right) \right| \\ &= \sum_{j=1}^n \left| x_j^* \left( \sum_{i=1}^n \alpha_i (x_i - y_i) \right) \right| \\ &\leq \sum_{j=1}^n \|x_j^*\| \left( \sum_{i=1}^n |\alpha_i| \varepsilon \right) \\ &= M\varepsilon \sum_{i=1}^n |\alpha_i|. \end{aligned}$$

Since  $M\varepsilon < 1$  we conclude that  $\alpha_i = 0$  for each  $i$ . Thus we can extend  $y_1, \dots, y_n$  to a basis, say  $y_1, \dots, y_n, x_{n+1}, \dots, x_p$ , of  $H$  with the property that  $\{x_{n+1}, \dots, x_p\} \subset \bigcap_{i=1}^n \mathcal{N}(x_i^*)$ .

Let  $G$  be the subspace of  $E$  spanned by  $x_1, \dots, x_n, x_{n+1}, \dots, x_p$ . Then  $F \subset G$ . If  $y = \sum_{i=1}^n \alpha_i y_i + \sum_{i=n+1}^p \alpha_i x_i \in H$  define  $Ty = \sum_{i=1}^n \alpha_i x_i + \sum_{i=n+1}^p \alpha_i x_i \in G$ . We have

$$\begin{aligned} \|y - Ty\| &= \left\| \sum_{i=1}^n \alpha_i (y_i - x_i) \right\| \leq \varepsilon \sum_{i=1}^n |\alpha_i| \\ &= \varepsilon \sum_{j=1}^n |x_j^*(Ty)| \\ &\leq M\varepsilon \|Ty\|. \end{aligned}$$

This gives  $(1 - M\varepsilon)\|Ty\| \leq \|y\| \leq (1 + M\varepsilon)\|Ty\| (y \in H)$ ; so that  $T$  is an isomorphism between  $F$  and  $H$  such that  $\|T\| \|T^{-1}\| \leq (1 + M\varepsilon)/(1 - M\varepsilon)$ . If  $\lambda > 1$  we can choose  $\varepsilon$  such that  $(1 + M\varepsilon)/(1 - M\varepsilon) < \lambda$ . Thus  $E$  is an  $\mathcal{L}_{p,\lambda}$ -space for all  $\lambda > 1$ .

*The proof that (2) implies (1).* Here the plan is first to embed  $E$ , isometrically, in an  $L_p$ -space, and then to use the theory of contractive projections of  $L_p$ -spaces.

This is carried out in detail for the real separable case in [7] and for the real nonseparable case in [5]. The generalizations to cover the complex case are mostly obvious. For  $1 < p < \infty$  our Theorem 4.1 is used. For  $p = 1$ , it follows as in the real case that  $E^*$  is a  $\mathcal{S}_1$  space whence by the complex version of Grothendieck's theorem [9]  $E$  is an  $L_1(\mu)$  space.

There is an aspect of the construction which needs a little elaboration. At one stage of the proof we have a complex vector space, say  $V$ , consisting of complex valued functions on a set  $U$ .  $V$  is a vector sublattice of the space of all complex functions on  $U$ . There

is a seminorm  $\pi$  on  $V$  such that  $\pi(f) \leq \pi(g)$  whenever  $|f| \leq |g|$ , and  $\pi(f + g)^p = \pi(f)^p + \pi(g)^p$  whenever  $|f| \wedge |g| = 0$ . We then need to embed the quotient  $V/N$ , where  $N = \{f \in V: \pi(f) = 0\}$ , isometrically in a concrete, complex,  $L_p$ -space. For this, let  $V_R$  and  $N_R$  denote the spaces of real-valued functions in  $V$  and  $N$  respectively. The quotient  $V_R/N_R$  with the norm induced by  $\pi$  is then linearly and lattice isomorphic, and isometric, to a vector sublattice of real  $L_p(X, \Sigma, \mu)$  just as in [7]. Let  $h_1$  denote the composition of the quotient map  $U_R \rightarrow V_R/N_R$  and the isometric isomorphism into real  $L_p(X, \Sigma, \mu)$ . Then  $h_1$  is a linear and lattice homomorphism and  $\|h_1 f\| = \pi(f)$  ( $f \in V_R$ ). We construct the required embedding of  $V/N$  into complex  $L_p(X, \Sigma, \mu)$  by defining

$$h(f + N) = h_1(\operatorname{Re} f) + ih_1(\operatorname{Im} f).$$

Then  $h$  is clearly well defined. To verify that  $h$  is an isometry we need the next lemma.

LEMMA 5.4. *The map  $h$  constructed above satisfies  $h|f| = |hf|$ , ( $f \in V$ ).*

*Proof.* For any real  $\theta$   $|f| \geq \operatorname{Re}(e^{i\theta} f)$  so

$$h|f| = h_1|f| \geq h_1(\operatorname{Re} e^{i\theta} f) = \operatorname{Re} h(e^{i\theta} f) = \operatorname{Re} e^{i\theta} hf.$$

Hence  $h|f| \geq |hf|$ . For the converse, let  $\omega$  be a complex  $n$ th root of unity and observe that for any complex  $z$

$$\max \{\operatorname{Re} \omega^r z: r = 1, 2, \dots, n\} \geq \cos(\pi/n)|z|.$$

Hence,

$$\begin{aligned} \cos(\pi/n)h|f| &\leq h(\sup \{(\operatorname{Re} \omega^r f): r = 1, \dots, n\}) \\ &= \sup \{\operatorname{Re} \omega^r hf: r = 1, \dots, n\} \\ &\leq |hf|. \end{aligned}$$

Letting  $n \rightarrow \infty$  we have  $h|f| = |hf|$  as required.

This completes our discussion of the proof of Theorem 5.1. We add a comment. It seems that a more elementary proof that a space which is an  $\mathcal{L}_{p,\lambda}$ -space for all  $\lambda > 1$ , is an  $L^p(\mu)$  space, should be possible. Certainly the result should not depend on the entire theory of contractive projections for such spaces. Indeed if  $p = 2$  the  $\mathcal{L}_{2,\lambda}$  condition already implies the parallelogram law and this makes the space a Hilbert space. For  $p \neq 2$  we can see that the Clarkson inequalities are valid and these with enough finite dimensional  $l_p$ -subspaces might give a more elementary proof.

**6. Appendix.** We prove two technical results used in [1], [10]. The first is also an extension of that in [1].

**LEMMA 6.1.** [1]. *Suppose  $0 < p < \infty$  and let  $M$  be a closed subspace of  $L_p(X, \Sigma, \mu)$ . If  $(f_n)$  is a sequence in  $M$ , then there exists  $f \in M$  such that  $S(f) = \bigcup_{n=1}^{\infty} S(f_n)$ . In particular if  $\mu$  is finite or  $M$  is separable there exists  $f \in M$  such that  $J_f = J_{M^{\perp}}$ ; that is,  $f$  is a function in  $M$  of maximum support.*

*Proof.* If  $f, g \in L_p$  and  $\alpha$  is a scalar, the zero sets  $\{t \in X: (f + \alpha g)(t) = 0\}$  have disjoint intersection with  $S(f) \cup S(g)$  for differing values of  $\alpha$ . Since  $S(f) \cup S(g)$  is  $\sigma$ -finite,  $\mu(S(f) \cup S(g) \sim S(f + \alpha g)) = 0$  except, perhaps for countably many values of  $\alpha$ .

Assume, as we may, that  $\int |f_n|^p = 1$  for all  $n$ . We define, inductively, two sequences  $(\alpha_n), (\varepsilon_n)$  of positive real numbers such that, if we write  $g_n = \alpha_1 f_1 + \dots + \alpha_n f_n$ ,  $A_n = \{t \in X: |g_n(t)| \leq \varepsilon_n\}$ , and  $B_n = \{t \in X: |\alpha_{n+1} f_{n+1}(t)| \geq \varepsilon_n/2\}$ , then

- (i)  $\alpha_{n+1} < 2^{-n/p}$  and  $\varepsilon_{n+1} < \varepsilon_n/2$ ;
- (ii)  $\mu(S(g_n) \cup S(f_{n+1}) \sim S(g_{n+1})) = 0$ ;
- (iii)  $\int_{A_n \cup B_n} |f_i|^p d\mu < 2^{-n}$  ( $i = 1, 2, \dots, n$ ).

Start with  $\alpha_1 = 1$ . Suppose  $\alpha_1, \dots, \alpha_n; \varepsilon_1, \dots, \varepsilon_{n-1}$  have been chosen. Note that  $\mu(S(f_i) \sim S(g_n)) = 0$  ( $i = 1, \dots, n$ ) so if  $C_\varepsilon = \{t \in X: |g_n(t)| \leq \varepsilon\}$ ,  $\int_{C_\varepsilon} |f_i|^p d\mu \rightarrow 0$  ( $\varepsilon \rightarrow 0+$ ) for  $i = 1, \dots, n$ . Also if

$$D_\eta = \{t \in X: |f_{n+1}(t)| \geq \eta\}, \int_{D_\eta} |f_i|^p d\mu \rightarrow 0$$
 ( $\eta \rightarrow \infty$ ) for  $i = 1, \dots, n$ .

Thus we choose  $\varepsilon_n$  such that  $0 < \varepsilon_n < \varepsilon_{n-1}/2$ , and  $\int_{A_n} |f_i|^p d\mu < 2^{-n-1}$  ( $i = 1, 2, \dots, n$ ); then choose  $\eta$  such that  $\int_{D_\eta} |f_i|^p d\mu < 2^{-n-1}$  ( $i = 1, 2, \dots, n$ ), and  $\alpha_{n+1}$  such that  $0 < \alpha_{n+1} < 2^{-n/p}$ , (ii) is satisfied, and  $\alpha_{n+1}\eta < \varepsilon_n/2$ . Since  $B_n \subset D_\eta$  we also have (iii) satisfied.

By (i)  $(g_n)$  converges in  $L_p$  to an element  $f \in M$ , and  $S(f) \subset \bigcup S(f_n)$ . Let  $E = \limsup (A_n \cup B_n) = \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} (A_n \cup B_n)$ . Fix  $i$  and let  $N > i$ , then, by (iii)

$$\begin{aligned} \int_E |f_i|^p d\mu &\leq \int_{\bigcup_{N^c} (A_n \cup B_n)} |f_i|^p d\mu \\ &\leq \sum_N \int_{A_n \cup B_n} |f_i|^p d\mu \\ &\leq \sum_N 2^{-n} \\ &= 2^{1-N} \longrightarrow 0 \quad (N \longrightarrow \infty). \end{aligned}$$

Thus  $\mu(E \cap S(f_i)) = 0$  for all  $i$  and  $\mu(E \cap \bigcup S(f_n)) = 0$ . We complete our proof by showing that  $X \sim E \subset S(f)$ . If  $t \in X \sim E$  choose the smallest integer  $n$  such that  $t \notin \bigcup_{k=n}^{\infty} (A_k \cup B_k)$ , then  $|g_n(t)| > \varepsilon_n$  and  $|\alpha_k f_k(t)| < \varepsilon_{k-1}/2 < \varepsilon_n/2^{k-n}$  ( $k \geq n+1$ ). Hence

$$\begin{aligned} |g_k(t)| &\geq |g_n(t)| - |\alpha_{n+1} f_{n+1}(t)| - \cdots - |\alpha_k f_k(t)| \\ &> |g_n(t)| - \varepsilon_n(2^{-1} + \cdots + 2^{-(k-n)}) \\ &> |g_n(t)| - \varepsilon_n \end{aligned} \quad (k > n).$$

Thus  $|f(t)| = \lim_{k \rightarrow \infty} |g_k(t)| \geq |g_n(t)| - \varepsilon_n > 0$ , and we are done.

**LEMMA 6.2. [10].** *Let  $M$  be a separable subspace of  $L_p(X, \Sigma, \mu)$  ( $p \geq 1$ ) and  $T$  a bounded linear operator on  $L_p$ . Then there is a  $\sigma$ -finite set  $X_0 \in \Sigma$  and a  $\sigma$ -subring  $\Sigma_0$  of  $\Sigma$  such that  $\Sigma_0$  consists of subsets of  $X_0$  and  $L_p(X_0, \Sigma_0, \mu)$  is separable,  $T$ -invariant and contains  $M$ .*

*Proof.* The subspace  $M + TM$  is separable,  $T$ -invariant and generates a separable vector sublattice  $M_1$  of  $L_p$ . Inductively construct separable vector sublattices  $M_n$  such that  $M_n + TM_n \subset M_{n+1}$ . Then  $\text{cl} \bigcup M_n$  is a separable  $T$ -invariant closed vector sublattice of  $L_p$ . Writing  $K_1 = \text{cl} \bigcup M_n$  we have  $K_1$  closed under all band projections  $J_x$  with  $x \in K_1$ . Let  $\Sigma_1 = \{S(x) : x \in K_1\}$  then  $\Sigma_1$  is a  $\sigma$ -subring of  $\Sigma$  and if  $x, y \in K_1$  with  $x \in y^{\perp\perp}$  then  $x/y$  is  $\Sigma_1$ -measurable. If  $(f_n)$  is dense in  $K_1$ ,  $f = \Sigma 2^{-n} \|f_n\|^{-1} |f_n| \in K_1$  and  $\mu(S(x) \sim S(f)) = 0$  ( $x \in K_1$ ). Consider  $L_p(S(f), \Sigma_1, \mu)$ . It is easy to see that this is the closure of the vector sublattice spanned by  $K_1$  and the functions  $\chi_{f^{-1}(\alpha, \infty]}$  with  $\alpha$  positive rational. Thus, writing  $X_1 = S(f)$  we have

$$K_1 \subset L_p(X_1, \Sigma_1, \mu)$$

with  $L_p(X_1, \Sigma_1, \mu)$  separable. Continue inductively, we obtain a sequence  $X_1 \subset X_2 \subset \cdots \subset X_n \subset \cdots$  of  $\sigma$ -finite subsets of  $X$  and a sequence  $\Sigma_1 \subset \Sigma_2 \subset \cdots \subset \Sigma_n \subset \cdots$  of  $\sigma$ -subrings of  $\Sigma$ , such that each  $\Sigma_n$  consists of subsets of  $X_n$ ,  $L_p(X_n, \Sigma_n, \mu) + TL_p(X_n, \Sigma_n, \mu) \subset L_p(X_{n+1}, \Sigma_{n+1}, \mu)$  and each  $L_p(X_n, \Sigma_n, \mu)$  is separable.

Let  $K_0 = \text{cl} \bigcup_{n=1}^{\infty} L_p(X_n, \Sigma_n, \mu)$ . Then  $K_0$  is a separable  $T$ -invariant closed vector sublattice of  $L_p(X, \Sigma, \mu)$ . Define  $\Sigma_0 = \{S(f) : f \in K_0\}$  and find, as for  $K_1$ ,  $f \in K_0$  such that  $\mu(S(x) \sim S(f)) = 0$  ( $x \in K_0$ ). It is routine to show that  $K_0 = L_p(S(f), \Sigma_0, \mu)$ . This proves our lemma with  $X_0 = S(f)$ .

*Added in Proof* (October 1974). In a manuscript, "A local characterization of complex Banach lattices with order continuous norm," submitted to *Studia Math.*, the authors have given a necessary and sufficient condition for a complex Banach space to admit a lattice

structure so that it is a complex Banach lattice with order continuous norm. The condition is automatically satisfied if the Banach space is an  $\mathcal{L}_{p,\lambda}$  space for every  $\lambda > 1$ . This does provide an elementary proof that such spaces are  $L_p$ -spaces.

## REFERENCES

1. T. Ando, *Contractive projections in  $L_p$ -spaces*, Pacific J. Math., **17** (1966), 391-405.
2. R. G. Douglas, *Contractive projections on an  $L_1$ -space*, Pacific J. Math., **15** (1965), 443-462.
3. C. V. Duplissey, *Contractive projections in abstract Banach function spaces*, Ph. D. Dissertation, University of Texas at Austin, 1971.
4. A. Grothendieck, *Une caractérisation vectorielle métrique des espaces  $L^1$* , Canad. J. Math., **7** (1955), 552-561.
5. H. Elton Lacey and S. J. Bernau, *Characterizations and classifications of some classical Banach spaces*, Advances in Math., **12** (1974), 367-401.
6. J. Lamperti, *On the isometries of certain spaces*, Pacific J. Math., **8** (1958), 459-466.
7. J. Lindenstrauss and A. Pelczynski, *Absolutely summing operators in  $\mathcal{L}_p$ -spaces and their applications*, Studia Math., **29** (1968), 275-326.
8. M. M. Rao, *Linear operations, tensor products, and contractive projections in function spaces*, Studia Math., **38** (1970), 131-186.
9. S. Sakai,  *$C^*$ -Algebras and  $W^*$ -Algebras*, Ergebnisse der Mathematik, Bd 60, Springer-Verlag, 1971.
10. L. Tzafriri, *Remarks on contractive projections in  $L_p$ -spaces*, Israel J. Math., **7** (1969), 9-15.
11. Daniel E. Wulbert, *A note on the characterization of conditional expectation operators*, Pacific J. Math., **34** (1970), 285-288.
12. M. Zippin, *On Bases in Banach Spaces*, Ph. D. thesis, Hebrew University, Jerusalem, 1968 (Hebrew).

Received May 22, 1973. The first author's research was supported in part by NSF Grant GP 27916. The second author's research was supported by a joint grant from the National Academy of Sciences and the Polish Academy of Sciences.

UNIVERSITY OF TEXAS-AUSTIN

