

## WHICH WEIGHTED SHIFTS ARE SUBNORMAL

J. G. STAMPFLI

Let  $H$  be a Hilbert space with orthonormal basis  $\{f_j\}_{j=1}^{\infty}$ . If the operator  $T$  is defined on  $H$  by  $Tf_i = a_j f_{i+1}$  for  $i = 1, 2, \dots$ , where  $|a_i| \leq |a_{i+1}| \leq M$  for  $i = 1, 2, \dots$ , then  $T$  will be called a monotone shift. The first section of the paper examines some of the elementary properties of such operators.

Every monotone shift is hyponormal. The central portion of the paper aims at discovering which monotone shifts are subnormal. Necessary and sufficient conditions are given in terms of the  $\{a_i\}$ . These conditions make it easy to show that even the first four coefficients ( $a_1 < a_2 < a_3 < a_4$ ) may "prevent" a shift from being subnormal. However, for any  $a_1 < a_2 < a_3$  there does exist a monotone shift with these as its initial terms. In fact, the unique minimal one is constructed.

A complete description is given of subnormal monotone shifts for which  $|a_{j_0}| = |a_{j_0+1}|$  for some  $j_0$ . The paper concludes with counter-examples constructed from the machinery developed.

We are tacitly assuming that  $\lim_{j \rightarrow \infty} |a_j|$  exists, i.e.,  $T$  is a bounded operator. If  $|a_j| = |a_{j+1}|$  for  $j = 1, 2, \dots$ , then  $T$  is (up to unitary equivalence) simply a multiple of the justly famous unilateral shift.

We recall that an operator  $T$  on a Hilbert space  $H$  is subnormal if it is the restriction of a normal operator to an invariant subspace. The terms "point", "continuous" and "residual spectrum" have their usual meaning and are designated by  $\sigma_p(\cdot)$ ,  $\sigma_c(\cdot)$  and  $\sigma_r(\cdot)$  respectively.

**THEOREM 1.** *Let  $T$  be a monotone shift on  $H$  where  $A = \lim_{j \rightarrow \infty} |a_j|$ , then*

- (i)  $\|T\| = A$
- (ii)  $\sigma_r(T) = \{z: |z| < A\}$
- (iii)  $\sigma_p(T^*) = \{z: |z| < A\}$
- (iv)  $\sigma_c(T) = \sigma_c(T^*) = \{z: |z| = A\}$ .

*Proof.* Surely (i) is clear.

For  $|z_0| < A$ , consider the vector  $g = \sum_{n=1}^{\infty} z_0^n b_n f_n$  where  $b_k = 1$  if  $a_{k-1} = 0$ ,  $a_k \neq 0$  and  $b_{n+1} = z_0 b_n / \bar{a}_n$  for  $n > k$ . Since  $|z_0 / \bar{a}_n| \leq r < 1$  for  $n$  sufficiently large,  $g \in H$ . But  $(T^* - zI)g = 0$  so  $g$  is the desired eigenvector, proving (iii). The relation  $b_{n+1} = z_0 b_n / \bar{a}_n$  is necessary which implies the eigenvalue  $z_0$  is of multiplicity one.

For any  $z$ , it is clear that  $(T - zI)h \neq 0$  for  $h \in H$ , and  $h \neq 0$ .

---

Received January 26, 1965. Supported by the Office of Naval Research under contract Nonr (G)-00013-65.

Now for an arbitrary operator,  $z_0 \in \sigma_P(T^*)$  implies  $\bar{z}_0 \in \sigma_P(T) \cup \sigma_R(T)$  and (ii) follows.

It is not hard to see that

$$\sigma_P(T) \cap \{z: |z| = A\} = \phi = \sigma_P(T^*) \cap \{z: |z| = A\}$$

and hence (iv) is proved.

REMARK. If one considered a two-sided monotone shift  $Tf_j = a_j f_{j+1}$  for  $j = 0, \pm 1, \dots$ , then the above theorem is easily modified, the spectrum of  $T$  now being the annulus  $B \leq |z| \leq A$  where  $A = \lim_{j \rightarrow \infty} |a_j|$  and  $B = \lim_{j \rightarrow -\infty} |a_j|$ .

THEOREM 2. *If  $T$  is a monotone shift then  $T$  is hyponormal; that is  $\|Tg\| \geq \|T^*g\|$  for  $g \in H$ .*

*Proof.* Clear.

THEOREM 3. *If  $T$  is a monotone shift then  $T$  is not a spectral operator, (for  $T \neq 0$ ).*

*Proof.* Assume  $T$  is a spectral operator; then  $T^*$  is also a spectral operator which can be written as  $S + N$  where  $S$  is a scalar operator and  $N$  a commuting quasi-nilpotent. By Theorem 1 we know that every point in the disc  $|z| < \|T\|$  is an eigenvalue of  $T^*$ . Invoking Corollary 5 to Theorem 1 of [2] every point in the disc  $|z| < \|T\|$  is an eigenvalue of  $S$ . But  $S$  is similar to a normal operator which must have an uncountable number of distinct eigenvalues. Thus our original assumption was ill founded.

Next we would like explicitly exhibit a normal extension  $B$  of  $T$  assuming  $T$  to be subnormal. The normal extension we construct will be minimal, (see [1], [3] for a discussion of subnormal operators). From now on we will assume  $a_i \neq 0$  for  $i = 1, 2, \dots$ . There is no real loss in generality if we do so. For if  $a_1 = a_2 = \dots = a_n = 0$ , then the subspace  $M$  spanned by  $\{\varphi_1, \dots, \varphi_n\}$  is a reducing subspace of  $T$  and moreover  $T$  is normal on  $M$ . Hence we are really only concerned with what takes place in  $M^\perp$ . Let us designate the original basis for  $H \equiv H^{(1)}$  by  $\{\varphi_j^{(1)}\}_{j=1}^\infty$ . We shall use the terms  $a_j$  and  $a_j^{(1)}$  interchangeably as a notational convenience. It will be to our advantage to assume that  $a_j^{(1)} > 0$ , for  $j = 1, 2, \dots$ , and we may do this by simply rotating the  $\varphi_j^{(1)}$ 's of the original basis by  $e^{i\theta_j}$ .

Now set  $B^*\varphi_j^{(1)} = a_{j-1}^{(1)}\varphi_{j-1}^{(1)} + b_j^{(2)}\varphi_j^{(2)}$  where  $\|\varphi_j^{(2)}\| = 1$  and  $\varphi_j^{(2)}$  is orthogonal to  $H^{(1)}$  for  $j = 1, 2, \dots$ . Since  $\|B^*\varphi_j^{(1)}\|^2 = (a_{j-1}^{(1)})^2 + |b_j^{(2)}|^2 = \|B\varphi_j^{(1)}\|^2 = (a_j^{(1)})^2$  we may conclude that  $|b_j^{(2)}|^2 = (a_j^{(1)})^2 - (a_{j-1}^{(1)})^2$ , ( $a_0^{(1)} = 0$ ). We claim that the  $\varphi_j^{(2)}$ 's are orthogonal, for

$$b_i^{(2)}b_j^{(2)}(\varphi_i^{(2)}, \varphi_j^{(2)}) = (B^*\varphi_i^{(1)}, B^*\varphi_j^{(1)}) = (B\varphi_i^{(1)}, B\varphi_j^{(1)}) = 0,$$



where  $b_j^{(2)} = ((\alpha_j^{(1)})^2 - (\alpha_{j-1}^{(1)})^2)^{1/2}$  and  $b_j^{(2)}\alpha_j^{(2)} = b_{j+1}^{(2)}\alpha_j^{(1)}$ . We have assumed that all the  $\alpha_j^{(1)}$ 's  $\neq 0$  and have shown that if  $b_{j_0}^{(2)} = 0$  then  $b_k^{(2)} = 0$  for  $k \geq j_0$ . In this case,  $H^{(2)}$  would be finite dimensional.

We will now assume  $B$  has been defined on  $H^{(1)}, \dots, H^{(n)}$  where  $\{\varphi_j^{(1)}\}, \dots, \{\varphi_j^{(n)}\}$  are the respective bases and  $B\varphi_j^{(k)} = \alpha_j^{(k)}\varphi_{j+1}^{(k)} + b_j^{(k)}\varphi_j^{(k-1)}$  for  $k = 2, \dots, n$ . We also note that all  $b_j^{(k)} \neq 0$ . That is, if  $b_1^{(k)}, \dots, b_{j_k}^{(k)}$  are nonzero and  $b_{j_k+1}^{(k)} = 0$  then  $b_j^{(k)} = 0$  for  $j \geq j_k + 1$  and  $H^{(k)}$  is spanned by  $\{\varphi_1^{(k)}, \dots, \varphi_{j_k}^{(k)}\}$ . Further  $\alpha_j^{(k)} \neq 0$  for  $j = 1, \dots, j_k - 1$ . The above can be considered to be the induction hypothesis.

We set  $B^*\varphi_j^{(n)} = \alpha_{j-1}^{(n)}\varphi_{j-1}^{(n)} + b_j^{(n+1)}\varphi_j^{(n+1)}$  for  $j = 1, 2, \dots$ , where  $\|\varphi_j^{(n+1)}\| = 1$ , and  $\varphi_j^{(n+1)}$  is orthogonal to  $H^{(n)} \oplus \dots \oplus H^{(1)}$ . Since  $\|B^*\varphi_j^{(n)}\| = \|B\varphi_j^{(n)}\|$  we have  $|b_j^{(n+1)}|^2 = (\alpha_j^{(n)})^2 + (b_j^{(n)})^2 - (\alpha_{j-1}^{(n)})^2 (\alpha_0^{(n)} = 0$  by definition). Now

$$\begin{aligned} (B^*\varphi_i^{(n)}, B^*\varphi_j^{(n)}) &= b_i^{(n+1)}b_j^{(n+1)}(\varphi_i^{(n+1)}, \varphi_j^{(n+1)}) \\ &= (B\varphi_i^{(n)}, B\varphi_j^{(n)}) = 0 \quad \text{for } i \neq j. \end{aligned}$$

Thus  $\varphi_i^{(n+1)}$  and  $\varphi_j^{(n+1)}$  are orthogonal for we may assume  $b_i^{(n+1)}, b_j^{(n+1)}$  are nonzero else  $\varphi_{i \text{ or } j}^{(n+1)}$  would not be defined. Thus  $b_j^{(n+1)}$  may be taken positive. Consider

$$\begin{aligned} (\varphi_{j+1}^{(n+1)}, B^*B\varphi_j^{(n)}) &= (\varphi_{j+1}^{(n+1)}, B^*(\alpha_j^{(n)}\varphi_{j+1}^{(n)} + b_j^{(n)}\varphi_j^{(n-1)})) \\ &= (\varphi_{j+1}^{(n+1)}, \alpha_j^{(n)}b_{j+1}^{(n+1)}\varphi_{j+1}^{(n+1)}) = \alpha_j^{(n)}b_{j+1}^{(n+1)} \\ &= (\varphi_{j+1}^{(n+1)}, BB^*\varphi_j^{(n)}) = (\varphi_{j+1}^{(n+1)}, B(\alpha_{j-1}^{(n)}\varphi_{j-1}^{(n)} + b_j^{(n+1)}\varphi_j^{(n+1)})) \\ &= (\varphi_{j+1}^{(n+1)}, b_j^{(n+1)}B\varphi_j^{(n+1)}). \end{aligned}$$

This implies  $B\varphi_j^{(n+1)} = b_j^{(n+1)}\varphi_j^{(n+1)} + \alpha_j^{(n+1)}\varphi_{j+1}^{(n+1)} + f_j$  where  $(f_j, \varphi_{j+1}^{(n+1)}) = 0$  and  $\alpha_j^{(n+1)}$  satisfies the relation  $\alpha_j^{(n+1)}b_j^{(n+1)} = \alpha_j^{(n)}b_{j+1}^{(n+1)}$ , for  $j = 1, 2, \dots$ .

Now let us clear up the zero, nonzero relation between the  $\alpha$ 's and the  $b$ 's. By the induction hypothesis either all  $\alpha_j^{(n)}$ 's  $\neq 0$  for  $j = 1, 2, \dots$ , or there are a finite number and all but the last is nonzero. In either case assume  $b_k^{(n+1)} = 0$  and  $b_{k+1}^{(n+1)} \neq 0$ ; then  $\alpha_k^{(n+1)}b_k^{(n+1)} = 0 = \alpha_k^{(n)}b_{k+1}^{(n+1)} \neq 0$  (for there are at least  $k$  nonzero  $\alpha_j^{(n)}$ 's since the  $\dim H^{(n+1)} \leq \dim H^{(n)}$ ). This is impossible. Now assume  $b_j^{(n+1)} \neq 0$  for  $j = 1, 2, \dots, k$ , and  $b_j^{(n+1)} = 0$  for  $j > k$ , i.e.,  $H^{(n+1)}$  is  $k$ -dimensional. Then  $\alpha_j^{(n+1)} = \alpha_j^{(n)}b_{j+1}^{(n+1)}/b_j^{(n+1)}$  is well defined for  $j \leq k$  and since  $H^{(n)}$  was at least  $k$ -dimensional,  $\alpha_j^{(n)} \neq 0$  for  $j = 1, \dots, k - 1$ ; hence  $\alpha_j^{(n+1)} \neq 0$  for  $j = 1, \dots, k - 1$  as we wished to show.

Let us go back to the defining relation

$$B\varphi_j^{(n+1)} = \alpha_j^{(n+1)}\varphi_{j+1}^{(n+1)} + b_j^{(n+1)}\varphi_j^{(n)} + f_j.$$

Since

$$(\varphi_i^{(n+1)}, B^*B\varphi_j^{(n)}) = 0 \quad \text{for } i \neq j + 1$$

we have

$$\begin{aligned}
 (\varphi_i^{(n+1)}, BB^* \varphi_j^{(n)}) &= (\varphi_i^{(n+1)}, b_j^{(n+1)}(a_j^{(n+1)} \varphi_{j+1}^{(n+1)} + b_j^{(n+1)} \varphi_j^{(n)} + f_j)) \\
 + (\varphi_i^{(n+1)}, Ba_{j-1}^{(n)} \varphi_{j-1}^{(n)}) &= (\varphi_i^{(n+1)}, f_j) = 0 \quad \text{for } i \neq j + 1.
 \end{aligned}$$

Now  $(\varphi_{j+1}^{(n+1)}, f_j) = 0$  by definition and hence  $f_j$  is orthogonal to  $H^{(1)} \oplus \dots \oplus H^{(n+1)}$ . But

$$\begin{aligned}
 (f_j, B^* B \varphi_j^{(n)}) &= (f_j, B^*(a_j^{(n)} \varphi_{j+1}^{(n)} + b_j^{(n)} \varphi_j^{(n-1)})) \\
 &= 0 = (f_j, BB^* \varphi_j^{(n)}) \\
 &= (f_j, B(a_{j-1}^{(n)} \varphi_{j-1}^{(n)} + b_j^{(n+1)} \varphi_j^{(n+1)})) \\
 &= (f_j, b_j^{(n+1)} B \varphi_j^{(n+1)}) \\
 &= (f_j, b_j^{(n+1)}(a_j^{(n+1)} \varphi_{j+1}^{(n+1)} b_j^{(n+1)} \varphi_j^{(n+1)} + f_j)) \\
 &= b_j^{(n+1)} \|f_j\| \quad \text{and since } b_j^{(n+1)} \neq 0
 \end{aligned}$$

we conclude  $f_j = 0$ . Thus  $B \varphi_j^{(n+1)} = a_j^{(n+1)} \varphi_{j+1}^{(n+1)} + b_j^{(n+1)} \varphi_j^{(n)}$  where  $b_j^{(n+1)} \neq 0$  for  $j = 1, \dots, (\dim H^{(n+1)})$  and  $a_j^{(n+1)} \neq 0$  for  $j = 1, \dots, (\dim H^{(n+1)} - 1)$ . It should help to present  $B$  as an infinite matrix with respect to the basis  $\{\varphi_j^{(n)}\}$  where  $j = 1, 2, \dots, \dim H^{(n)}$  and  $n = 1, 2, \dots$ .

$$B = \begin{vmatrix} F^{(1)} & G^{(2)} & 0 & & \\ 0 & F^{(2)} & G^{(3)} & & 0 \\ 0 & 0 & F^{(3)} & \cdot & \\ & & & \cdot & \cdot \\ & 0 & & & \cdot \\ & & & & \cdot \end{vmatrix}$$

where

$$F^{(n)} = \begin{vmatrix} 0 & & & & \\ \alpha_1^{(n)} & 0 & & & \\ & \alpha_2^{(n)} & 0 & & \\ & & \alpha_3^{(n)} & \cdot & \\ & & & \cdot & \cdot \\ & & & & \cdot \\ & & & & \cdot \end{vmatrix}$$

and

$$G^{(n)} = \begin{vmatrix} b_1^{(n)} & & & & \\ & b_2^{(n)} & & & \\ & & b_3^{(n)} & & \\ & & & \cdot & \\ & & & & \cdot \\ & & & & \cdot \end{vmatrix}$$

Now that we have a normal extension of  $T$  in concrete form it is not difficult to read off some necessary conditions for the subnormality of  $T$ . In fact, it is clear that:

- ( I )  $(a_j^{(n)})^2 + (b_j^{(n)})^2 \geq (a_{j-1}^{(n)})^2$  for  $n = 2, 3, \dots$ , and  $j = 1, 2, \dots$ , (otherwise it would not be possible to define  $b_j^{(n+1)}$  compatibly with the normality of  $B$ ).

- (II) If  $b_j^{(n)} = 0$  then  $b_{j+1}^{(n)} = 0$  for  $j = 1, 2, \dots$ , and  $n = 2, 3, \dots$ , (else the equation  $\alpha_j^{(n)} b_j^{(n)} = \alpha_j^{(n-1)} b_{j+1}^{(n)}$  could not be satisfied).
- (III) There exists a constant  $M$  such that  $|\alpha_j^{(n)}| \leq M$  and  $|b_j^{(n)}| \leq M$  for  $n = 2, 3, \dots$ , and  $j = 1, 2, \dots$ , (else the normal extension would not be bounded). These conditions are also sufficient. More precisely:

**THEOREM 4.** *Let  $\{\alpha_j^{(1)}\}$  be a sequence such that  $0 < \alpha_1^{(1)} \leq \alpha_2^{(1)} \leq \dots$  and define  $T\varphi_j^{(1)} = \alpha_j^{(1)}\varphi_{j+1}^{(1)}$  for  $j = 1, 2, \dots$ , where  $\{\varphi_j^{(1)}\}_{j=1}^\infty$  is an orthonormal basis for  $H^{(1)}$ . The operator  $T$  is subnormal if and only if*

( I )  $(\alpha_j^{(n)})^2 + (b_j^{(n)})^2 - (\alpha_{j-1}^{(n)})^2 \geq 0$

( II )  $b_j^{(n)} = 0 \Rightarrow b_{j+1}^{(n)} = 0$

(III) *there exists a constant  $M$  such that  $|\alpha_j^{(n)}| \leq M$  and  $|b_j^{(n)}| \leq M$  for  $n = 2, 3, \dots$ , and  $j = 1, 2, \dots$ , where*

$$(b_j^{(n+1)})^2 = + [(\alpha_j^{(n)})^2 + (b_j^{(n)})^2 - (\alpha_{j-1}^{(n)})^2]^{1/2}$$

and  $\alpha_j^{(n+1)} = \alpha_j^{(n)} b_{j+1}^{(n+1)} / b_j^{(n+1)}$  (if  $b_{j_0}^{(n)} = 0$  then  $\alpha_{j_0}^{(n)}$  is taken to be zero).

*Proof.* The necessity of (I), (II) and (III) has already been proved. In the other direction, since conditions (I) and (II) are satisfied the recursive definitions of the  $\alpha_j^{(n+1)}$  and  $b_j^{(n+1)}$  make sense. We now set  $B\varphi_j^{(n)} = \alpha_j^{(n)}\varphi_{j+1}^{(n)} + b_j^{(n)}\varphi_j^{(n-1)}$  where  $\{\varphi_j^{(n)}\}$  is an orthonormal basis for  $H^{(n)}$ . If  $b_{j_0}^{(n)} = 0$  then  $\varphi_{j_0}^{(n)}$  is omitted from the basis. With this definition it should be clear from the former construction that

$$(B\varphi_j^{(n)}, B\varphi_k^{(m)}) = (B^*\varphi_j^{(n)}, B^*\varphi_k^{(m)}) \quad \text{for all } n, m, j, k.$$

(Here as elsewhere all should be interpreted as vacuously covering the case when  $\varphi_j^{(n)}$  has been omitted from the basis.) Thus  $B$  is defined and satisfies  $\|Bx\| = \|B^*x\|$  on finite linear combinations of the basis elements. From (III) we may infer that  $B$  is bounded: certainly  $\|B\| \leq 2M$ . Hence  $B$  is defined and is normal on all of  $\hat{H} = \sum \oplus H^{(n)}$ .

We remark that the  $\alpha_j^{(1)}$ 's were taken to be real only as a notational convenience. To modify the theorem for an arbitrary monotone shift one need only add absolute value signs in the appropriate places.

Let  $\alpha_1, \dots, \alpha_n$  be given. If there exist elements  $\alpha_{n+1}, \alpha_{n+2}, \dots$ , such that  $T\varphi_j = \alpha_j \varphi_{j+1}$  for  $j = 1, 2, \dots$ , is a subnormal operator, we will call  $T$  a subnormal completion of  $\alpha_1, \dots, \alpha_n$ .

In Theorem 5 we will show that for any  $\alpha_1, \alpha_2, \alpha_3$  where  $|\alpha_1| < |\alpha_2| < |\alpha_3|$ , there exists a subnormal completion. However, one can specify  $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ , where  $\alpha_1 < \alpha_2 < \alpha_3 < \alpha_4$ , in such a manner that they can not be completed to form a subnormal monotone shift. This can be easily done by making use of necessary condition (I).

Consider  $(a_3^{(2)})^2 + (b_3^{(2)})^2 \geq (a_2^{(2)})^2$  which is equivalent to:

$$(a_3^2 - a_2^2) + a_3^2 \frac{a_4^2 - a_3^2}{a_3^2 - a_2^2} \geq a_2^2 \frac{a_3^2 - a_2^2}{a_2^2 - a_1^2}$$

or

$$(1) \quad a_4^2 \geq a_3^2 + \frac{a_1^2}{a_3^2} \frac{(a_3^2 - a_2^2)^2}{a_2^2 - a_1^2}.$$

If  $a_4$  is chosen so as not to satisfy (1) then clearly no choice of the remaining  $a_j$ 's will make the resulting shift subnormal. For example, if  $a_1 = 1$ ,  $a_2 = 2^{1/2}$ ,  $a_3 = 3^{1/2}$ , then for  $3^{1/2} \leq a_4 < (10/3)^{1/2}$  there exists no subnormal completion.

**THEOREM 5.** *Given  $a_1, a_2, a_3$  where  $0 < |a_1| < |a_2| < |a_3|$  then there exists a subnormal completion of  $a_1, a_2, a_3$ . Moreover, if  $S$  is any subnormal completion of  $a_1, a_2, a_3$ , then*

$$\begin{aligned} \|S\|^2 \geq & 1/2 \left[ |a_2|^2 \frac{|a_3|^2 - |a_1|^2}{|a_2|^2 - |a_1|^2} \right. \\ & \left. + \left( \left[ |a_2|^2 \frac{|a_3|^2 - |a_1|^2}{|a_2|^2 - |a_1|^2} \right]^2 - 4 |a_1|^2 |a_2|^2 \frac{|a_3|^2 - |a_2|^2}{|a_2|^2 - |a_1|^2} \right)^{1/2} \right]. \end{aligned}$$

Further there is, up to unitary equivalence, exactly one subnormal completion for which equality holds.

*Proof.* We may take  $a_1, a_2, a_3$  to be positive. The  $a_n$ 's for the minimal completion are given by the following rule:

$$(2) \quad a_n^2 \frac{a_{n+1}^2 - a_n^2}{a_n^2 - a_{n-1}^2} = a_{n-1}^2 \frac{a_n^2 - a_{n-1}^2}{a_{n-1}^2 - a_{n-2}^2} - (a_n^2 - a_{n-1}^2).$$

Since  $a_{n-1}^2 / (a_{n-2}^2 - a_{n-3}^2) > 1$  the right side is strictly positive and  $a_{n+1} > a_n$ . The defining relation (2) is simply the necessary condition  $(a_j^{(2)})^2 + (b_j^{(2)})^2 - (a_{j-1}^{(2)})^2 \geq 0$  with equality holding.

Next we will find a more convenient expression for  $a_n$ . From (2) we have:

$$(3) \quad \frac{a_{n+1}^2 - a_n^2}{a_n^2 - a_{n-1}^2} = \frac{a_{n-2}^2}{a_n^2} \cdot \frac{a_n^2 - a_{n-1}^2}{a_{n-1}^2 - a_{n-2}^2}$$

and so by induction,

$$(4) \quad \frac{a_{n+1}^2 - a_n^2}{a_n^2 - a_{n-1}^2} = \frac{a_2^2 \cdot a_1^2}{a_n^2 \cdot a_{n-1}^2} \cdot \frac{a_3^2 - a_2^2}{a_2^2 - a_1^2} = \frac{C}{a_n^2 a_{n-1}^2}$$

<sup>1</sup> Note that  $a_j^2$  means  $(a_j)^2$  not  $a_j^{(2)}$



There is no difficulty in defining  $a_1^{(n)}$  from (i) and  $b_1^{(n+1)}$  from (ii). From (ii) we see that  $b_1^{(n+1)}$ 's are increasing and  $b_1^{(n+1)} > a_1^{(n)}$ . Hence from (i) we have  $1 < b_1^{(n)}/a_1^{(n-1)} = b_2^{(n)}/a_1^{(n)}$  or  $(b_2^{(n)})^2 - (a_2^{(n)})^2 - (a_1^{(n)})^2 > 0$  so there is no obstacle to defining  $b_2^{(n+1)}$  by (iii). Now to show boundedness. From (iii) we see  $\{b_2^{(n)}\}$  is decreasing and thus converges which also implies that  $\{a_1^{(n)}\}$  converges to zero. Combining (ii) and (iii) we have

$$(b_1^{(n+1)})^2 - (b_1^{(n)})^2 = (b_2^{(n)})^2 - (b_2^{(n+1)})^2.$$

Thus

$$(b_1^{(n+1)})^2 - (b_2^{(3)})^2 = (b_2^{(3)})^2 - (b_2^{(n+1)})^2$$

which implies  $\{b_1^{(n)}\}$  converges since  $\{b_2^{(n)}\}$  does. Now  $B$  the normal extension of  $T$  must be a bounded operator.

We will compute the norm of this completion from (6). Let  $A = \lim_{j \rightarrow \infty} \|a_j\|^2 = \|T\|^2$ . Then (6) yields

$$(7) \quad A = a_2^2 + C \left[ \frac{1}{a_1^2} - \frac{1}{A} \right]$$

or

$$A^2 - (a_2^2 + C/a_1^2)A - C = 0.$$

Recalling that  $C = a_1^2 a_2^2 [(a_3^2 - a_2^2)/(a_2^2 - a_1^2)]$  and selecting the proper root of the quadratic equation yields

$$(8) \quad A = \frac{1}{2} \left[ a_2^2 \frac{a_3^2 - a_1^2}{a_2^2 - a_1^2} + \left\{ \left[ a_2^2 \frac{a_3^2 - a_1^2}{a_2^2 - a_1^2} \right]^2 - 4a_1^2 a_2^2 \frac{a_3^2 - a_2^2}{a_2^2 - a_1^2} \right\}^{1/2} \right]$$

(the expression inside the radical is positive).

Before we show that the subnormal operator  $T$  which we constructed to complete  $a_1, a_2, a_3$  is minimal in norm, we will prove that it is an efficient subnormal completion in another sense. Let  $S\varphi_j = c_j\varphi_{j+1}$  be any subnormal operator with  $c_i = a_i$  for  $i = 1, 2, 3$ . Rather than be inundated by absolute value signs we will assume that both the  $a_i$ 's and the  $c_j$ 's are positive.

Now the necessary condition (I)

$$(a_j^{(2)})^2 + (b_j^{(2)})^2 - (a_{j-1}^{(2)})^2 \geq 0$$

applied to the operator  $S$  yields the following analogue to (2):

$$(9) \quad \frac{c_n^2 c_{n+1}^2 - c_n^2}{c_n^2 - c_{n-1}^2} \geq c_{n-1}^2 \frac{c_n^2 - c_{n-1}^2}{c_{n-1}^2 - c_{n-2}^2} - (c_n^2 - c_{n-1}^2)$$

and if we argue as before we can reduce this to

$$(10) \quad c_{n+1}^2 \geq a_2^2 + C \left[ \frac{1}{a_1^2} - \frac{1}{c_n^2} \right].$$

If the completion  $c_n$  is different from the one given above then for some first  $k$  we have  $c_k > a_k$  (it can not be less by the necessary condition (I)). But then we have

$$c_{k+1}^2 \geq a_2^2 + C \left[ \frac{1}{a_1^2} - \frac{1}{c_k^2} \right] > a_2^2 + C \left[ \frac{1}{a_1^2} - \frac{1}{a_k^2} \right] = a_{k+1}^2.$$

Hence  $c_j > a_j$  for  $j \geq k$ .

Now we will prove the last statement of the theorem. Let  $S$  and  $c_j$  be as above. Let  $k$  be the smallest integer such that  $a_{k+2} = c_{k+2}$  but  $a_{k+3} < c_{k+3}$ . Using the necessary condition (I) we arrive at

$$(11a) \quad a_{n+1}^2 = a_n^2 + D \left[ \frac{1}{a_{n-1}^2} - \frac{1}{a_n^2} \right]$$

and

$$(11b) \quad a_{n+1}^2 \geq c_n^2 + \hat{D} \left[ \frac{1}{c_{n-1}^2} - \frac{1}{c_n^2} \right]$$

for  $n \geq k+3$  where

$$D = a_{k+2}^2 a_{k+1}^2 \frac{a_{k+3}^2 - a_{k+2}^2}{a_{k+2}^2 - a_{k+1}^2} \quad \text{and} \quad \hat{D} = c_{k+2}^2 c_{k+1}^2 \frac{c_{k+3}^2 - c_{k+2}^2}{c_{k+2}^2 - c_{k+1}^2}.$$

These are obtained by a restricted telescoping of (3) and (9). Since  $c_{k+1} = a_{k+1}$ ,  $c_{k+2} = a_{k+2}$  and  $c_{k+3} > a_{k+3}$ , certainly  $\hat{D} > D$ . Now from (11a) and (11b) we obtain

$$(12a) \quad a_{n+1}^2 = a_{k+2}^2 + D \left[ \frac{1}{a_{k+1}^2} - \frac{1}{a_n^2} \right]$$

and

$$(12b) \quad c_{n+1}^2 \geq c_n^2 + \hat{D} \left[ \frac{1}{c_{k+1}^2} - \frac{1}{c_n^2} \right] \quad \text{for } n \geq k+3.$$

If we let  $n \rightarrow \infty$  in (12a) and (12b) we obtain

$$(13a) \quad A = a_{k+2}^2 + D \left[ \frac{1}{a_{k+1}^2} - \frac{1}{A} \right]$$

and

$$(13b) \quad \hat{A} \geq c_{k+2}^2 + \hat{D} \left[ \frac{1}{a_{k+1}^2} - \frac{1}{\hat{A}} \right]$$

or

$$\hat{A} \geq a_{k+2}^2 + \hat{D} \left[ \frac{1}{a_{k+1}^2} - \frac{1}{\hat{A}} \right].$$

Now in (13a) the solution  $A$  is a strictly increasing function of  $D$  and since  $\hat{D} > D$  we must have  $\hat{A} > A$ . But  $\|T\|^2 = A < \hat{A} = \|S\|^2$  and thus we have shown there is exactly one subnormal completion of  $a_1, a_2, a_3$  with the norm given by (8) (again we note that operators are specified up to unitary equivalence only).

REMARK. The above construction of a subnormal completion of  $a_1, a_2, a_3$  can be used in slightly more general circumstances. It has been pointed out that  $a_1, a_2, a_3, a_4$  may not have a subnormal completion. If however  $a_4$  satisfies the necessary condition

$$a_4^2 \geq a_3^2 + a_1^2 \frac{(a_3^2 - a_2^2)^2}{a_2^2 - a_1^2}$$

then one can obtain a subnormal completion by defining  $a_n$  for  $n \geq 5$  as in (2). When one has  $a_1, \dots, a_5$  this definition does not work for reasons sufficiently complicated to remain unmentioned here.

THEOREM 6. Let  $T$  be a monotone shift,  $T\varphi_j = a_j\varphi_{j+1}$ . Assume further that

- (i)  $a_j \neq 0$  for  $j = 1, 2, \dots$
- (ii)  $T$  is subnormal
- (iii)  $|a_k| = |a_{k+1}|$  for some  $k$

then  $|a_j| = |a_{j+1}|$  for  $j = 2, 3, \dots$ , and  $a_1$  is arbitrary.

Proof. We will assume that the  $a_j$ 's are positive. Let  $k$  be the smallest integer such that  $a_k = a_{k+1}$ . If  $k \leq 2$  then by Theorem 4 of [5] we are through. For the case  $k \geq 3$  we now make use of Theorem 4. We observe that

$$a_k^{(2)} = a_k \left[ \frac{a_{k+1}^2 - a_k^2}{a_k^2 - a_{k-1}^2} \right]^{1/2} = 0$$

and

$$a_{k-1}^{(2)} = a_{k-1}^2 \left[ \frac{a_k^2 - a_{k-1}^2}{a_{k-1}^2 - a_{k-2}^2} \right]^{1/2} \neq 0$$


and

$$b_k^{(2)} = [a_k^2 - a_{k-1}^2]^{1/2}.$$

Now a necessary condition for the subnormality of  $T$  is just that

$$\begin{aligned}
 0 &\leq (a_k^{(2)})^2 + (b_k^{(2)})^2 - (a_{k-1}^{(2)})^2 \\
 &= (a_k^2 - a_{k-1}^2) - \frac{a_{k-1}^2}{a_{k-1}^2 - a_{k-2}^2} (a_k^2 - a_{k-1}^2) .
 \end{aligned}$$

But that implies  $(a_{k-1}^2/a_{k-1}^2)(a_{k-2}^2) \leq 1$  or  $a_{k-2} = 0$  contrary to hypothesis. This still leaves the existence of monotone shifts unsettled for  $a_1 < a_2 = a_j$  for  $j = 2, 3, \dots$ . We will show that these are all subnormal by writing down the subnormal extension. (The reader may do this by simply following the “directions” in Theorem 4.)



$$p = (a_2^2 - a_1^2)^{1/2}$$

**COROLLARY.** *The space  $H^{(2)}$  is either infinite dimensional or  $\dim H^{(2)} \leq 2$ . This is just another way of stating Theorem (6).*

**REMARK.** This corollary has an analogue for the spaces  $H^{(n)}$ . We state without proof that either  $H^{(n)}$  is infinite dimensional or  $\dim H^{(n)} \leq n$ . In the proof of Theorem 5 we constructed a subnormal operator where  $\dim H^{(3)} = 2$ . Given  $a_1, a_2, a_3, a_4$ , where  $a_4$  satisfies the necessary condition (I), the construction yields an operator where  $\dim H^{(3)} = 3$ .

Using the last theorem we can construct several counter-examples. In [4] we encounter the following:

**THEOREM.** *If  $T$  is hyponormal and  $T^n$  is normal for some integer  $n$  then  $T$  is normal.*

The question arises whether the theorem remains valid with normal replaced by subnormal (in both hypothesis and conclusion). The answer is no. Define  $T\varphi_1 = (1/4)\varphi_2$ ,  $T\varphi_2 = (1/2)\varphi_3$ ,  $T\varphi_i = \varphi_{i+1}$  for  $i \geq 3$ . Clearly  $T$  is hyponormal, but not subnormal by the previous theorem, We claim  $T^n$  is subnormal for  $n \geq 2$ . For  $T^2\varphi_1 = (1/8)\varphi_3$ ,  $T^2\varphi_2 = (1/2)\varphi_4$

and  $T^2\varphi_i = \varphi_{i+2}$  for  $i \geq 3$ . Thus if we let  $g_j = \varphi_{2j-1}$  and  $h_j = \varphi_{2j}$  for  $j = 1, 2, \dots$ , then we may write  $T^2$  as the direct sum of operators acting on  $H_1 = \{\text{span of } g_j\}$  and  $H_2 = \{\text{span of } h_j\}$ . Now if  $Fg_1 = (1/8)g_2$  and  $F_1g_j = g_{j+1}$  for  $j \geq 2$ , then  $F_1$  is subnormal on  $H_1$ . Similarly, if  $F_2h_1 = (1/2)h_2$  and  $F_2h_j = h_{j+1}$  for  $j \geq 2$ , then  $F_2$  is subnormal on  $H_2$ . But  $T^2$  on  $H$  equals  $F_1 \oplus F_2$  on  $H_1 \oplus H_2$  and hence  $T^2$  is subnormal. For  $n \geq 3$ ,  $T^n$  can be shown to be subnormal by a similar argument.

We will now exhibit two subnormal operators which are similar but whose (minimal) normal extensions are not. Let  $\{\varphi_j\}_{j=1}^\infty$  be an orthonormal basis for  $H$ . Define  $T\varphi_j = \varphi_{j+1}$  for  $j = 1, 2, 3, \dots$ , and  $S\varphi_1 = (1/2)\varphi_2$ ,  $S\varphi_j = \varphi_{j+1}$  for  $j = 2, 3, \dots$ . Then  $S$  and  $T$  are subnormal by Theorem 6, they are similar and, in fact, the operator  $Q$  given by  $Q\varphi_1 = (1/2)\varphi_1$ ,  $Q\varphi_j = \varphi_j$  for  $j \geq 2$ , effects the similarity. We claim that the minimal normal extensions are not similar. If they were, they would have to be unitarily equivalent since they are normal. The normal extension of  $T$  is unitary while the normal extension of  $S$  clearly is not which completes the argument.

#### BIBLIOGRAPHY

1. J. Bram, *Subnormal operators*, Duke Math. J. **22** (1955), 75-94.
2. S. R. Foguel, *The relation between a spectral operator and its scalar part*, Pacific J. Math. **8** (1958), 51-65.
3. P. R. Halmos, *Normal dilations and extensions of operators*, Summa Bras. Math. **2** (1950), 124-134.
4. J. G. Stampfli, *Hyponormal operators*, Pacific J. Math. **12** (1962), 1453-1458.
5. ———, *Hyponormal operators and spectral density*, Trans. Amer. Math. Soc., (to appear).

UNIVERSITY OF MICHIGAN

