

## ON $p$ -ADIC PROPERTIES OF THE EICHLER-SELBERG TRACE FORMULA II

M. KOIKE

### Introduction

Let  $\mathfrak{S}_k$  be the space of cusp forms of weight  $k$  with respect to  $SL(2, \mathbf{Z})$ . Let  $p$  be a prime number and let  $T_k(p)$  be the Hecke operator of degree  $p$  acting on  $\mathfrak{S}_k$  as a linear endomorphism. Put  $H_k(X) = \det(I - T_k(p)X + p^{k-1}X^2I)$ , where  $I$  is the identity operator on  $\mathfrak{S}_k$ .  $H_k(X)$  is a polynomial with coefficients of rational integers, which is called the Hecke polynomial.

In this paper, we shall prove the congruences between Hecke polynomials:

**THEOREM.** *Let  $p \geq 5$  be a prime number and let  $\alpha$  be a positive integer. Let  $k$  be an even positive integer such that  $k \geq 2\alpha + 2$  and  $\dim_C \mathfrak{S}_{k+p\alpha-p\alpha-1} < p^{k-\alpha-1}$ . Then we have*

$$H_{k'}(X) \equiv H_k(X) \pmod{p^\alpha \mathbf{Z}[X]}$$

for every even positive integer  $k' > k$  satisfying  $k' \equiv k \pmod{p^\alpha - p^{\alpha-1}}$ .

In the case of  $\alpha = 1$ , our theorem is a weaker version of the property of contraction of  $U_p$ , which was proved by Serre. The proof of our theorem makes essential use of the  $p$ -adic properties of the Eichler-Selberg trace formula which is finer than what was proved in our previous paper [2].

### § 1. Congruences between traces of Hecke operators.

We fix a prime number  $p$  once and for all. For each positive integer  $n$ , let  $T_k(n)$  be the Hecke operator of degree  $n$  acting on  $\mathfrak{S}_k$  as a linear endomorphism. The Eichler-Selberg trace formula for  $T_k(n)$  reads as follows:

---

Received May 12, 1976.

$$(1) \quad \begin{aligned} \operatorname{tr} T_k(n) = & \sum_{\{\rho, \rho'\}} \sum_{\mathfrak{o} \ni \rho} - \frac{h_{\mathfrak{o}}}{w_{\mathfrak{o}}} F^{(k-2)}(\rho, \rho') - \sum'_{\substack{d|n \\ d>0, d \leq \sqrt{n}}} d^{k-1} \\ & + \delta(\sqrt{n}) \frac{k-1}{12} n^{k/2-1} + \begin{cases} 0 & (k > 2), \\ \sum_{\substack{d|n \\ d>0}} d & (k = 2), \end{cases} \end{aligned}$$

where we use the same notations as in [2].

We shall prove finer congruences between traces of Hecke operators than what was proved in our previous paper [2]. Our result is as follows:

**PROPOSITION.** *We assume  $p \geq 5$ . Let  $m$  and  $\alpha$  be positive integers. Put  $\operatorname{ord}_p m = \beta$ . Let  $k'$  and  $k$  be even positive integers satisfying (1)  $k' \equiv k \pmod{p^\alpha - p^{\alpha-1}}$  and (2)  $k' > k \geq \operatorname{Max}\{2\alpha + 2, \alpha + \beta + 2\}$ . Then we have*

$$\operatorname{tr} T_{k'}(p^m) \equiv \operatorname{tr} T_k(p^m) \pmod{p^{\alpha+\beta}}.$$

*Remark.* In order to prove congruences between traces of Hecke operators in our previous paper, we made use of the property that  $h_{\mathfrak{o}}$  is merely a rational integer. On the other hand, the proof of Proposition makes essential use of the fact that  $h_{\mathfrak{o}}$  is the number of proper  $\mathfrak{o}$ -ideal classes.

*Proof.* We consider the trace formula for  $T_k(p^m) \pmod{p^{\alpha+\beta}}$ . Since  $k \geq 4$ , the fourth summand is equal to zero. By the condition (2), the second (resp. third) summand is proved to be congruent to one (resp. zero)  $\pmod{p^{\alpha+\beta}}$ . Let us deal with the first summand. Let  $K$  be an imaginary quadratic field which contains  $\rho$  and  $\rho'$  and let  $\left(\frac{K}{p}\right)$  denote Kronecker's symbol. In the case of  $\left(\frac{K}{p}\right) = -1$  or  $0$ ,  $F^{(k-2)}(\rho, \rho')$  is easily proved to be congruent to zero  $\pmod{p^{\alpha+\beta}}$ . So we may assume  $\left(\frac{K}{p}\right) = 1$ ,  $p = \mathfrak{p} \cdot \mathfrak{p}'$  with two prime ideals in  $K$ . If the conductor of  $\mathfrak{o}$  is divisible by  $p$ ,  $F^{(k-2)}(\rho, \rho')$  is congruent to zero  $\pmod{p^{\alpha+\beta}}$ . Hence we may assume the conductor of  $\mathfrak{o}$  is not divisible by  $p$ . Put  $\mathfrak{p}_0 = \mathfrak{p} \cap \mathfrak{o}$  and  $\mathfrak{p}'_0 = \mathfrak{p}' \cap \mathfrak{o}$ . Let  $d$  be the smallest positive integer such that  $\mathfrak{p}_0^d$  is principal. Put  $\gamma = \operatorname{ord}_p d$ . We may put  $\mathfrak{p}_0^d = \pi \mathfrak{o}$  with  $\pi \in \mathfrak{o}$ , or what is the same as  $\mathfrak{p}^d = \pi \mathfrak{o}_1$ ,  $\mathfrak{o}_1$  being the maximal order of  $K$ . If  $\rho$  is not primitive,  $F^{(k-2)}(\rho, \rho')$  is congruent to zero  $\pmod{p^{\alpha+\beta}}$ . So we may also assume that

$\rho$  is primitive and that  $\rho' \equiv 0 \pmod{\mathfrak{p}}$ . Since  $\rho \cdot \rho' = p^m$ , we have  $(\rho) = \mathfrak{p}'^m$ . Hence  $\mathfrak{p}'^m p^{\gamma-\beta}$  is principal and it is proved that there exists an imaginary quadratic integer  $\rho_1$  such that  $\rho_1^{p^{\beta-\gamma}} = \rho$ . Therefore we have

$$\begin{aligned} F^{(k'-2)}(\rho, \rho') &\equiv \frac{1}{\rho - \rho'} \cdot \rho^{k-1} \cdot \rho_1^{(k'-k)p^{\beta-\gamma}} \pmod{\mathfrak{p}^{\alpha+\beta-\gamma}}, \\ &\equiv \frac{\rho^{k-1}}{\rho - \rho'} \pmod{\mathfrak{p}^{\alpha+\beta-\gamma}}, \\ &\equiv F^{(k-2)}(\rho, \rho') \pmod{\mathfrak{p}^{\alpha+\beta-\gamma}}. \end{aligned}$$

Since  $h_o$  is divisible by  $d$ , we have  $\text{ord}_p h_o \geq \gamma$ . Hence we have

$$\frac{h_o}{w_o} F^{(k'-2)}(\rho, \rho') \equiv \frac{h_o}{w_o} F^{(k-2)}(\rho, \rho') \pmod{p^{\alpha+\beta}}.$$

Thus Proposition is completely proved.

Q.E.D.

In cases of  $p = 2, 3$ , we can prove following propositions by the same arguments as above:

**PROPOSITION.** (Case of  $p = 2$ .) *Let  $m$  and  $\alpha$  be positive integers. Put  $\text{ord}_2 m = \beta$ . Let  $k'$  and  $k$  be even positive integers satisfying (1)  $k' \equiv k \pmod{2^\alpha}$  and (2)  $k' > k \geq \text{Max}\{2\alpha + 6, \alpha + \beta + 4\}$ . Then we have*

$$\text{tr } T_{k'}(2^m) \equiv \text{tr } T_k(2^m) \pmod{2^{\alpha+\beta}}.$$

**PROPOSITION.** (case of  $p = 3$ .) *Let  $m$  and  $\alpha$  be positive integers. Put  $\text{ord}_3 m = \beta$ . Let  $k'$  and  $k$  be even positive integers satisfying (1)  $k' \equiv k \pmod{3^\alpha - 3^{\alpha-1}}$  and (2)  $k' > k \geq \text{Max}\{2\alpha + 4, \alpha + \beta + 3\}$ . Then we have*

$$\text{tr } T_{k'}(3^m) \equiv \text{tr } T_k(3^m) \pmod{3^{\alpha+\beta}}.$$

## § 2. Preliminary lemmas

Let  $x_1, \dots, x_N$  be indeterminates. For each positive integer  $n$ , we define  $S_n(x_1, \dots, x_N) = \sum_{i=1}^N x_i^n$  and  $F_n(x_1, \dots, x_N) = (-1)^n \sum_{1 \leq i_1 < \dots < i_n \leq N} x_{i_1} \dots x_{i_n}$ . We simply write  $S_n$  and  $F_n$  instead of  $S_n(x_1, \dots, x_N)$  and  $F_n(x_1, \dots, x_N)$ . It is obvious that  $F_n = 0$  if  $n$  is greater than  $N$ . It is well known that there exist following relations between two functions  $S_n$  and  $F_n$ , which are called Newton's formulae;

$$S_n + S_{n-1}F_1 + \dots + S_1F_{n-1} + nF_n = 0.$$

By means of Newton's formulae,  $F_n$  (resp.  $S_n$ ) can be described as a polynomial of  $S_i$  (resp.  $F_i$ ) with  $1 \leq i \leq n$  as follows:

$$F_n = \sum_{r=1}^n \sum_{\substack{1 \leq i_1 < \dots < i_r \leq n \\ 1 \leq j_s \\ \sum_{s=1}^r i_s j_s = n}} a_{\binom{i_1, \dots, i_r}{j_1, \dots, j_r}}^{(n)} S_{i_1}^{j_1} \cdots S_{i_r}^{j_r},$$

$$S_n = \sum_{r=1}^n \sum_{\substack{1 \leq i_1 < \dots < i_r \leq n \\ 1 \leq j_s \\ \sum_{s=1}^r i_s j_s = n}} b_{\binom{i_1, \dots, i_r}{j_1, \dots, j_r}}^{(n)} F_{i_1}^{j_1} \cdots F_{i_r}^{j_r},$$

where  $a^{(n)}$  and  $b^{(n)}$  are rational numbers. All these coefficients can be calculated as follows:

LEMMA 1. *We have*

$$(2) \quad a_{\binom{i_1, \dots, i_r}{j_1, \dots, j_r}}^{(n)} = \left( (-1)^{\sum_{s=1}^r j_s} \prod_{s=1}^r j_s! i_s^{j_s} \right)^{-1},$$

and

$$(3) \quad b_{\binom{i_1, \dots, i_r}{j_1, \dots, j_r}}^{(n)} = (-1)^{\sum_{s=1}^r j_s} \frac{\left( \left( \sum_{s=1}^r j_s \right) - 1 \right)!}{\prod_{s=1}^r j_s!} n.$$

*Proof.* We use induction on  $n$ . It is obvious that (2) is valid for  $n = 1$ . Suppose that (2) is valid for all  $a^{(\ell)}$  with  $1 \leq \ell \leq n - 1$ . By Newton's formulae, we have  $F_n = -\frac{1}{n} \left( S_n + \sum_{k=1}^{n-1} S_{n-k} F_k \right)$ . If  $i_1 = n$ , (2) is obviously valid. So we may assume  $i_1 < n$ . Then we have

$$\begin{aligned} a_{\binom{i_1, \dots, i_r}{j_1, \dots, j_r}}^{(n)} &= -\frac{1}{n} \left( (-1)^{\left( \sum_{s=1}^r j_s \right)} \left[ \sum_{s=1}^r \left\{ (j_s - 1)! i_s^{j_s-1} \prod_{k \neq s} j_k! i_k^{j_k} \right\}^{-1} \right] \right), \\ &= (-1)^{\sum_{s=1}^r j_s} \left( \prod_{s=1}^r j_s! i_s^{j_s} \right)^{-1} \frac{1}{n} \sum_{s=1}^r i_s j_s, \\ &= (-1)^{\sum_{s=1}^r j_s} \left( \sum_{s=1}^r j_s! i_s^{j_s} \right)^{-1}. \end{aligned}$$

Hence (2) is proved to be valid. Let us prove that (3) is valid. We also use induction on  $n$ . It is obvious that (3) is valid for  $n = 1$ . Suppose that (3) is valid for all  $b^{(\ell)}$  with  $1 \leq \ell \leq n - 1$ . By Newton's

formulae, we have  $S_n = -\left(nF_n + \sum_{k=1}^{n-1} S_{n-k}F_k\right)$ . If  $i_1 = n$ , it is obvious that (3) is valid. So we may assume  $i_1 < n$ . Then we have

$$\begin{aligned} b_{\binom{i_1, \dots, i_r}{j_1, \dots, j_r}}^{(n)} &= -\sum_{s=1}^r (-1)^{\binom{\sum_{s=1}^r j_s}{s}-1} \frac{\left(\left(\sum_{s=1}^r j_s\right) - 2\right)!}{(j_s - 1)! \prod_{k \neq s} j_k!} (n - i_s), \\ &= (-1)^{\sum_{s=1}^r j_s} \frac{\left(\left(\sum_{s=1}^r j_s\right) - 2\right)!}{\prod_{s=1}^r j_s!} \left(\sum_{s=1}^r j_s n - j_s i_s\right), \\ &= (-1)^{\sum_{s=1}^r j_s} \frac{\left(\left(\sum_{s=1}^r j_s\right) - 1\right)!}{\sum_{s=1}^r j_s!} n. \end{aligned}$$

Therefore (3) is proved to be valid.

Q.E.D.

By making use of Lemma 1, we can prove the following lemma:

LEMMA 2. Let  $G(X) = \prod_{i=1}^k (1 - a_i X)$  and  $H(X) = \prod_{j=1}^{\ell} (1 - b_j X)$  be polynomials with coefficients of rational integers. Put  $s_n = S_n(a_1, \dots, a_k)$ ,  $t_n = S_n(b_1, \dots, b_{\ell})$ ,  $\sigma_n = F_n(a_1, \dots, a_k)$  and  $\tau_n = F_n(b_1, \dots, b_{\ell})$ . Let  $\alpha$  be a positive integer. Then the following statements are equivalent:

- (1)  $s_n \equiv t_n \pmod{p^{\alpha + \text{ord}_p n}}$  for every  $n \geq 1$ ,
- (2)  $\sigma_n \equiv \tau_n \pmod{p^{\alpha}}$  for every  $n$  with  $1 \leq n \leq \text{Max}\{k, \ell\}$ ,
- (3)  $F(X) \equiv G(X) \pmod{p^{\alpha} \mathbf{Z}[X]}$ .

*Proof.* It is obvious that the statements (2) and (3) are equivalent. So we shall show that the statements (1) and (2) are equivalent. Let  $N$  be any positive integer. We assume that (1) $_{N-1}$ :  $s_n \equiv t_n \pmod{p^{\alpha + \text{ord}_p n}}$  for every  $n \leq N - 1$  and (2) $_{N-1}$ :  $\sigma_n \equiv \tau_n \pmod{p^{\alpha}}$  for every  $n \leq N - 1$ . Under this assumption, we show that the following statements are equivalent:

- (1) $_N$   $s_n \equiv t_n \pmod{p^{\alpha + \text{ord}_p n}}$  for every  $n \leq N$ ,
- (2) $_N$   $\sigma_n \equiv \tau_n \pmod{p^{\alpha}}$  for every  $n \leq N$ .

By making use of (3) in Lemma 1, we have

$$s_N = -N\sigma_N + \sum_{r=1}^N \sum_{\substack{1 \leq i_1 < \dots < i_r \leq N \\ 1 \leq j_s}} b_{\binom{i_1, \dots, i_r}{j_1, \dots, j_r}}^{(N)} \sigma_{i_1}^{j_1} \cdots \sigma_{i_r}^{j_r},$$

$$t_N = -N\tau_N + \sum_{r=1}^N \sum_{\substack{1 \leq i_1 < \dots < i_r \leq N \\ 1 \leq j_s \\ \sum_{s=1}^r i_s j_s = N}} b_{(j_1, \dots, j_r)}^{(i_1, \dots, i_r)} \tau_{i_1}^{j_1} \cdots \tau_{i_r}^{j_r}.$$

Since  $\frac{\left(\sum_{s=1}^r j_s\right)!}{\prod_{s=1}^r j_s!}$  is a rational integer,  $\frac{j_s}{N} b_{(j_1, \dots, j_r)}^{(i_1, \dots, i_r)}$  and  $\frac{\left(\sum_{s=1}^r j_s\right)}{N} b_{(j_1, \dots, j_r)}^{(i_1, \dots, i_r)}$

are rational integers. Put  $\beta = \text{ord}_p N$  and  $\gamma = \text{Min} \left\{ \text{ord}_p j_1, \dots, \text{ord}_p j_s, \text{ord}_p \sum_{s=1}^r j_s \right\}$ . Then we have  $\text{ord}_p b_{(j_1, \dots, j_r)}^{(i_1, \dots, i_r)} \geq \beta - \gamma$ . By the condition  $(2)_{N-1}$ , we have  $\sigma_{i_s} \equiv \tau_{i_s} \pmod{p^\alpha}$  for every  $i_s$  with  $1 \leq i_s \leq N-1$ . Hence we have  $\sigma_{i_s}^{j_s} \equiv \tau_{i_s}^{j_s} \pmod{p^{\alpha + \text{ord}_p j_s}}$  for every  $i_s$  with  $1 \leq i_s \leq N-1$ . Therefore we have  $s_N - N\sigma_N \equiv t_N - N\tau_N \pmod{p^{\alpha + \text{ord}_p N}}$ , so  $s_N - t_N \equiv N(\sigma_N - \tau_N) \pmod{p^{\alpha + \text{ord}_p N}}$ . From this, it follows immediately that  $(1)_N$  and  $(2)_N$  are equivalent under the assumption that  $(1)_{N-1}$  and  $(2)_{N-1}$  are valid. Hence it is proved that (1) and (2) are equivalent. Q.E.D.

### § 3. Congruences between Hecke polynomials

For any even positive integer  $k$ , we put  $C_k(X) = \det(I - T_k(p)X)$  and  $H_k(X) = \det(I - T_k(p)X + p^{k-1}X^2I)$  where  $I$  is the identity operator on  $\mathfrak{S}_k$ .  $C_k(X)$  and  $H_k(X)$  are polynomials with coefficients of rational integers.  $H_k(X)$  is usually called the Hecke polynomial.

Combining results in § 1 and 2, we can prove the following:

**THEOREM 1.** *We assume  $p \geq 5$ . Let  $\alpha$  be a positive integer. Let  $k$  be an even positive integer such that (1)  $k \geq 2\alpha + 2$  and (2)  $\dim_{\mathcal{C}} \mathfrak{S}_{k+p^\alpha-p^{\alpha-1}} < p^{k-\alpha-1}$ . Then we have*

$$\begin{aligned} H_{k'}(X) &\equiv H_k(X) \pmod{p^\alpha \mathcal{Z}[X]}, \\ C_{k'}(X) &\equiv C_k(X) \pmod{p^\alpha \mathcal{Z}[X]}, \end{aligned}$$

for every even positive integer  $k' > k$  satisfying  $k' \equiv k \pmod{p^\alpha - p^{\alpha-1}}$ .

*Proof.* Since  $k \geq 2\alpha + 2$ , we have  $H_k(X) \equiv C_k(X) \pmod{p^\alpha \mathcal{Z}[X]}$ . So we shall prove only  $C_{k'}(X) \equiv C_k(X) \pmod{p^\alpha \mathcal{Z}[X]}$ . By the dimension formula for  $\mathfrak{S}_k$ , it is easily proved that  $k + p^\alpha - p^{\alpha-1}$  also satisfies the condition (2) if  $k$  satisfies it. Hence we may prove our theorem only in case of  $k' = k + p^\alpha - p^{\alpha-1}$ . Let  $m$  be any positive integer such that

$m < \dim_C \mathfrak{S}_{k'}$ , and put  $\beta = \text{ord}_p m$ . By the condition (2), we have  $\beta < k - \alpha - 1$ , so we have  $\alpha + \beta + 2 \leq k$ . Hence, making use of Proposition 1, we have  $\text{tr } T_{k'}(p^m) \equiv \text{tr } T_k(p^m) \pmod{p^{\alpha+\beta}}$ . On the other hand, by the recursion formula for  $T_k(p^m)$ , we have  $\text{tr } T_k(p^m) \equiv \text{tr } T_k(p)^m \pmod{p^{k-1}}$ . Therefore we have  $\text{tr } T_{k'}(p)^m \equiv \text{tr } T_k(p)^m \pmod{p^{\alpha+\beta}}$ . Combining these congruences with Lemma 2, we obtain the proof of Theorem 1.

Q.E.D.

In cases of  $p = 2, 3$ , we can prove following theorems by the same arguments as above:

**THEOREM 1** (Case of  $p = 2$ ). *Let  $\alpha$  be a positive integer. Let  $k$  be an even positive integer such that  $k \geq 2\alpha + 6$  and  $\dim_C \mathfrak{S}_{k+2\alpha} < 2^{k-\alpha-3}$ . Then we have*

$$H_{k'}(X) \equiv H_k(X) \pmod{2^\alpha \mathbf{Z}[X]},$$

for every even positive integer  $k' > k$  satisfying  $k' \equiv k \pmod{2^\alpha}$ .

**THEOREM 1** (Case of  $p = 3$ ). *Let  $\alpha$  be a positive integer. Let  $k$  be an even positive integer such that  $k \geq 2\alpha + 4$  and  $\dim_C \mathfrak{S}_{k+3\alpha-3\alpha-1} < 3^{k-\alpha-2}$ . Then we have*

$$H_{k'}(X) \equiv H_k(X) \pmod{3^\alpha \mathbf{Z}[X]},$$

for every even positive integer  $k' > k$  satisfying  $k' \equiv k \pmod{3^\alpha - 3^{\alpha-1}}$ .

We give an application of Theorem 1. In the rest of this section, we assume  $p \geq 5$  for the sake of simplicity. Let  $k' > k$  be even positive integers such that  $k' \equiv k \pmod{p-1}$  and  $k \geq 4$ . Then, it is obvious that  $k$  satisfies the condition (2) in Theorem 1 for  $\alpha = 1$ . Put  $n = \dim_C \mathfrak{S}_k$  and  $n' = \dim_C \mathfrak{S}_{k'}$ . It is clear that  $\det(XI - T_k(p)) = X^n \det\left(I - \frac{1}{X} T_k(p)\right)$ , where  $I$  is the identity operator on  $\mathfrak{S}_k$ . Therefore, from Theorem 1 follows

**COROLLARY.** *Under the above conditions, we have*

$$\det(XI_{k'} - T_{k'}(p)) \equiv X^{n'-n} \det(XI_k - T_k(p)) \pmod{p\mathbf{Z}[X]}.$$

This result is equivalent to Serre's result [3, (i), Corollary to Theorem 6].

#### §4. $p$ -adic Hecke polynomials

Let  $\alpha$  be a positive integer. Put  $X_\alpha = \mathbf{Z}/(p^\alpha - p^{\alpha-1})\mathbf{Z}$  if  $p \neq 2$ , and  $X_\alpha = \mathbf{Z}/2^{\alpha-2}\mathbf{Z}$  if  $p = 2$ .  $\{X_\alpha\}$  forms a projective system naturally. We have

$$X = \lim_{\leftarrow} X_\alpha = \begin{cases} \mathbf{Z}_p \times \mathbf{Z}/(p-1)\mathbf{Z} & \text{if } p \neq 2, \\ \mathbf{Z}_2 & \text{if } p = 2, \end{cases}$$

where  $\mathbf{Z}_p$  is the ring of  $p$ -adic integers. The canonical homomorphism  $\mathbf{Z} \rightarrow X$  is injective. We identify  $\mathbf{Z}$  with a dense subgroup of  $X$  through this homomorphism.

Let  $\mathcal{O}$  denote the ring of formal power series in  $X$  with coefficients in  $\mathbf{Z}_p$ . Let  $\mathfrak{m}$  be the maximal ideal of  $\mathcal{O}$ . The powers of  $\mathfrak{m}, \mathfrak{m}^n, n \geq 0$  define the  $\mathfrak{m}$ -adic topology on  $\mathcal{O}$ .

We assume  $p \geq 5$ . Let  $\{k_\alpha\}_{\alpha=1}^\infty$  be a sequence of monotonically increasing, even positive integers satisfying  $k_\alpha \equiv k_{\alpha'} \pmod{p^\alpha - p^{\alpha-1}}$  if  $\alpha' > \alpha$ ,  $k_\alpha \geq 2^\alpha + 2$  and  $\dim_{\mathcal{O}} \mathcal{S}_{k_\alpha + p^\alpha - p^{\alpha-1}} < p^{k_\alpha - \alpha - 1}$ . Then  $\{k_\alpha\}_{\alpha=1}^\infty$  has a limit in  $X$ , which is denoted by  $\tilde{k}$ . By means of Theorem 1, there exists a common  $\mathfrak{m}$ -adic limit of  $\{H_{k_\alpha}(X)\}$  and of  $\{C_{k_\alpha}(X)\}$  in  $\mathcal{O}$ . Put  $\tilde{H}_{\tilde{k}}(X) = \lim_{\alpha \rightarrow \infty} H_{k_\alpha}(X)$ . It is clear that  $\tilde{H}_{\tilde{k}}(X)$  depends only on  $\tilde{k}$ , but not on the choice of sequences  $\{k_\alpha\}$  with  $\lim k_\alpha = \tilde{k}$ . We call  $\tilde{H}_{\tilde{k}}(X)$  the  $p$ -adic Hecke polynomial.

In the case where  $\tilde{k}$  belongs to  $2\mathbf{Z}$ , we shall show that  $\tilde{H}_{\tilde{k}}(X)$  coincides with the Fredholm determinant of the  $p$ -adic Hecke operator  $\tilde{U}_{\tilde{k}}(p)$  and that  $\tilde{H}_{\tilde{k}}(X)$  is an entire function.

Before this, we extend Lemma 1 as follows:

LEMMA 3. Let  $G(X) = 1 + \sum_{n \geq 1} \sigma_n X^n$  be a formal power series in  $X$  with coefficients  $\sigma_n$  in a field  $K$ , so that  $\log G(X) = \sum_{n \geq 1} (-1)^n \frac{(G(X) - 1)^n}{n}$  is also a formal power series in  $X$  with coefficients in  $K$ , which we write  $-\sum_{n \geq 1} \frac{s_n}{n} X^n$ , with  $s_n \in K$ . Then there exist following relations between  $\sigma_n$  and  $s_n$ ;

$$(4) \quad S_n = \sum_{r=1}^n \sum_{\substack{1 \leq i_1 < \dots < i_r \leq n \\ 1 \leq j_s \\ \sum_{s=1}^r i_s j_s = n}} b_{\binom{i_1, \dots, i_r}{j_1, \dots, j_r}}^{(n)} \sigma_{i_1}^{j_1} \cdots \sigma_{i_r}^{j_r},$$

$$\sigma_n = \sum_{r=1}^n \sum_{\substack{1 \leq i_1 < \dots < i_r \leq n \\ \sum_{s=1}^r i_s = n}} a_{(i_1, \dots, i_r)}^{(n)} s_{i_1}^{j_1} \cdots s_{i_r}^{j_r},$$

where  $a^{(n)}$  and  $b^{(n)}$  are the same as in Lemma 1.

*Proof.* If  $G(X)$  is a polynomial in  $X$  with coefficients in  $K$ , (4) is equal to (2) and (3) in Lemma 1. Put  $G_n(X) = 1 + \sum_{i=1}^n \sigma_i X^i$  and  $\log G_n(X) = (-1) \sum_{i \geq 1} \frac{s_i^{(n)}}{i} X^i$ . Then it is clear that  $s_i^{(n)} = s_i$  for all  $i$  with  $i \leq n$ . Hence, from Lemma 1, (4) follows immediately. Q.E.D.

Let  $\tilde{k}$  be an even integer and let  $D_{\tilde{k}}^{(p)}(X)$  be the Fredholm determinant of the  $p$ -adic Hecke operator  $\tilde{U}_{\tilde{k}}(p)$  which is defined in [2].

**THEOREM 2.** *We have*

$$\tilde{H}_{\tilde{k}}(X) = D_{\tilde{k}}^{(p)}(X), \quad \text{for } \tilde{k} \in 2\mathbf{Z}.$$

*Proof.* Let  $\{k_\alpha\}$  be a sequence of monotonically increasing, even positive integers satisfying  $k_\alpha \equiv k_{\alpha'} \pmod{p^\alpha - p^{\alpha-1}}$  for every  $\alpha' \geq \alpha$ ,  $k_\alpha \leq 2\alpha + 2$ ,  $\dim_{\mathcal{O}} \mathfrak{S}_{k_\alpha + p^\alpha - p^{\alpha-1}} < p^{k_\alpha - \alpha - 1}$  and  $\lim k_\alpha = \tilde{k}$ . Put  $H_{k_\alpha}(X) = 1 + \sum_{n \geq 1} \sigma_n^{(\alpha)} X^n$  and  $\log H_{k_\alpha}(X) = - \sum_{n \geq 1} \frac{s_n^{(\alpha)}}{n} X^n$  with  $\sigma_n^{(\alpha)}$  and  $s_n^{(\alpha)}$  in  $\mathbf{Z}$ . When  $\alpha \rightarrow \infty$ ,  $\{\sigma_n^{(\alpha)}\}$  and  $\{s_n^{(\alpha)}\}$  have  $p$ -adic limits which we denote by  $\sigma_n$  and  $s_n$  respectively. Then we have  $\tilde{H}_{\tilde{k}}(X) = 1 + \sum_{n \geq 1} \sigma_n X^n$ . Since  $\sigma_n^{(\alpha)}$  and  $s_n^{(\alpha)}$  satisfy the relations (4),  $\sigma_n$  and  $s_n$  also satisfy the relations (4). Hence we have  $\log H_{\tilde{k}}(X) = - \sum_{n \geq 1} \frac{s_n}{n} X^n$ . On the other hand, we have  $s_n^{(\alpha)} = \text{tr } U_{k_\alpha}(p^n)$  by (41) in [1]. Hence, from Theorem 1 in [2], it follows that  $s_n = \text{tr } \tilde{U}_{\tilde{k}}(p)^n$ . Therefore we have  $\tilde{H}_{\tilde{k}}(X) = D_{\tilde{k}}^{(p)}(X)$ . Q.E.D.

Since  $D_{\tilde{k}}^{(p)}(X)$  is a  $p$ -adic entire function, we have the following:

**COROLLARY.**  $\tilde{H}_{\tilde{k}}(X)$  is a  $p$ -adic entire function for  $\tilde{k} \in 2\mathbf{Z}$ .

*Remark.* It is obvious that the  $p$ -adic Hecke polynomials converge for all  $x \in \mathbf{Z}_p$ .

*Remark.* In cases of  $p = 2, 3$ , the same argument as above can be applied.

*Remark.* Recently, Prof. B. Dwork kindly let me know a direct proof of Theorem is obtained from Adolphson's thesis and, at the same time, the condition on  $\dim_C \mathfrak{S}_{k+p^a-p^{a-1}}$  can be discarded.

#### REFERENCES

- [ 1 ] Y. Ihara, Hecke polynomials as congruence  $\zeta$  functions in elliptic modular case, *Ann. of Math.*, **85** (1967), 267-295.
- [ 2 ] M. Koike, On some  $p$ -adic properties of the Eichler-Selberg trace formula, *Nagoya Math. J.*, vol. **56** (1974), 45-52.
- [ 3 ] J.-P. Serre, *Formes modulaires et fonctions zeta  $p$ -adiques*, Modular functions of one variable III, Lecture note in math., Springer, Berlin-Heidelberg-New York, 1973.

*Nagoya University*