

On the asymptotic geometry of abelian-by-cyclic groups

by

BENSON FARB

and

LEE MOSHER

*University of Chicago
Chicago, IL, U.S.A.*

*Rutgers University
Newark, NJ, U.S.A.*

Contents

1. Introduction
 - 1.1. The abelian-by-cyclic group Γ_M
 - 1.2. Statement of results
 - 1.3. Homogeneous spaces
 - 1.4. Outline of proofs
2. Preliminaries
3. Linear algebra
 - 3.1. Jordan forms
 - 3.2. Growth of vectors under a linear transformation
4. The solvable Lie group G_M
5. Dynamics of G_M , Part I: Horizontal-respecting quasi-isometries
 - 5.1. Theorem 5.2 on horizontal-respecting quasi-isometries
 - 5.2. Step 1a: Hyperbolic dynamics and the shadowing lemma
 - 5.3. Step 1b: Foliations rigidity
 - 5.4. Step 2: Time rigidity
 - 5.5. Interlude: The induced boundary map
 - 5.6. Step 3: Reduction to Theorem 5.11 on 1-parameter subgroup rigidity
6. Dynamics of G_M , Part II: 1-parameter subgroup rigidity
7. Quasi-isometries of Γ_M via coarse topology
 - 7.1. A geometric model for Γ_M
 - 7.2. Proof of Proposition 7.1 on induced quasi-isometries of G_M
8. Finding the integers
 - 8.1. The first half of the classification
 - 8.2. Quasi-isometric implies that integral powers have the same absolute Jordan forms
9. Quasi-isometric rigidity

The first author was supported in part by NSF Grant DMS 9704640, by IHES and by the Alfred P. Sloan Foundation. The second author was supported in part by NSF Grant DMS 9504946 and by IHES.