

Unitary derived functor modules with small spectrum

by

T. J. ENRIGHT⁽¹⁾

*University of California, San Diego
La Jolla, CA, U.S.A.*

R. PARTHASARATHY

*Tata Institute of Fundamental Research
Bombay, India*

and

N. R. WALLACH⁽¹⁾

*Rutgers University
New Brunswick, N.J., U.S.A.*

J. A. WOLF⁽¹⁾

*University of California
Berkeley, CA, U.S.A.*

Contents

§ 1. Introduction	105
§ 2. θ -stable parabolic subalgebras	108
§ 3. Splitting criteria and signature	110
§ 4. Signature results when u_c is abelian	114
§ 5. Signature results when u is abelian	116
§ 6. Zuckerman functors	119
§ 7. The main results	122
§ 8. Ladder representations for orthogonal groups $SO(p, q)$ with $p+q$ even	124
§ 9. Ladder representations for $Sp(r, s)$ and special representations for $Sp(n, \mathbb{R})$	125
§ 10. Special multiplicity free unitary representations of $Sp(n, \mathbb{R})$	128
§ 11. The Speh representations for $SL(2n, \mathbb{R})$ and their analogues for $SU^*(2n)$	129
§ 12. An application to unitary highest weight modules	131
§ 13. Coherent continuation of Borel de Siebenthal discrete series representations	132

§ 1. Introduction

For a real semisimple Lie group G , the description of the unitary dual remains an elusive question. One of the difficulties has been the lack of technique for constructing

⁽¹⁾ The first, third and fourth authors have been supported in part by NSF Grants #MCS-8300793, #MCS-7903153 and #MCS-8200235 respectively.