

The transformation T from $P, (\xi, \eta)$, to $P', (\xi', \eta')$, where $\xi' = y(Z_3 + p) - H$, $\eta' = \dot{y}(Z_3 + p)$, is topological in $\bar{\Delta}$. We shall show that there is a fixed point of T in Δ , which then corresponds to the desired periodic Γ . Suppose there is no fixed point of T in Δ . Then a continuous vector, or arrow, $P \rightarrow P'$, exists for all points P of $\bar{\Delta}$. Now the disposition of the arrows at boundary points of Δ is as follows. If P is a B_+ point, TP (considered as a point of \mathfrak{H} at Z_3) corresponds to a Γ' through the + end of (the first) G_1 ; further since Γ is in $S^*(Z_2 + p)$ [Lemma 34], it has arrived at G_1 from an S^* . By Lemma 35 (i) its r.p. is distant $O(\zeta)$ from P_+ . The arrow from such a P points nearly at P_+ . Similarly for B_- points. A boundary point on XY corresponds to a Γ through all the G, G' ; hence its $|\dot{y}(Z_3 + p)| < L_3^* k^{-1} = \eta_0$, by Lemma 34. TP has accordingly $|\eta| < \eta_0$, and the arrow from such a P has a downward component. Similarly one from a boundary point on ZW has an upward one. It follows from these facts, and the continuity of the arrow in $\bar{\Delta}$, alone, that when P describes a simple closed contour whose maximum distance from the boundary of Δ is small, the arrow rotates either through $+2\pi$ or -2π (which it depends on the disposition of the signs on the two continua joining XY, ZW , and the sense of description). This is incompatible with there being no fixed point in Δ .

ERRATA

CORRECTIONS TO THE PAPER: "On non-linear differential equations of the second order. III. The equation $\ddot{y} - k(1 - y^2)\dot{y} + y = b\mu k \cos(\mu t + \alpha)$ for large k , and its generalizations" BY J. E. LITTLEWOOD:

Page 277, line 11 Read $O(A(d)k^{-1})$ for $O(A(d, d')k^{-1})$

286, line 16 should read

$$V' + V = -\left(\frac{v}{3} - 2b\right)k - \int \frac{y}{v} dt, \quad (1)$$

299, Fig. 5 $(V^* + M)'$ should be higher.