

Symplectic nonsqueezing of the Korteweg–de Vries flow

by

JAMES COLLIANDER

*University of Toronto
Toronto, ON, Canada*

MARKUS KEEL

*University of Minnesota
Minneapolis, MN, U.S.A.*

GIGLIOLA STAFFILANI

*Massachusetts Institute of Technology
Cambridge, MA, U.S.A.*

HIDEO TAKAOKA

*Kobe University
Kobe, Japan*

and

TERENCE TAO

*University of California
Los Angeles, CA, U.S.A.*

Contents

1. Introduction	198
1.1. Summary of local and global well-posedness theory	199
1.2. Low-frequency approximation of KdV	200
1.3. Application to symplectic nonsqueezing	203
2. Inverting the Miura transform	211
3. The Fourier restriction spaces Y^s and Z^s	218
4. An improved trilinear estimate	219
4.1. The F_0 (resonant) estimate	220
4.2. The $F_{\neq 0}$ (nonresonant) estimate	221
5. Proof of Theorem 1.3: KdV low frequencies are stable under high-frequency perturbations of data	226
6. Proof of Theorem 1.2: BKdV approximates KdV at low frequencies	231
7. Proof of Theorem 1.5: Symplectic nonsqueezing of KdV	238
8. Proof of Theorem 1.1: $P_{\leq N}$ KdV does not approximate KdV	239

The first author was supported in part by N.S.E.R.C. Grant RGPIN 250233-03 and the Sloan Foundation. The second author was supported in part by N.S.F. Grant DMS 9801558 and the Sloan Foundation. The third author was supported in part by N.S.F. Grant DMS 0100345 and the Sloan Foundation. The fourth author was supported in part by J.S.P.S. Grant No. 13740087. The fifth author was a Clay Prize Fellow and was supported in part by grants from the Packard Foundation.