

CORRECTION

QUANTILE-ADAPTIVE MODEL-FREE VARIABLE SCREENING FOR HIGH-DIMENSIONAL HETEROGENEOUS DATA

Ann. Statist. **41** (2013) 342–369

BY XUMING HE¹, LAN WANG² AND HYOKYOUNG GRACE HONG³

University of Michigan, University of Minnesota and Michigan State University

There was a typo in the expression of the model used in Example 3. The model should be

$$Y = 2(X_1^2 + X_2^2) + \{10^{-1} \exp(X_1 + X_2 + X_{18} + X_{19} + \cdots + X_{30})\} \cdot \varepsilon.$$

X. HE
DEPARTMENT OF STATISTICS
UNIVERSITY OF MICHIGAN
ANN ARBOR, MICHIGAN 48109
USA
E-MAIL: xmhe@umich.edu

L. WANG
SCHOOL OF STATISTICS
UNIVERSITY OF MINNESOTA
MINNEAPOLIS, MINNESOTA 55455
USA
E-MAIL: wangx346@umn.edu

H. G. HONG
DEPARTMENT OF STATISTICS AND PROBABILITY
MICHIGAN STATE UNIVERSITY
EAST LANSING, MICHIGAN 48824
USA
E-MAIL: hhong@stt.msu.edu

Received August 2013.

¹Supported in part by NSF Grant DMS-10-07396, NIH Grant R01GM080503 and National Natural Science Foundation of China Grant 11129101.

²Supported in part by NSF Grant DMS-10-07603.

³Supported in part by a Baruch College Eugene M. Lang Fellowship.