

CORRECTION
**STRONG ORACLE OPTIMALITY OF FOLDED CONCAVE
PENALIZED ESTIMATION**

BY JIANQING FAN, LINGZHOU XUE AND HUI ZOU

Princeton University, Penn State University and University of Minnesota

In this note we include a correction to equation (19) on page 840, which is a step in the proof of Theorem 4 of Fan, Xue and Zou [*Ann. Statist.* **42** (2014) 819–849]. There is no change in the statement of Theorem 4, and the rest of the proof stays unchanged. Equation (19) on page 840 should be corrected in the following way:

We apply the coordinatewise mean-value theorem with respect to each coordinate of β (i.e., β_j) to obtain that

$$\nabla_j \ell_n(\widehat{\beta}^{\text{oracle}}) = \nabla_j \ell_n(\beta^*) + [\nabla^2 \ell_n(\beta^*)]_j \cdot \widehat{\Delta} + R_j(\widetilde{\Delta}^{(j)}),$$

where $[\nabla^2 \ell_n(\beta^*)]_j$ denotes the j th row of $\nabla^2 \ell_n(\beta^*)$, $\widetilde{\beta}^{(j)}$ is on the line segment joining $\widehat{\beta}^{\text{oracle}}$ and β^* and $R_j(\widetilde{\Delta}^{(j)}) = [\nabla^2 \ell_n(\widetilde{\beta}^{(j)}) - \nabla^2 \ell_n(\beta^*)]_j \cdot \widehat{\Delta}$.

Define

$$\mathbf{R}(\widetilde{\Delta}) = (R_1(\widetilde{\Delta}^{(1)}), R_2(\widetilde{\Delta}^{(2)}), \dots, R_p(\widetilde{\Delta}^{(p)}))',$$

and rewrite $\mathbf{R}(\widetilde{\Delta})$ as $(\mathbf{R}'_{\mathcal{A}}(\widetilde{\Delta}), \mathbf{R}'_{\mathcal{A}^c}(\widetilde{\Delta}))'$.

The rest of the proof stays the same.

REFERENCES

FAN, J., XUE, L. and ZOU, H. (2014). Strong oracle optimality of folded concave penalized estimation. *Ann. Statist.* **42** 819–849. [MR3210988](#)

J. FAN
DEPARTMENT OF OPERATIONS RESEARCH
AND FINANCIAL ENGINEERING
PRINCETON UNIVERSITY
PRINCETON, NEW JERSEY 08544
USA
E-MAIL: jqfan@princeton.edu

L. XUE
DEPARTMENT OF STATISTICS
PENNSYLVANIA STATE UNIVERSITY
UNIVERSITY PARK, PENNSYLVANIA 16802
USA
E-MAIL: lzxue@psu.edu

H. ZOU
SCHOOL OF STATISTICS
UNIVERSITY OF MINNESOTA
MINNEAPOLIS, MINNESOTA 55414
USA
E-MAIL: zouxx019@umn.edu