

A GENERALIZATION OF THE WEAK VERSION OF THE GLIVENKO-CANTELLI THEOREM

BY JONATHAN SHUSTER

University of Florida

0. Introduction. In this note, a theorem, generalizing the weak version of the Glivenko-Cantelli Theorem, is presented. It is furthermore pointed out that the result is of interest in the area of Goodness of Fit Tests of the Seshadri, Csörgö, Stephens type, [1].

1. The Theorem. Let $X_1, X_2, \dots, X_n, \dots$ be a sequence of positive random variables satisfying the Strong Law of Large Numbers. Let $S_n = X_1 + \dots + X_n$ and

$$T_{j,n} = S_j/S_n, \quad 1 \leq j < n < \infty.$$

Let $\mu > 0$ be the a.e. limit of S_n/n .

Then: for every $\varepsilon > 0$,

$$(1.1) \quad P[\lim_{N \rightarrow \infty} \sup_{n > N} (\max_{1 \leq j < n} |T_{j,n} - (j/n)|) > \varepsilon] = 0.$$

PROOF. Let ε, η be arbitrary positive numbers with $\varepsilon < 4$ and $\eta < 1$. By Egoroff's Theorem, [2], page 339, there exists a set A , of measure at least $(1 - \eta)$, such that $S_n/n \rightarrow \mu$, uniformly on A . We now restrict our attention to the set A .

Set $\varepsilon' = \varepsilon\mu/8$. Then there exists a positive integer $M = M(\varepsilon, \eta)$, such that whenever $n > j \geq M$, $\mu - \varepsilon' < S_j/j < \mu + \varepsilon'$ and $\mu - \varepsilon' < S_n/n < \mu + \varepsilon'$, on A . The above easily gives:

$$|T_{j,n} - (j/n)| < \varepsilon/2 \quad \text{on } A, \text{ for } n > j \geq M.$$

Next, choose $N_0 = N_0(\varepsilon, M)$, such that

$$M/N_0 < \varepsilon/2.$$

Let $k < M$, $n \geq N_1 \geq N_0$.

$$\begin{aligned} |T_{k,n} - (k/n)| &\leq \max [T_{M,N_0}, (M/N_0)] \\ &\leq |T_{M,N_0} - (M/N_0)| + \varepsilon/2 \\ &< \varepsilon \quad \text{on } A, \text{ for all } k < M. \end{aligned}$$

Hence:

$$P[\sup_{n \geq N_0} (\max_{1 \leq j < n} |T_{j,n} - (j/n)|) > \varepsilon] < \eta.$$

Since N_0 is a function of ε, η , the arbitrary constants, the proof is complete.

2. Specialization. Let the X_i be independent, identically distributed positive random variables with mean, μ . Under these conditions, a necessary and sufficient condition that the random variables $T_{j,n}$ behave as $(n - 1)$ order statistics from the uniform pdf on $(0, 1)$, is that the X_i have pdf $f(x, \lambda) = \lambda \exp(-\lambda x)$. This result is proved in [1].

Received March 10, 1970; revised November 24, 1971.