

A SHORT PROOF OF A KNOWN LIMIT THEOREM FOR SUM  
OF INDEPENDENT RANDOM VARIABLES WITH INFINITE  
EXPECTATIONS<sup>1</sup>

BY BERT FRISTEDT

University of Minnesota

The following theorem is proved by Feller ([1]) with slightly more general hypotheses. He proves it using Kronecker's theorem and a special case of the three series theorem. We shall prove it using an elementary application of the law of large numbers.

**THEOREM.** Let  $X_1, X_2, \dots$  be a sequence of independent, identically distributed random variables with common distribution function  $V$ . Let  $S_n = X_1 + \dots + X_n$ . Let  $0 = a_0 < a_1 < \dots$  be a convex sequence of numbers. Assume that  $\int |x| dV(x) = \infty$ . Then,

$$P\{|S_n| > a_n \text{ infinitely often}\} = 0 \text{ or } 1$$

according as

$$\sum_{n=1}^{\infty} \int_{|x| > a_n} dV(x) < \infty \text{ or } = \infty.$$

**PROOF.** Assume first that

$$\sum_{n=1}^{\infty} \int_{|x| > a_n} dV(x) = \infty.$$

Since  $2a_n \leq a_{2n}$  (which follows from the convexity of  $\{a_n\}$ ), we conclude that

$$\sum_{n=1}^{\infty} \int_{|x| > 2a_n} dV(x) = \infty.$$

Hence

$$P\{|X_n| > 2a_n \text{ infinitely often}\} = 1$$

which implies the desired conclusion.

For the other half of the proof we can, and do, assume, with no loss of generality, that  $X_n \geq 0$  for all  $n$ . Of course, we assume that

$$\sum_{n=1}^{\infty} \int_{a_n}^{\infty} dV(x) < \infty.$$

For fixed  $k$  we define a new sequence:

$$\begin{aligned} b_n &= nk^{-1}a_k, \quad n = 0, 1, \dots, k; \\ b_n &= a_n, \quad n = k + 1, \dots \end{aligned}$$

The sequence  $0 = b_0 < b_1 < \dots$  is convex. Let  $b(x)$  be defined for all  $x \geq 0$  such that  $b$  is strictly increasing and convex, and such that  $b(n) = b_n$  if  $n$  is a non-negative integer.

Received 12 November 1965.

<sup>1</sup> Supported in part by NSF Grant GP-7490.