

ON OBTAINING BALANCED INCOMPLETE BLOCK DESIGNS FROM PARTIALLY BALANCED ASSOCIATION SCHEMES

BY PETER W. M. JOHN

University of California, Davis

1. Shrikhande and Singh [6] pointed out that if there exists a PBIB design with two associate classes for v varieties such that $p_{i,i}^1 = p_{i,i}^2 = \lambda$ ($i = 1$ or 2), then we can construct a symmetric BIB design for v varieties with parameter $v, r = n_i, \lambda$. The design is obtained by letting the j th block consist of all those varieties which are i th associates of the j th variety.

Their procedure can be extended to PBIB designs with m associate classes, $m > 2$, with the requirement that for some $i, p_{i,i}^k = \lambda$ for all k , where $k = 1, \dots, m$. Consider the following association scheme for $(s+2)(s+3)(s+4)/6$ varieties. Denote each variety by one of the triplets (x, y, z) where x, y, z are integers such that $1 \leq x < y < z \leq s+4$.

Two varieties are i th associates if their representations have $(3-i)$ integers in common. This is an association scheme of a class of PBIB designs of three associate classes, John [3], with $p_{22}^1 = s^2, p_{22}^2 = (s-1)(s+6)/2, p_{22}^3 = 9(s-2)$. For $s = 3$ we have $v = 35, p_{22}^1 = p_{22}^2 = p_{22}^3 = 9$. Thus we obtain a BIBD $b = v = 35, r = k = 18, \lambda = 9$ by assigning one block to each variety and letting the j th block contain all the second associates of the j th variety. For example, the block corresponding to variety $(1, 2, 3)$ consists of the eighteen varieties represented by $(1, y, z), (2, y, z)$ or $(3, y, z)$ where $y, z = 4, 5, 6$ or 7 and $y < z$. (The existence of a design for $v = 35, k = 18, \lambda = 9$ is well known. A difference set for the complementary design $(35, 17, 8)$ is given by Hall [2]).

2. Other symmetric PBIB designs are obtained by assigning to the j th block the j th variety and its i th associates. For these designs $\lambda_i = p_{i,i}^1 + 2$, and $\lambda_k = p_{i,i}^k (k \neq i), k = n_i + 1$ and BIB designs are obtained when we can find association schemes with $\lambda_i, \lambda_k = \lambda$ for all k . This procedure produces nothing new from PBIB schemes with two associate classes because the design of this type for $i = 1$ is merely the complement of the design obtained by the procedure of the previous section for $i = 2$, and vice versa. However, for $m > 2$ new designs occur.

The cubic designs of Raghavarao and Chandrasekhararao [4] have $v = s^3, p_{22}^1 = 2(s-1)(s-2), p_{22}^2 = 2(s-1) + (s-2)^2, p_{22}^3 = 6(s-2), n_2 = 3(s-1)^2$. Each variety is represented by a set of three integers $(x, y, z), 1 \leq x, y, z \leq s$. The integers in any representation need not be all different and any two varieties are i th associates if their representations have exactly $(3-i)$ coordinates equal. With $s = 4$ we have $p_{22}^1 = p_{22}^2 + 2 = p_{22}^3 = 12, v = 64, n_2 + 1 = 28$. This gives a symmetric BIBD with $v = 64, k = 28, \lambda = 12$. The j th block consists of v_j and all its second associates. The block corresponding to variety $(1, 2, 3)$ consists of

Received 23 June 1966; revised 15 August 1966.