REFERENCES

[1] R. C. Bose and K. R. Nair, "Partially balanced incomplete block designs," *Sankya*, Vol. 4 (1939), pp. 337–372.

[2] R. C. Bose and T. Shimamoto, "Classification and analysis of partially balanced designs with two associate classes," *J. Amer. Stat. Assn.*, Vol. 47 (1952), pp. 151–190.

[3] W. S. Connor, "The uniqueness of the triangular association scheme," *Ann. Math. Stat.*, Vol. 29 (1958), pp. 262–266.

[4] A. J. Hoffman, "On the exceptional case of a characterization of the arcs of a complete graph," to appear in *IBM Journal of Research*.

[5] A. J. Hoffman and R. R. Singleton, "On a graph-theoretic problem of E. F Moore," to appear in *IBM Journal of Research*.

[6] S. S. Shrikhande, "On a characterization of the triangular association scheme," *Ann. Math. Stat.*, Vol. 30 (1959), pp. 39–47.

NOTE

The results of this paper have also been obtained, using different methods, by Chang, L. C., "The Uniqueness and Nonuniqueness of the Triangular Association Schemes," *Science Record*, Vol. III, New Series, 1959, pp. 604–613. Chang has also shown that there are exactly three counterexamples when $n = 8$ ("Association Schemes of Partially Balanced Designs with Parameters $v = 28$, $n_1 = 12$, $n_2 = 15$ and $p_{11}^2 = 4$," *Science Record*, Vol. IV, New Series, 1960, pp. 12–18).