

## A NOTE ON SHEPPARD'S CORRECTIONS

BY SOLOMON KULLBACK

In this note we shall derive a simple relation between the characteristic function of the grouped distribution and the characteristic function of the original continuous distribution, assuming that the frequency curve has high contact with the x-axis at both ends.

If we set  $p_s = \int_{x_s - \frac{w}{2}}^{x_s + \frac{w}{2}} f(x) dx$ , then the characteristic function of the grouped distribution is given by

$$(1) \quad \psi(t) = \sum e^{itx_s} p_s$$

where  $i = \sqrt{-1}$ . Replacing  $p_s$  by its value as given above, we have

$$\begin{aligned} (2) \quad \psi(t) &= \sum e^{itx_s} \int_{x_s - \frac{w}{2}}^{x_s + \frac{w}{2}} f(x) dx \\ &= \sum e^{itx_s} \int_{-\frac{w}{2}}^{\frac{w}{2}} f(x + x_s) dx \\ &= \int_{-\frac{w}{2}}^{\frac{w}{2}} dx \sum e^{itx_s} f(x + x_s) \\ &= \sum e^{itx_s} f(x_s) \int_{-\frac{w}{2}}^{\frac{w}{2}} e^{-itx} dx. \end{aligned}$$

There is no difficulty about justifying the inversion of the order of integration and summation.

Because of the assumption of high-contact with the axis of  $x$  at both ends of the frequency curve, we have

$$(3) \quad \varphi(t) = \int e^{itx} f(x) dx = w \sum e^{itx_s} f(x_s)$$

so that

$$(4) \quad \psi(t) = \frac{2}{wt} \sin \frac{tw}{2} \varphi(t).$$