

- BICKEL, P. J. and LEHMANN, E. L. (1975). Descriptive statistics for non-parametric models. II. *Ann. Statist.* **3** 1045–1069.
- BROWNLEE, K. A. (1965). *Statistical Theory and Methodology in Science and Engineering*, 2nd ed. Wiley, New York.
- CARROLL, R. J. (1979). On estimating variances of robust estimators when the errors are asymmetric. *J. Amer. Statist. Assoc.* **74** 674–679.
- DE WET, T. and VENTER, J. H. (1974). An asymptotic representation of trimmed means with applications. *S. African Statist. J.* **8** 127–134.
- HILL, M. A. and DIXON, W. J. (1982). Robustness in real life: a study of clinical laboratory data. *Biometrics* **38** 377–396.
- HUBER, P. J. (1972). Robust statistics: a review. *Ann. Math. Statist.* **43** 1041–1067.
- HUBER, P. J. (1981). *Robust Statistics*. Wiley, New York.
- JAECKEL, L. A. (1971). Some flexible estimates of location. *Ann. Math. Statist.* **42** 1540–1552.
- KOENKER, R. W. and BASSETT, G. W. (1978). Regression quantiles. *Econometrica* **46** 33–50.
- KOUL, H. L. (1969). Asymptotic behavior of Wilcoxon type confidence regions in multiple linear regression. *Ann. Math. Statist.* **40** 1950–1979.
- ROCKE, D. M., DOWNS, G. W. and ROCKE, A. J. (1982). Are robust estimators really necessary? *Technometrics* **24** 95–101.
- RUPPERT, D. and CARROLL, R. J. (1980). Trimmed least squares estimation in the linear model. *J. Amer. Statist. Assoc.* **75** 828–838.
- SHORACK, G. R. (1974). Random means. *Ann. Statist.* **2** 661–675.
- SKOROKHOD, A. V. (1956). Limit theorems for stochastic processes. *Theory Probab. Appl.* **1** 261–290.
- SPJØTVOLL, E. and AASTREIT, A. H. (1980). Comparison of robust estimators on data from field experiments. *Scand. J. Statist.* **7** 1–13.
- STIGLER, S. M. (1973). The asymptotic distribution of the trimmed mean. *Ann. Statist.* **1** 472–477.
- STIGLER, S. M. (1977). Do robust estimators work with real data? *Ann. Statist.* **5** 1055–1077.
- TUKEY, J. W. and McLAUGHLIN, G. (1963). Less vulnerable confidence and significance procedures for location based on a single sample. *Sankhyā Ser. A* **35** 331–352.

DEPARTMENT OF STATISTICS
UNIVERSITY OF CHICAGO
5734 UNIVERSITY AVENUE
CHICAGO, ILLINOIS 60637

DISCUSSION

P. J. DE JONGH AND T. DE WET

Institute for Maritime Technology

This paper proposes a new way of defining trimmed means in the linear model, which differs from earlier proposals by Bickel (1973), Koenker and Bassett (1978) and Ruppert and Carroll (1980). We find the idea of the proposal very interesting. It has the “right” equivariance and asymptotic properties and is thus an attractive (large sample) extension of the trimmed mean in the location case. These properties also hold for the Koenker–Bassett (1978) estimator, but the Welsh estimator has the potential advantage of computational simplicity (if least squares is used as a preliminary estimator). Our remarks will concern the small sample behaviour of the proposed estimator. We wish to

