

ASYMPTOTIC BEHAVIOR OF HIGH ORDER MEANS¹

BY JAMES W. DANIEL

The University of Texas at Austin

The following simple but interesting question was suggested by S. W. Joshi. Given a nonnegative Borel measure μ on $[0, 1]$ with $\mu([0, 1]) = 1$, the mean m_1 can be defined as the unique root in $[0, 1]$ of the equation $\int_0^1 (x-m) d\mu(x) = 0$; the root m_r in $[0, 1]$ of $f_r(m) = \int_0^1 |x-m|^r \text{sign}(x-m) d\mu(x) = 0$, for $r \geq 1$, can be considered a generalized mean, in the spirit of the more general ϕ -means of [1] for which results similar to these being presented here can be obtained. The question arises as to how m_r depends on r and μ ; in this note we show that m_r converges to the midpoint of the interval of essential support of the measure μ as r tends to infinity.

Let $a = \sup \{l; \mu([l, 1]) > 0\} = \mu([0, 1])$ and let $b = \inf \{r; \mu([0, r]) = \mu([0, 1])\}$. Clearly then $\mu([a, b]) = \mu([0, 1])$, $\mu([a, a+\varepsilon]) > 0$ for every ε in $(0, b-a]$, and $\mu([b-\varepsilon, b]) > 0$ for every ε in $(0, b-a]$; we call the interval $[a, b]$ the interval of essential support of the measure μ , although one might discard an endpoint if it itself has μ -measure zero. Our result can now be stated as follows. *For each $r \geq 1$, a unique m_r exists, and $\lim_{r \rightarrow \infty} m_r = m_\infty \equiv (a+b)/2$.*

The proof is simple. Since $f_r(m)$ is a continuous function of m and satisfies $f_r(0) = f_r(a) \geq 0$ and $f_r(1) = f_r(b) \leq 0$, at least one root exists in $[0, 1]$. Furthermore, by using the Lebesgue dominated convergence theorem we can show easily that $f_r(m)$ is differentiable and $f'_r(m) = -r \int_0^1 |x-m|^{r-1} d\mu(x)$ which is less than zero for all m unless $a = b = m$, in which case $m_r = m_\infty$ for all r and there is nothing to prove. Thus a unique root m_r exists and lies in $[a, b]$. Now it remains to prove that m_r tends to m_∞ when $a \neq b$. Let m be a fixed number satisfying $m_\infty < m \leq b$. We shall show that for all large r we have $f_r(m) < 0$ which implies $m_r < m$ which in turn implies $\limsup_{r \rightarrow \infty} m_r \leq m_\infty$ since $m > m_\infty$ was arbitrary; for this purpose let ε satisfy $0 < \varepsilon < 2(m-m_\infty)$, $\varepsilon < m-a$. Then

$$\begin{aligned} f_r(m) &= -\int_a^{a+\varepsilon} (m-x)^r d\mu(x) - \int_{a+\varepsilon}^m (m-x)^r d\mu(x) + \int_m^b (x-m)^r d\mu(x) \\ &\leq -\int_a^{a+\varepsilon} (m-x)^r d\mu(x) + \int_m^b (x-m)^r d\mu(x) \\ &\leq -(m-a-\varepsilon)^r \mu([a, a+\varepsilon]) + (b-m)^r \mu([m, b]) \\ &= (m-a-\varepsilon)^r \mu([a, a+\varepsilon]) \left\{ -1 + \left(\frac{b-m}{m-a-\varepsilon} \right)^r \frac{\mu([m, b])}{\mu([a, a+\varepsilon])} \right\}. \end{aligned}$$

Since $m-a-\varepsilon > 0$, since $\mu([a, a+\varepsilon]) > 0$, and since $(b-m)/(m-a-\varepsilon) < 1$ because $\varepsilon < 2(m-m_\infty) = 2(m-(b+a)/2)$, we conclude that $f_r(m) < 0$ for sufficiently

Received October 12, 1970.

¹ Research supported in part under Contract Number N000-14-67-A-0128-0004 at the University of Wisconsin.