

## ERRATA

H. W. Gould, *A series transformation for finding convolution identities*, vol. 28 (1961), p. 200. The last line should read

$$C_k(a, b) = \frac{a}{a + bk} G_k(a, b).$$

David Dean and Ralph A. Raimi, *Permutations with comparable sets of invariant means*, vol. 27 (1960). In Theorems 3.3 and 4.2 it is necessary to add the hypothesis that  $F_\sigma = F_\mu$ , where  $F_\sigma$  is the collection of finite cycles in  $\sigma$ , as in the definition preceding Lemma 4.3. This error does not affect what follows Theorem 4.2 and is irrelevant to what precedes Theorem 3.3.

Page 468, line 6: The displayed formula should read

$$l_{p_\alpha} x_\alpha = \bigcap_{\alpha \circ \epsilon A} (\bigcup_{\alpha > \alpha_\circ} \{x_\alpha\}).$$

Page 468, line 19: The displayed formula should read

$$M'_\sigma = \{ \bigcup [l_{p_\alpha} S'_\alpha p'_\alpha] \}^{\wedge}.$$

Jack Levine, *Coefficient identities derived from expansions of elementary symmetric function products in terms of power sums*, vol. 28(1961).

In (2.2) read  $\frac{\partial}{\partial s_m}$  instead of  $\frac{\partial}{\partial s_m}$ .

In first line under (2.10) read  $1^{n_i}$  instead of  $1^{n_i}$ .

Page 95, in (6) read "entries" instead of "entires".

In (3.1), (3.5), (3.9), (4.9) read  $\sum_m$  instead of  $\sum_m$ .

In (3.10) read  $\sum_{m-1}$  instead of  $\sum_{m-1}$ .

In (6.5) read  $\sum_m$  on left and  $\sum_{m'_\alpha}$  on right.

Page 102, line 8 from bottom, read  $\sum_3, \sum_4, \sum_5$ , instead of  $\sum_3, \sum_4, \sum_5$ .

Eckford Cohen, *Representations of even functions (mod r)*, III. *Special topics*, vol. 26(1959), pp. 491-500. *Remark.* In §4 of this paper the function  $G_s(n, r)$  was defined to be the number of solutions of  $n \equiv p_1 x_1 + \dots + p_s x_s \pmod{r}$  such that  $(x_i, r) = 1$ ,  $p_i \mid r$ ,  $p_i$  prime ( $i = 1, \dots, s$ ). In §1 this function was interpreted to be the number of representations of  $n$  as a sum of  $s$  primes  $\pi_i$  in the residue class ring  $J_r$  of the integers (mod  $r$ ). Actually,  $G_s(n, r)$  represents the number of *weighted* compositions of  $n$  in  $J_r$  as a sum of  $s$  primes  $\pi_i$  with each  $\pi_i$  counted  $p_i$  or  $p_i - 1$  times ( $p_i$  being the prime divisor associate  $d$  with  $\pi_i$ ), according as  $p_i$  does or does not divide  $r$  to a power higher than the first.

M. Lees and M. H. Protter, *Unique continuation for parabolic differential equations and inequalities*, vol. 28(1961), page 369, line 2,  $L = A - \frac{\partial}{\partial t}$  instead of  $L = A = \frac{\partial}{\partial t}$ .