

## ON COMPLETE INTERSECTION REAL CURVES

M. G. MARINARI AND M. RAIMONDO.

Dedicated to the memory of Gus Eforymson

Let  $C \subset \mathbf{R}^n$  be an embedded smooth real algebraic curve. Let  $\mathbf{R}[C]$  (resp.  $\Gamma_C = \Gamma(C, \mathcal{O}_C)$ ) denote the ring of real polynomial functions (resp. of global regular functions) on  $C$ . If  $I_C$  (resp.  $J_C$ ) is the ideal of  $C$  in  $\mathbf{R}[X_1, \dots, X_n]$  (resp. in  $\Gamma_{\mathbf{R}^n} = \Gamma(\mathbf{R}^n, \mathcal{O}_{\mathbf{R}^n})$ ), then we have  $\Gamma_C \simeq \Gamma_{\mathbf{R}^n}/J_C \simeq N^{-1}\mathbf{R}[C]$  where  $N = \{s \in \mathbf{R}[C] \mid s(x) \neq 0 \ \forall x \in C\}$  and  $\mathbf{R}[C] \simeq \mathbf{R}[X_1, \dots, X_n]/I_C$ .

Let  $\mathcal{C}$  be the abstract curve of which  $C$  is a realization; to each embedding  $\varphi: \mathcal{C} \rightarrow \mathbf{R}^N$  corresponds a f.g.  $\mathbf{R}$ -algebra  $P = \mathbf{R}[\varphi(\mathcal{C})] = \mathbf{R}[X_1, \dots, X_N]/I_{\varphi(\mathcal{C})}$  which is called an affine representation of  $\mathcal{C}$  and has the property that  $\Gamma_{\varphi} \simeq N_P^{-1}P$  where  $N_P = \{s \in P \mid s(x) \neq 0 \text{ for each } x \in \varphi(\mathcal{C})\}$ . The various affine representations of  $\mathcal{C}$  can be compared by introducing on the set of isomorphism classes of affine representations of  $\mathcal{C}$  the following ordering relation  $<$ : "given two affine representations  $P, Q$  of  $\mathcal{C}$  then  $P < Q$  if there exists a homomorphism of  $\mathbf{R}$ -algebras  $P \hookrightarrow Q$  which extends to an isomorphism  $\Gamma_{\varphi} \simeq N_P^{-1}P \simeq N_Q^{-1}Q \simeq \Gamma_{\psi}$ . We assume that all the affine representations that we consider are regular as schemes (this is always possible). We say that  $P$  is "the canonical" affine representation if  $P < Q$ , for every other  $Q$ . Since to each affine representation corresponds a complexification (uniquely, up to isomorphism), we mix up the two notions. So, given two complexifications  $\tilde{C}', \tilde{C}''$ , we say that  $\tilde{C}' < \tilde{C}''$  if there is a real open immersion  $\tilde{C}'' \hookrightarrow \tilde{C}'$ .

In this setting the main result we know is the following (cf. [3]).

**THEOREM 1.** *A smooth affine real curve  $\mathcal{C}$  has a canonical complexification if and only if it is either rational or embeddable as a non-compact subspace of  $\mathbf{R}^3$ .*

The above machinery can be used in order to find some results on the problem of complete intersection. We recall the following definition (cf. [2]): "an integral domain  $R$  which is a f.g. algebra over a field  $k$ , is called an abstract complete intersection (ACI) if there exists a polynomial ring