

INTEGRABILITY OF PLANAR POLYNOMIAL DIFFERENTIAL SYSTEMS THROUGH LINEAR DIFFERENTIAL EQUATIONS

H. GIACOMINI, J. GINÉ AND M. GRAU

ABSTRACT. In this work we consider rational ordinary differential equations $dy/dx = Q(x, y)/P(x, y)$, with $Q(x, y)$ and $P(x, y)$ coprime polynomials with real coefficients. We give a method to construct equations of this type for which a first integral can be expressed from two independent solutions of a second-order homogeneous linear differential equation. This first integral is, in general, given by a non Liouvillian function.

We show that all the known families of quadratic systems with an irreducible invariant algebraic curve of arbitrarily high degree and without a rational first integral, can be constructed by using this method. We also present a new example of this kind of family.

We give an analogous method for constructing rational equations but by means of a linear differential equation of first order.

1. Introduction. This paper deals with rational ordinary differential equations such as

$$(1) \quad \frac{dy}{dx} = \frac{Q(x, y)}{P(x, y)},$$

where $Q(x, y)$ and $P(x, y)$ are coprime polynomials with real coefficients. We associate to this rational equation a planar polynomial differential system by introducing an independent variable t usually called *time*. Denoting by $\dot{} = d/dt$, we have

$$(2) \quad \dot{x} = P(x, y), \quad \dot{y} = Q(x, y),$$

2000 AMS *Mathematics Subject Classification.* Primary 34A05, 34C05, 34C20.
Key words and phrases. Planar polynomial system, first integral, invariant curves, Darboux integrability.

The second and third authors are partially supported by a MCYT grant number BFM 2002-04236-C02-01. The second author is partially supported by DURSI of Government of Catalonia's Acció Integrada ACI2002-24 and by the Distinció de la Generalitat de Catalunya per a la promoció de la recerca universitària.

Received by the editors on May 26, 2003, and in revised form on October 8, 2003.