

SINGULAR LIMITS IN FREE BOUNDARY PROBLEMS

ANDREW STUART

ABSTRACT. We analyze the following class of nonlinear eigenvalue problems: find $(\underline{u}, \mu) \in B \times \mathfrak{R}$ satisfying

$$\begin{aligned} (1) \quad & D\underline{u} + \mu H(\underline{a} \cdot \underline{u} - 1) f(\underline{u}) = 0 \quad \text{in } \Omega \subseteq \mathfrak{R}^N, \\ (2) \quad & \underline{u} = 0 \quad \text{on } \partial\Omega. \end{aligned}$$

Here $H(X)$ is the Heaviside step-function defined by

$$\begin{aligned} H(X) &= 0, & X &\leq 0 \\ H(X) &= 1, & X &> 0. \end{aligned}$$

B is some Banach space appropriate to the problem. D is taken to be a (possibly nonlinear) differential operator with the property that, when $\mu = 0$, equations (1,2) have the unique solution $\underline{u} \equiv 0$.

The problem (1,2) is of free boundary type, since determination of the sets on which $\underline{a} \cdot \underline{u} = 1$ is necessary to determine the solution. Our motivation is the study of porous medium combustion where equations of the form (1,2) represent equilibrium states of the coupled chemical and heat-transfer processes governing the combustion [3, 4]. The step function H arises from the diffusion limited reaction rate, in the limit of large activation energy. The reaction behaves as a switch, triggered when the temperature of the solid phase reaches a threshold value. Problems such as (1,2) also arise in a variety of other applications such as the study of vortex motion in ideal fluids [1] and plasma physics [7].

Clearly $\underline{u} \equiv 0$ satisfies (1,2) for all μ . However, there is no classical bifurcation from this trivial solution, since all other solutions must satisfy $\underline{a} \cdot \underline{u} > 1$ at some point in Ω , and hence cannot be of arbitrarily small supremum norm. Nonetheless, it is useful to develop a constructive approach to the solution of (1,2) since explicit constructions are useful both as the basis for numerical continuation procedures and as the basis for local time-dependent stability calculations.

Received by the editors on May 20, 1989.

Copyright ©1991 Rocky Mountain Mathematics Consortium