

SEPARATING HYPERPLANES FOR CONVEX SETS OVER ORDERED FIELDS

JONATHAN LEE MERZEL

Dedicated to the memory of Gus Efroymson

We work with ordered fields $(F, <)$, the ordering being fixed. For an extension K of F , we may use the same symbol $<$ for an extension of the ordering on F to K . F^n denotes the space of n -tuples of elements of F , whose elements in turn may be designated by vector notation \mathbf{x} . For $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{y} = (y_1, \dots, y_n)$, $\mathbf{x} \cdot \mathbf{y}$ is the "usual" dot product $x_1 y_1 + \dots + x_n y_n$.

DEFINITION. A subset A of F^n is called convex iff for all $\mathbf{a}, \mathbf{b} \in A$ and all $\lambda \in F$: $0 < \lambda < 1$ implies $(1 - \lambda)\mathbf{a} + \lambda\mathbf{b} \in A$.

DEFINITION. A hyperplane in F^n with equation $\mathbf{a} \cdot \mathbf{x} = b$ is said to strongly separate subsets A, B of F^n if either

$$\begin{aligned} \mathbf{a} \cdot \mathbf{y} < b \text{ for all } \mathbf{y} \in A \text{ and } \mathbf{a} \cdot \mathbf{y} > b \text{ for all } \mathbf{y} \in B \text{ or} \\ \mathbf{a} \cdot \mathbf{y} > b \text{ for all } \mathbf{y} \in A \text{ and } \mathbf{a} \cdot \mathbf{y} < b \text{ for all } \mathbf{y} \in B. \end{aligned}$$

(Weaker versions of separation allow points of A, B , or possibly both to lie on the hyperplane.)

The question to be studied is this: given disjoint convex sets A, B in F^n , is there an ordered field extension $(K, <)$ of $(F, <)$ such that A, B are (weakly or strongly) separated by a hyperplane in K^n ? Of course, if K exists it may be taken to have transcendence degree at most n over F (for we need only adjoin to F the coefficients in the equation of the hyperplane, one of which can be taken to be 1).

In fact, we have the following theorem.

THEOREM. *If $(F, <)$ is an ordered field and A, B are disjoint convex sets in F^n , then there is an ordering on $K = F(t_1, \dots, t_n)$ (t_1, \dots, t_n independent transcendentals over F) extending that on F such that A, B are strongly separated by a hyperplane in K^n (with respect to the extended ordering).*

The gist of the argument is to first replace A, B with a maximal disjoint pair of convex sets, and then to identify the intercepts of the separating