

THE MODULI OF WEIERSTRASS FIBRATIONS OVER \mathbf{P}^1 : RATIONALITY

PABLO LEJARRAGA

1. The moduli of Weierstrass fibrations over \mathbf{P}^1 : Rationality.

Let K be an algebraically closed field of characteristic $\neq 2, 3$. A Weierstrass fibration is a pair $(p : Y \rightarrow S, s)$ where p is a morphism of algebraic varieties over K whose fibers are elliptic curves or rational curves with a node or a cusp, and where s is a section of S not passing through the nodes or cusps of the fibers. As is well known, a Weierstrass fibration can be defined by an equation of the form $y^2 = x^3 + ax + b$ due to Weierstrass (see [1]).

In [3], Miranda defines the moduli W_N , $N \geq 1$ of Weierstrass fibrations over $S = \mathbf{P}_K^1$ whose section has self-intersection $-N$ in the associated elliptic surface. Seiler [4] generalizes this work and defines moduli for Weierstrass fibrations over an arbitrary complete nonsingular connected curve. In spite of their obvious interest very little is known about the structure of these moduli. Here we prove:

Theorem 1.1. *When $K = \mathbf{C}$, W_N is a rational variety for all $N \geq 1$.*

Define V_n to be the set of binary homogeneous forms of degree n in t, s . Let $f = f(t, s)$ be an element of V_n and $g = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ be an element of \mathbf{GL}_2 . Then

$$f \cdot g = f(\alpha t + \beta s, \gamma t + \delta s)$$

defines an action of \mathbf{GL}_2 on V_n . There are induced actions of \mathbf{GL}_2 on products $V_m \times V_n$. Define $(V_{4N} \times V_{6N})^s$ and V_{6N}^s to be the open sets of SL_2 -stable (= finite stabilizer and closed orbit) points of $V_{4N} \times V_{6N}$ and V_{6N} , respectively. By means of the Weierstrass equation, W_N is defined as the quotient $(V_{4N} \times V_{6N})^s / \mathbf{GL}_2$. On the other side the quotient

Received by the editors on May 31, 1991.
1980 *Mathematics Subject Classification* (1985 Revision). Primary 14J27, Secondary 14M20.