

A NOTE ON THE STRUCTURE OF THE POWER SET

G. F. SCHUMM

In the theory of sets, one commonly finds existence theorems of the form

*For every infinite set  $S$  of cardinality  $m$  there exists a subset  $T$  of  $\mathcal{P}(S)$  such that  $\text{card}(T) = 2^m$  and  $\mathbf{P}$ ,*

where  $\mathbf{P}$  is some property on the elements of  $T$  and ' $\mathcal{P}(S)$ ' denotes the power set of  $S$ . Thus, for example, it is known that there exist, for every infinite set  $S$  of cardinality  $m$ , subsets  $F, G, H$  and  $L$  of  $\mathcal{P}(S)$  each of cardinality  $2^m$  and such that

- (1)  $\text{card}(X \cap Y) < m$  for distinct  $X, Y \in F$ ;
- (2)  $\text{card}(X - Y) = m$  for distinct  $X, Y \in G$ ;
- (3)  $\left(\bigcap_{i=1}^s A_i\right) \cap \left(\bigcap_{i=1}^t S - B_i\right) \neq \emptyset$  for distinct  $A_1, \dots, A_s, B_1, \dots, B_t \in H$ ;
- (4)  $X \subseteq Y$  or  $Y \subseteq X$  for each  $X, Y \in L$ .<sup>1</sup>

Following Sierpiński ([2], p. 29), let us call a property  $\mathbf{P}$  *hereditary* if, whenever  $\mathbf{P}$  holds for a set, it also holds for each of its subsets. Now the purpose of this note is to point out that any existence theorem of the above form can be strengthened to read ". . . there exist  $2^{2^m}$  subsets of  $\mathcal{P}(S)$  of cardinality  $2^m$  such that  $\mathbf{P}$ ," provided that  $\mathbf{P}$  is hereditary (as in (1)-(4)). This observation, which does not seem to appear in the literature, is based upon the following

**Theorem.** *For every infinite set of cardinality  $\aleph$  there exists  $2^{\aleph}$  subsets of cardinality  $\aleph$ .*

*Proof.* Let  $S$  be any infinite set of cardinality  $\aleph$ , and put

$$Z = \{X : X \subseteq S \text{ and } \text{card}(X) = \aleph\}$$

---

1. These and similar results can be found in [1], [3] and [4]. In the proof of (4), Wolk [4] employs the generalized continuum hypothesis.