

## A NOTE ON R-MINGLE AND SOBOCIŃSKI'S THREE-VALUED LOGIC

R. ZANE PARKS

The system **R-Mingle (RM)** of Dunn [2] is the result of adding the axiom schema  $A \rightarrow . A \rightarrow A$  to the system **R** of relevant implication—cf. Belnap [1]. Consider the matrices

$\rightarrow$	0	1	2	$\sim$	$\&$	0	1	2	$\vee$	0	1	2
0	2	2	2	2	0	0	0	0	0	0	1	2
*1	0	1	2	1	*1	0	1	1	*1	1	1	2
*2	0	0	2	0	*2	0	1	2	*2	2	2	2

The values 1 and 2 are designated. Since axioms of **RM** always take designated values and *modus ponens* and *adjunction* preserve this property, we have

**Lemma 1.** *Theorems of RM uniformly take designated values when evaluated by the above matrices.*

In [3], Sobociński proved that the system **S** based on the above matrices for  $\rightarrow$  and  $\sim$  is axiomatized by the following schemas together with *modus ponens*:

- S1.  $A \rightarrow B \rightarrow . B \rightarrow C \rightarrow . A \rightarrow C$
- S2.  $A \rightarrow . A \rightarrow B \rightarrow B$
- S3.  $(A \rightarrow . A \rightarrow B) \rightarrow . A \rightarrow B$
- S4.  $A \rightarrow . B \rightarrow . \sim B \rightarrow A$
- S5.  $\sim A \rightarrow \sim B \rightarrow . B \rightarrow A$

Lemma 1, together with Sobociński's result, yields

**Lemma 2.** *If A is a theorem of the pure theory of implication and negation of the calculus RM, then A is a theorem of S.*

In as much as the axioms of **S** are theorems of **RM** (proofs are either routine or given in Dunn [2]) and *modus ponens* is a rule of **RM**, we have

**Lemma 3.** *If A is a theorem of S, then A is a theorem of RM.*