

A SET-THEORETICAL FORMULA EQUIVALENT TO
THE AXIOM OF CHOICE

BOLESŁAW SOBOCIŃSKI

It is obvious that the following set-theoretical formula:¹

S1 For any cardinal numbers m and n which are not finite, if $\aleph(m)$ and $\aleph(n)$ are the least Hartogs' alephs with respect to m and n respectively, and such that $\aleph(m) = \aleph(n)$, then there is no cardinal \wp such that $m < \wp < n$.

is a simple consequence of the theorem:

\aleph . For any cardinal numbers m and n which are not finite, if $\aleph(m)$ and $\aleph(n)$ are the least Hartogs' alephs with respect to m and n respectively, and such that $\aleph(m) = \aleph(n)$, then $m = n$.

which, as it is proved in [3], p. 230, is inferentially equivalent to the axiom of choice. Although at first glance it appears that formula **S1** is weaker than \aleph , in fact, as I shall show in this note, the former formula implies the axiom of choice, and, therefore, it is inferentially equivalent to \aleph . For, a proof is given here that the following theorem:

A. For any cardinal number m which is not finite, if $\aleph(m)$ is the least Hartogs' aleph with respect to m , then there is no cardinal \wp such that $\aleph(m) < \wp < m + \aleph(m)$.

which is inferentially equivalent to the axiom of choice, as it is proved in [2], follows from **S1** without the aid of the said axiom.

Proof: Let us assume **S1** and consider that

(i) m is an arbitrary cardinal number which is not finite,

and that

(ii) $\aleph(m)$ is the least Hartogs' aleph with respect to m .

Then, obviously, we have

Received November 23, 1961