

A NOTE ON UDEs IN AN n -VALUED LOGIC

T. C. WESSELKAMPER

Professor Sobociński has shown that in a 2-valued logic there exists a function with four inputs which is a Universal Decision Element (UDE) and there exists no three-place function with this property [1]. J. C. Muzio has shown recently that in an n -valued logic there is a UDE with $n^2 + n + 1$ inputs. He has conjectured that in an n -valued logic there is a UDE with $n^2 + 1$ inputs [2]. This note exhibits a UDE with $n^2 + 2$ inputs.

Both this note and Muzio's result are obtained by a generalization of the operation called "conditioned disjunction" [3] or "the McCarthy conditional" [4]. Church defines:

$$[x, y, z] = \begin{cases} x, & \text{if } y = \mathbf{T}; \\ z, & \text{if } y = \mathbf{F}. \end{cases}$$

For $[x, y, z]$ McCarthy writes: $(y \rightarrow x, z)$. Note that the generalizations given here do not coincide with McCarthy's generalization to the three-valued case ([4], p. 54).

Let $X(n)$ be the space consisting of the values $0, 1, 2, \dots, n - 1$. Over $X(n)$ define:

$$(A) \quad (x; y_0, y_1, \dots, y_{n-1}) = y_i \text{ if } x = i, 0 \leq i \leq n - 1.$$

Muzio's construction consists of defining:

$$(B) \quad F = (x; c_0, c_1, \dots, c_{n-1}), \text{ where } c_i = (y, z_{i1}, z_{i2}, \dots, z_{i(n-1)}).$$

It is clear that if a two-place function fxy is defined by an n by n operation table of values a_{ij} ($0 \leq i, j \leq n - 1$), then fxy may be obtained from the definition of F by the substitutions:

$$z_{ij} = a_{ij} \quad (0 \leq i, j \leq n - 1).$$

The n repetitions of y in the definition (B) can be eliminated in a simple way. Let $N = n^2$. We define:

$$(C) \quad G = (x, y; z_0, \dots, z_{N-1}) = z_i, \text{ where } i = nx + y.$$

Received October 12, 1973