

A NOTE TO MY PAPER:  
ON CHARACTERIZATIONS OF THE FIRST-ORDER  
FUNCTIONAL CALCULUS

JULIUSZ REICHBACH

In [1] I have presented two characterizations of theses of the first-order functional calculus; the first characterization may be modified in the following way:<sup>1</sup>

- D.0.  $Q(k) \cdot \equiv . Q$  is a non-empty set of tables of the rank  $k$ .
- D.1.  $Q/T, i_1, \dots, i_m \cdot \equiv . (\exists T_1) (\exists j_1) \dots (\exists j_m) \{(T_1 \in Q) \wedge ([T_1|j_1, \dots, j_m] = [T|i_1, \dots, i_m])\}$ .
- $Q/T, i_1, \dots, i_m$  asserts that  $[T|i_1, \dots, i_m]$  is a submodel of some  $T_1 \in Q$  in the meaning of homomorphism.
- D.2.  $T, Q/T_1, i_1, \dots, i_m; i \cdot \equiv . ([T|i_1, \dots, i_m] = [T_1|i_1, \dots, i_m]) \wedge Q/T_1, i_1, \dots, i_m, i$ .
- D.3.  $Q(r, k) \cdot \equiv . (r \leq k) \wedge Q(k) \wedge (i_1 \dots (i_{m+1}) (T) \{(m < r) \wedge (i_1, \dots, i_{m+1} \text{ are different numbers} \leq k) \wedge Q/T, i_1, \dots, i_m \wedge Q/T, i_{m+1} \rightarrow (\exists T_1) (T, Q/T_1, i_1, \dots, i_m; i_{m+1} \wedge (j_1) \dots (j_{m-i}) \{(j_1, \dots, j_s \text{ is a subsequence of } i_1, \dots, i_m) \wedge Q/T, j_1, \dots, j_s, i_{m+1} \rightarrow ([T_1|j_1, \dots, j_s, i_{m+1}])\})\})$ .

The meaning of D.2. and D.3. is clear, see D.1.

For an arbitrary  $T$  of the rank  $k$ , for an arbitrary  $Q$  such that  $Q(k)$  and for an arbitrary formula  $E$  whose indices of free variables occurring in it are  $\leq k$ , we introduce the inductive definition of the functional  $V$ :

- (1d)  $V\{T, Q, f_j^m(x_{r_1}, \dots, x_{r_m})\} = 1 \cdot \equiv . F_j^m(r_1, \dots, r_m),$
- (2d)  $V\{T, Q, F^i\} = 1 \cdot \equiv . \sim V\{T, Q, F\} = 1 \cdot \equiv . V\{T, Q, F\} = 0,$
- (3d)  $V\{T, Q, F + G\} = 1 \cdot \equiv . V\{T, Q, F\} = 1 \vee V\{T, Q, G\} = 1,$
- (4d)  $V\{T, Q, \Pi aF\} = 1 \cdot \equiv . (i) (T_1) \{(i \leq k) \wedge T, Q/T_1, i_1, \dots, i_{w(F)}; i \rightarrow V\{T_1, Q, F(x_i/a)\} = 1\}.$
- D.4.  $E \in P(Q) \cdot \equiv . (T) \{(H \in A \{E\}) \rightarrow Q/T, i_1, \dots, i_{w(H)}\} \rightarrow V\{T, Q, E\} = 1\}.$

*Received April 28, 1961*