

NOTE TO MY PAPER: "A DIAGRAM OF THE FUNCTORS
OF THE TWO-VALUED PROPOSITIONAL CALCULUS"

THOMAS W. SCHARLE

Theorem 6 of my paper [1] gives a criterion for determining which functors of the two-valued propositional calculus are Sheffer functors, viz. that a functor lie in the N quadrant and not be on the vertical axis. From the nature of the diagram presented, it is obvious that precisely one fourth of the n -ary functors lie in the N quadrant, which means that there are $2^{2^{n-2}}$ of them. (Note that, since the I , U , O , and N quadrants in the diagram correspond respectively to the α , β , γ , and δ functors of Post [2], we have a simple formula for the number of such functors)

On the other hand, we can see that the total number of n -ary functors lying on the vertical axis is the square of the number of $(n-1)$ -ary functors on the vertical axis, which yields almost immediately that there are $2^{2^{n-1}}$ of them. Exactly half of these are in the N quadrant, which gives us this equation for the number of n -ary functors satisfying the criterion:

$$(1) \quad S(n) = 2^{2^{n-2}} - 2^{2^{n-1}-1},$$

where $S(n)$ is the number of n -ary Sheffer functors. This formula has also been given without proof in [4].

In a recent paper [3], A. R. Turquette investigates the same problem and gives the answer

$$(2) \quad S(n) = 2^{2^{n-3}} + 2^{2^{n-4}} + 2^{2^{n-5}} + \dots + 2^{2^{n-1}-1}.$$

Having the well known equation for finite sums

$$(3) \quad \sum_{i=1}^n 2^i = 1 - 2^{n+1}$$

we can show the identity of the solutions (1) and (2). For from (2) we have

Received October 20, 1962