

COUNTEREXAMPLES FOR THE SPACE OF MINIMAL
SOLUTIONS OF THE EQUATION $\Delta u = Pu$
ON A RIEMANN SURFACE

Y. K. Kwon and J. L. Schiff

1. Let $P \geq 0$ ($P \neq 0$) be a C^1 -density on an open Riemann surface R . The space of nonnegative C^2 -solutions on R of the elliptic equation $\Delta u = Pu$ is denoted by $PN(R)$. A function $u \in PN(R)$ is said to be PN -minimal if for every $v \in PN(R)$ with $0 \leq v \leq u$, there exists a constant c_v such that $v = c_v u$ on R .

It was established in a former work by the second author [6] that *if the space $PB(R)$ of bounded C^2 -solutions of $\Delta u = Pu$ is of dimension at least 2, then every PN -minimal function on R has zero infimum*. The purpose of this note is to demonstrate that the conclusion is *no longer* valid in the remaining two cases: $\dim PB(R) = 0$ or 1 .

2. First consider the case $\dim PB(R) = 0$. Take R to be the complex plane. It is well-known that $\dim PB(R) = 0$ since R is parabolic (see H. L. Royden [4]). For a constant $M \geq 2$ and the density

$$(1) \quad P(z) = M^2 |z|^{M-2} (1 + |z|^M)^{-1}$$

on R , it is not difficult to see that the function $v(z) = |z|^M + 1$ belongs to the class $PN(R)$.

We claim that $v(z)$ is PN -minimal on R . A bit more strongly, it is true that $PN(R)$ is generated by $v(z)$. For a function $u \in PN(R)$, set $\phi = u \cdot v^{-1}$. We need to show that ϕ is a constant. In view of the conditions $\Delta u = Pu$ and $\Delta v = Pv$, the function ϕ must satisfy the partial differential equation

$$\Delta \phi + \frac{2M |z|^{M-2}}{|z|^M + 1} \left(x \frac{\partial \phi}{\partial x} + y \frac{\partial \phi}{\partial y} \right) = 0$$

on R . It follows from Liouville's theorem (see for example M. Prother and G. Weinberger [3, p. 120]) that every nonnegative solution of this equation is a constant. Thus the plane R with the density (1) carries a PN -minimal function $v(z) = |z|^M + 1$ (≥ 1), although $\dim PB(R) = 0$.

3. Turning now to the case $\dim PB(R) = 1$, we base our argument on the remarkable examples of Y. Tôki [7], [8] (see also L. Sario [5]) of a hyperbolic Riemann surface carrying no nonconstant positive harmonic functions. We take such a surface $R \in O_{HB} - O_G$ and construct a C^1 -density $Q \geq 0$ ($Q \neq 0$) on R such that $\int_R Q(z) dx dy < \infty$. Thus $HB(R)$ and $QB(R)$ are isomorphic (Royden [4], also M. Nakai [1]), which implies that $\dim QB(R) = 1$. Moreover, $\dim HBD(R) = 1$ implies

Received December 11, 1974.

Michigan Math. J. 21 (1974).