

CAPACITY AND MEASURE

Jussi Väisälä

1. *Introduction.* A condenser in the euclidean space R^n is a pair $E = (A, C)$, where A is open in R^n and C is compact in A . For $p \geq 1$, we define the p -capacity of E as

$$\text{cap}_p E = \inf_u \int |\nabla u|^p \, dm,$$

where the infimum is taken over all functions u in $C_0^\infty(A)$ such that $u(x) = 1$ for all $x \in C$. It is well known that if $\text{cap}_p(A_0, C) = 0$ for some bounded A_0 , then $\text{cap}_p(A, C) = 0$ for all open sets A containing C . In this case, we write $\text{cap}_p C = 0$, and otherwise $\text{cap}_p C > 0$. The case $p = n$ is particularly important in the theory of quasiregular maps, and here we write $\text{cap} = \text{cap}_n$. If $p > n$, then $\text{cap}_p C = 0$ only in the case $C = \emptyset$.

The capacity of a condenser can also be defined with the aid of moduli of path families. Given a bounded condenser $E = (A, C)$, we let Γ_E be the family of all paths $\alpha: [a, b] \rightarrow A$ such that $\alpha(a) \in C$ and $\alpha(t) \rightarrow \partial A$ as $t \rightarrow b$. Then

$$\text{cap}_p E = M_p(\Gamma_E),$$

by W. P. Ziemer [10]. Here M_p denotes the p -modulus. Instead of Γ_E , we may take the family of all paths joining C and ∂A in $A \setminus C$.

In this note we shall give a new proof for the following result: If a compact set $C \subset R^n$ has a finite h -measure for $h(r) = (\log(1/r))^{1-n}$, then $\text{cap} C = 0$. The corresponding result holds for the p -capacity with $h(r) = r^{n-p}$.

The earliest result of this type is due to J. W. Lindeberg [4]. He showed that for $n = p = 2$, $\text{cap} C = 0$ for every compact set C of h -measure zero, $h(r) = (\log(1/r))^{-1}$. This result was extended for sets of finite h -measure by P. Erdős and J. Gillis [2]. A simple proof of their result was given by L. Carleson [1]. His proof is also applicable in higher dimensions. These authors used a potential-theoretic definition for capacity. For $p = 2$, this is equivalent to our definition. For $p \neq 2$, this is no longer true, although there are close connections (see [8, p. 332]). Our results are contained in papers of N. Meyers [6, Theorem 21] and, V. G. Mazja and V. P. Havin [5, Section 7], who formulated them in a very general framework. The present formulation is from H. Wallin [9, Theorem 4.3]. For related results, see [8, Remark on p. 335] and [7, Theorem 4.2].

2. *Notation.* If $C \subset R^n$ and $r > 0$, we let $B(C, r)$ be the set of all x in R^n such that $\text{dist}(x, C) < r$. In particular, $B(x, r)$ is the open ball with center at x and radius r . If C is compact, $E(C, r)$ will denote the condenser $(B(C, r), C)$.

3. **LEMMA.** *If $p > 1$ and C is a compact set in R^n with $\text{cap}_p C > 0$, then $\lim_{r \rightarrow 0} \text{cap}_p E(C, r) = \infty$.*

Received January 30, 1975.

Michigan Math. J. 22 (1975).