

CERTAIN ALGEBRAIC FUNCTIONS AND EXTREME POINTS OF S

Louis Brickman

Let S be the usual set of holomorphic, univalent, normalized ($f(0) = 0$, $f'(0) = 1$) functions on the unit disk $\Delta = \{z \in \mathbb{C}: |z| < 1\}$. In [2] it was shown that if $f \in S$ and the set $\mathbb{C} \setminus f(\Delta)$ contains two points of equal modulus, then f is a convex combination of two other members of S . A simple topological argument leads to the further conclusion that if f is an extreme point of S (see [3, p. 439]), then $\mathbb{C} \setminus f(\Delta)$ is an arc tending to infinity with increasing modulus. (Interesting variations of this result are obtained by W. Hengartner and G. Schober in [4].) In the present note we obtain a generalization of the two-point theorem of [2]. In this generalization we assume that $\mathbb{C} \setminus f(\Delta)$ contains a finite set of points of a certain description, and we conclude that f can be written as a nontrivial convex combination of finitely many members of S . In particular, f is not an extreme point of $\overline{\text{co}} S$ (the closure of the convex hull of S). Consequently, the extreme points g of $\overline{\text{co}} S$ have the property that the arc $\mathbb{C} \setminus g(\Delta)$ contains no set E of the type described in the theorem below. (The theorem is applicable because, by [3, p. 440], extreme points of $\overline{\text{co}} S$ must belong to S .)

THEOREM. *Let $P(z) = \prod_{j=1}^n (z - \alpha_j)$, where $n \geq 2$ and where the α_j are distinct complex numbers. Let*

$$Q(z) = \sum_{j=1}^n \frac{\lambda_j P(z)}{z - \alpha_j},$$

where the λ_j are nonzero complex numbers, all having the same argument. Finally, let E be the set of complex numbers w such that $P - wQ$ has a multiple zero. Then E consists of $2n - 2$ points at most, and any $f \in S$ such that $\mathbb{C} \setminus f(\Delta) \supset E$ admits an equation of the form $f = \sum_{j=1}^n t_j f_j$ ($\sum_{j=1}^n t_j = 1$, $t_j > 0$, $f_j \in S$, $f_j \neq f$).

Proof. We begin by noting that Q is a polynomial of degree $n - 1$ and that $Q(\alpha_j) = \lambda_j P'(\alpha_j)$ ($1 \leq j \leq n$), so that $Q = P'$, in the special case where $\lambda_j = 1$ for each j . In particular, we observe that P and Q have no common zeros. Now suppose $w \in E$ and z is a multiple zero of $P - wQ$. Then $P(z) - wQ(z) = 0$, $Q(z) \neq 0$, and $w = P(z)/Q(z)$. Also, $P'(z) - wQ'(z) = 0$, and hence $(QP' - PQ')(z) = 0$. Since $QP' - PQ'$ is a nontrivial polynomial of degree at most $2n - 2$, there are at most $2n - 2$ such numbers z , and since $w = P(z)/Q(z)$, it follows that there are at most $2n - 2$ such w .

If $|w|$ is sufficiently small, $P - wQ$ has distinct zeros $\phi_j(w)$ ($1 \leq j \leq n$) (the branches of the algebraic function defined by the equation $P(z) - wQ(z) = 0$). We number these root functions in the natural way so that $\phi_j(0) = \alpha_j$ ($1 \leq j \leq n$). Each ϕ_j is analytic, and each admits unrestricted analytic continuation in $\mathbb{C} \setminus E$ [1, p. 294]. But $f(\Delta)$ is a simply connected subregion of $\mathbb{C} \setminus E$. Therefore it follows from the monodromy theorem that ϕ_j is analytic and single-valued in $f(\Delta)$ ($1 \leq j \leq n$).

Received June 16, 1975.

Michigan Math. J. 22 (1975).