

ON THE RANK OF AN OPEN MANIFOLD

Chao-Chu Liang

Let M denote a differentiable manifold. J. Milnor defined the rank of M to be the maximal number of vector fields X_1, \dots, X_k on M , everywhere linearly independent, such that their Lie brackets $[X_i, X_j] = 0$ for all i, j . (X_1, \dots, X_k are called *commuting vector fields* [8].) From Hirsch's immersion theorem, we see that any open parallelizable n -dimensional manifold has rank n . By a *tangent k -field* on a manifold M , we mean k linearly independent vector fields X_1, \dots, X_k on M . In this short note, we will prove the following result.

THEOREM. *Let M be an n -dimensional open manifold. Then M admits k linearly independent commuting vector fields if and only if M has a tangent k -field which is homotopic to a foliation.*

Proof. Let $f_M: M \rightarrow BGL_n$ denote the classifying map for the tangent bundle TM of M . If M has a tangent k -field which is homotopic to a foliation, then f_M has a lifting to $BGL_1 \times \dots \times BGL_1 \times B\Gamma_{n-k}$; that is, the lifting is homotopic to $(\text{constant}) \times \dots \times (\text{constant}) \times \phi$, where $\phi: M \rightarrow B\Gamma_{n-k}$. ($B\Gamma_{n-k}$ is Haefliger's classifying space for codimension $n - k$ foliations [1].)

Considering the topological groupoid $V\Gamma_p$ consisting of those local diffeomorphisms of \mathbb{R}^p such that their first derivatives lie in $SL(p, \mathbb{R})$, we may construct a classifying space $BV\Gamma_p$ for the codimension p volume-preserving foliations [7]. (Here we used the fact that an $SL(n, \mathbb{R})$ -structure on an n -dimensional manifold is integrable [3, p. 6].) The argument in [2, p. 148] showed that an open manifold admits a codimension p volume-preserving foliation if and only if the classifying map for its tangent bundle lifts to $BGL_{n-p} \times BV\Gamma_p$ [7].

A *multifoliation* F on M is a collection of foliations $\{F_1, \dots, F_t\}$, with codimension $F_j = k_j$ and TF_j the tangent bundle of F_j , such that

$$\text{codim}(TF_{i_1} \cap \dots \cap TF_{i_s}) = k_{i_1} + \dots + k_{i_s} \quad \text{for any subset } \{i_1, \dots, i_s\} \subseteq \{1, \dots, t\}$$

[5, p. 406]. In [6], we called such an F a *multifoliation of type* (k_1, \dots, k_t) , and showed that an open manifold admits a multifoliation of type (k_1, \dots, k_t) with

$$k = \sum k_j \leq n \text{ if and only if the classifying map for its tangent bundle lifts to } B\Gamma_{k_1} \times \dots \times B\Gamma_{k_t} \times BGL_{n-k}.$$

Our classifying map f_M clearly has a lifting to $BV\Gamma_1 \times \dots \times BV\Gamma_1 \times B\Gamma_{n-k}$. Equivalently, TM can be considered as a normal bundle of a particular codimension n Haefliger Γ -structure [2], where $\Gamma = V\Gamma_1 \times \dots \times V\Gamma_1 \times \Gamma_{n-k}$. Then the argument in the proof of the result mentioned above ([6]) can be used here to show that M admits a multifoliation $F = \{F_1, \dots, F_{k+1}\}$ of type $(1, 1, \dots, 1, n - k)$ such that F_1, \dots, F_k are codimension 1 volume-preserving foliations with respect to the same volume form on M [7].

Received July 19, 1976.

Supported by the University of Kansas General Research Fund.

Michigan Math. J. 23 (1976).