

# AMBIGUOUS POINTS OF A FUNCTION HARMONIC INSIDE A SPHERE

F. Bagemihl

Let  $x, y, z$  denote the Cartesian coordinates of a point in three-dimensional Euclidean space, and set

$$S = \{(x, y, z): x^2 + y^2 + z^2 < 1\}, \quad T = \{(x, y, z): x^2 + y^2 + z^2 = 1\}.$$

**THEOREM.** *There exists a harmonic function  $h(P)$  ( $P \in S$ ) such that, for every  $Q \in T$  and every real number  $r$ , including the values  $+\infty$  and  $-\infty$ , there is a Jordan arc  $J_r^Q$  lying wholly in  $S$  except for its end point  $Q$ , with the property that*

$$\lim_{P \rightarrow Q, P \in J_r^Q} h(P) = r.$$

*Proof.* Piranian has shown [2, Remark 2] that there exists a continuous function  $f(P)$  ( $P \in S$ ) such that the assertion we have made concerning the boundary behavior of  $h(P)$  holds for  $f(P)$ ; he has constructed a tree  $G$  in  $S$  such that, for every  $Q \in T$  and every  $r$ ,  $J_r^Q$  is, except for its end point  $Q$ , a subarc of  $G$ . To define  $h(P)$ , we shall make use of  $G$  and  $f(P)$ .

Let

$$0 < r_0 < r_1 < \dots < r_n < \dots < 1, \quad \lim_{n \rightarrow \infty} r_n = 1,$$

$$S_n = \{(x, y, z): x^2 + y^2 + z^2 < r_n^2\}, \quad T_n = \{(x, y, z): x^2 + y^2 + z^2 = r_n^2\} \quad (n = 0, 1, 2, \dots),$$

$$K_n = (S_n \cup T_n \cup G) \cap (S_{n+1} \cup T_{n+1}) \quad (n = 0, 1, 2, \dots).$$

For every nonnegative integer  $n$ ,  $K_n$  is a compact set with the property that any continuous function on  $K_n$  that is harmonic at every interior point of  $K_n$  can be uniformly approximated on  $K_n$  as closely as desired by a harmonic polynomial (see [1]; I am indebted to Professor J. L. Walsh for this reference).

We define, by induction on  $n$ , a harmonic polynomial  $h_n(P)$ , as follows. Let

$$g_0(P) = 0 \quad (P \in S_0 \cup T_0),$$

$$g_0(P) = f(P) \quad (P \in G \cap T_1),$$

and let  $g_0(P)$  be linear on each segment of  $G$  which extends from  $T_0$  to  $T_1$ . Then  $g_0(P)$  is continuous on  $K_0$  and harmonic at every interior point of  $K_0$ , and hence there exists a harmonic polynomial  $h_0(P)$  for which

$$|h_0(P) - g_0(P)| < 1 \quad (P \in K_0).$$

Suppose that  $n > 0$ , and that we have defined the harmonic polynomial  $h_{n-1}(P)$ . Let