

ON THE MILNOR-SPANIER AND ATIYAH DUALITY THEOREMS

Hans Samelson

The theorems of Milnor and Spanier [5] and of Atiyah [1] state that the Thom spaces of certain pairs of vector bundles over a manifold are (stably) dual to each other in the sense of Spanier and Whitehead. In this note, we give a slight but natural generalization of these theorems, and we relate the result to Poincaré duality. For a different approach, see P. Holm [2].

THEOREM. *Let M be an m -dimensional compact manifold whose boundary ∂M is the union of two manifolds A and B with $\partial A = \partial B = A \cap B$, and suppose that M is embedded in R^m (the same m). Then M/A is m -dual to M/B .*

Here M/A means M with A collapsed to a point; R^m is real m -space. Two (reasonable) spaces are m -dual if they can be embedded (up to homotopy type) disjointly in the sphere S^{m+1} in such a way that for each the inclusion in the complement of the other is a stable homotopy equivalence (it is, in fact, sufficient that this hold for one of them).

The proof goes along the lines of [1] and [5]. There are the inclusions $M \subset R^m = R^m \times 0 \subset R^{m+1}$. Define $T = \{x \in R^{m+1}: x_{m+1} \geq 1\}$ and, as usual, $I = [0, 1] \subset R$. The subset $P = M \times 0 \cup A \times I \cup T$ of R^{m+1} is easily seen to be of the homotopy type of M with a cone erected over A (T is contractible, and the contraction extends to a deformation of P ; thus we may collapse it to a point) and thus of the homotopy type of M/A , since M and A are ANR's. The same is true for the (closed) subset P' of S^{m+1} obtained from P by addition of the point at infinity.

We show next that the complement $Q = S^{m+1} - P' = R^{m+1} - P$ is of the homotopy type of M/B . We write Q as union of two subsets C and D , defined by

$$C = (M - A) \times (0, 1) \quad \text{and} \quad D = (R^{m+1} - (T \cup M \times I)) \cup (B^\circ \times (0, 1));$$

here $(0, 1)$ is the open interval, and B° means the interior $B - \partial B$. One sees easily that (i) D is contractible and the contraction extends to a deformation of Q , so that Q is of the homotopy type of Q/D ; (ii) C is of the homotopy type of M ; and (iii) $Q/D [= C/(B^\circ \times (0, 1))]$ is of the homotopy type of M/B .

For this, we use an exterior collar of ∂M , that is, a homeomorphism f of $\partial M \times I$ onto a neighborhood of ∂M in $R^m - M^\circ$, with $f(y, 0) = y$; also a collar of B in M that over ∂B gives a collar of ∂B in A ; and a similar collar for A . To prove the existence of the exterior collar, we may have to shrink M , using an interior collar (which exists, by M. Brown's theorem). Now for the constructions: The first term of D has an obvious contraction, which begins by pushing down until $x_{m+1} < 0$. To contract all of D , we first pull $B^\circ \times (0, 1)$ out of $M \times I$, by moving all points $(f(y, t), x_{m+1})$ of D in the t -direction, so that the (t, x_{m+1}) -unit square is contracted onto the triangle $x_{m+1} \leq t$.

The open set Q contains a copy, say Q' , of M with a cone erected over B : we shrink $M \times 1/2$ slightly, using the collar over A , and contract B to a point in D . The inclusion $Q' \subset Q$ is a stable homotopy equivalence (one sees easily, *via*

Received December 2, 1968.

This work was partly supported by NSF grant GP-8522.