

ON THE ASYMPTOTIC BEHAVIOR OF FUNCTIONS HOLOMORPHIC IN THE UNIT DISC

Frank B. Ryan

An *asymptotic value* of a function f meromorphic in $D = \{ |z| < 1 \}$ is defined as a limit value α of $f(z)$ as $|z| \rightarrow 1$ on an arc γ in D . In terms of the associated Riemann surface \mathcal{F} over the extended complex w -plane \mathcal{W} , the concept of asymptotic value has the following geometric interpretation: γ is the inverse image of a noncompact arc Γ on \mathcal{F} whose projection into \mathcal{W} ends at the point $w = \alpha$.

A set constitutes the *asymptotic set* of some meromorphic function f (that is, the set of asymptotic values of f) if and only if it is an analytic subset (possibly empty) of \mathcal{W} (see [1], [2]).

The characterization of the asymptotic sets of holomorphic functions is more difficult, because many analytic sets in \mathcal{W} must be excluded (see [4], [5]). The following theorem gives a trivial necessary condition. In the statement of the theorem, ∂G denotes the boundary of G , and the bar $-$ indicates closure. We can easily verify the necessity of the condition by defining G as the image of D under f and using properties of the Riemann surface of f (see [4]).

THEOREM. *If A is the asymptotic set of a function f holomorphic in D , then A is an analytic set and there exists a domain G such that:*

- (1) $\partial G \subset A^- \subset G^-$,
- (2) if $\xi \in \partial G$ is inaccessible from G , then $\xi \notin A$,
- (3) if $\xi \in \partial G - A$, then every arc in G to ξ meets A .

The complexities of the holomorphic case are illustrated by the following example, which shows that the condition in the theorem is not sufficient. At the same time, the example answers a question posed by the author [3]. *There exists an analytic subset A of $\{ |w| < 1 \}$ that meets every arc in $\{ |w| < 1 \}$ ending at a point of $\{ |w| = 1 \}$ but is not the asymptotic set of any function holomorphic in D .*

To construct the example, let S be the finite domain bounded by the triangle with vertices at $(0, 1/4)$, $(0, -1/4)$, and $(1, 0)$. Define

$$C_n = \{ |w| = 1 - 2^{-n} \} \quad \text{and} \quad C_{n,m} = \{ |w| = 1 - 2^{-n} + 2^{-m} \}.$$

Now put

$$A_n = \{ C_n - S \} \cup \bigcup_{m \geq n+2} \{ C_{n,m} \cap S \}$$

and

$$A = \bigcup_{n=1}^{\infty} A_n \cup \{0\}.$$

Received March 23, 1972.

Michigan Math. J. 19 (1972).