

A FIXED-POINT THEOREM FOR HOMOGENEOUS CONTINUA

Charles L. Hagopian

A space is *homogeneous* if for each pair x, y of its points there exists a homeomorphism of the space onto itself that takes x to y . The union S of a finite collection \mathcal{F} of arcs is called a *star* provided that one point of S is the common part of each pair of elements of \mathcal{F} . A continuum X is *star-like* if for each $\varepsilon > 0$ there exists an ε -mapping of X onto a star. Here we prove that every homogeneous star-like continuum has the fixed-point property for homeomorphisms.

A *topological transformation group* (G, X) is a topological group G together with a topological space X and a continuous mapping $(g, x) \rightarrow gx$ of $G \times X$ into X such that $ex = x$ for all $x \in X$ (e denotes the identity of G) and $(gh)x = g(hx)$ for all $g, h \in G$ and $x \in X$.

For each $x \in X$, let G_x be the stabilizer subgroup of x in G (that is, the set of all $g \in G$ such that $gx = x$). If we let G/G_x denote the left-coset space with the usual topology, the mapping of G/G_x onto Gx sending gG_x to gx is one-to-one and continuous. We call the set Gx the *orbit* of x .

Henceforth, X is a continuum (that is, a nondegenerate, compact, connected metric space) and G is the topological group of homeomorphisms of X onto itself with the compact open topology. It follows from a theorem of E. G. Effros [2, Theorem 2.1] that each orbit is a G_δ -set in X if and only if for each $x \in X$, the mapping $gG_x \rightarrow g(x)$ of G/G_x onto Gx is a homeomorphism. In [5], G. S. Ungar pointed out that if X is homogeneous, then $Gx = X$ for each $x \in X$, and therefore $T_x: g \rightarrow g(x)$, being the composition of the natural open mapping of G onto G/G_x and a homeomorphism of G/G_x onto X , is an open mapping of G onto X .

A continuous function f of X onto a space Y is called an ε -mapping if for each $y \in Y$, the diameter of $f^{-1}(y)$ in X is less than ε . A finite sequence $\{L_i\}_{i=1}^n$ of open sets in X is a *chain* provided that $L_i \cap L_j \neq \emptyset$ if and only if $|i - j| \leq 1$.

THEOREM. *Suppose that X is a homogeneous star-like continuum. Then for each homeomorphism h of X onto itself, there exists a point x of X such that $h(x) = x$.*

Proof. Assume there is a homeomorphism h of X onto itself that moves each point of X . There exist a positive number ε and an open set U in G containing h such that for each $f \in U$ and each $x \in X$, the distance from x to $f(x)$ in X is greater than ε .

Since X is star-like, there is a sequence $\{\mathcal{C}_i\}_{i=1}^\infty$ of open covers of X such that for each i , (1) each element of \mathcal{C}_i has diameter less than i^{-1} and (2) there is an element Y_i of \mathcal{C}_i such that $\mathcal{C}_i - \{Y_i\}$ consists of finitely many mutually disjoint chains, each having only one element, an end-link, that meets Y_i . For each i , define y_i to be a point of Y_i .

Let y be a limit point of $\{y_i\}_{i=1}^\infty$, and let $x = h^{-1}(y)$. Note that $T_x[U]$ is an open set in X that contains y . Let j be an integer such that $j^{-1} < \varepsilon/2$ and the closure of Y_j is a subset of $T_x[U]$.

Received October 30, 1974.

Michigan Math. J. 21 (1974).